

MH1101 Calculus II

Tutorial 2 (Week 3) – Questions & Solutions

Academic Year 2025/2026, Semester 2

Quantitative Research Society @NTU

February 20, 2026

Overview of This Tutorial

This tutorial focuses on the Fundamental Theorem of Calculus, definite integrals with absolute values and piecewise functions, substitution techniques, symmetry identities, and average value / mean value theorems for integrals.

- **Q1:** Differentiation of integral-defined functions (FTC Part 1 + Leibniz rule).
- **Q2:** Definite integrals with $|x|$ and piecewise functions.
- **Q3:** Rewriting integrals using an antiderivative function f .
- **Q4:** Substitution practice (six integrals).
- **Q5:** Symmetry identity $\int_0^\pi x f(\sin x) dx$ and a target evaluation.
- **Q6–7:** Average value and weighted-average property across subintervals.

Question 1

Problem

Use the Fundamental Theorem of Calculus (Part 1) to find the derivative of the following functions:

$$(i) \quad g(x) = \int_1^x \cos(t^2) dt.$$

$$(ii) \quad F(s) = \int_s^0 \sqrt{1 + \sec t} dt.$$

$$(iii) \quad R(y) = \int_{2y}^{3y} \frac{u^2 - 1}{u^2 + 1} du.$$

Solution

Method 1: FTC Part 1 + sign handling + Leibniz rule

(i) By FTC Part 1,

$$g'(x) = \cos(x^2).$$

(ii) First rewrite to standard orientation:

$$F(s) = \int_s^0 \sqrt{1 + \sec t} dt = - \int_0^s \sqrt{1 + \sec t} dt.$$

Then by FTC Part 1,

$$F'(s) = -\sqrt{1 + \sec s}.$$

(iii) Use Leibniz rule for variable limits:

$$\frac{d}{dy} \int_{a(y)}^{b(y)} f(u) du = f(b(y))b'(y) - f(a(y))a'(y).$$

Here $f(u) = \frac{u^2-1}{u^2+1}$, $a(y) = 2y$, $b(y) = 3y$. Thus

$$R'(y) = f(3y) \cdot 3 - f(2y) \cdot 2 = 3 \cdot \frac{(3y)^2 - 1}{(3y)^2 + 1} - 2 \cdot \frac{(2y)^2 - 1}{(2y)^2 + 1}.$$

So

$$R'(y) = 3 \cdot \frac{9y^2 - 1}{9y^2 + 1} - 2 \cdot \frac{4y^2 - 1}{4y^2 + 1}.$$

Method 2: Differential-quotient intuition (brief) / consistency check

- (i) Increasing the upper limit x by h adds approximately $\cos(x^2)h$ to the integral, so the derivative is $\cos(x^2)$.
- (ii) Since s appears as a *lower* limit, increasing s reduces the integral by approximately $\sqrt{1 + \sec s} h$, giving $F'(s) = -\sqrt{1 + \sec s}$.
- (iii) A change $y \mapsto y + h$ shifts both bounds; the net effect is approximately “(integrand at top) \times change in top” minus “(integrand at bottom) \times change in bottom”, matching the Leibniz formula above.

Question 2

Problem

Evaluate the following definite integrals:

(i) $\int_{-1}^2 (x - 2|x|) dx.$

(ii) $\int_{-2}^2 f(x) dx$, where

$$f(x) = \begin{cases} x^3, & x \geq 0, \\ -x, & x < 0. \end{cases}$$

Solution

Method 1: Split at sign changes (piecewise evaluation)

(i) Split at $x = 0$ since $|x|$ changes form:

$$\int_{-1}^2 (x - 2|x|) dx = \int_{-1}^0 (x - 2(-x)) dx + \int_0^2 (x - 2x) dx.$$

So

$$\int_{-1}^0 (3x) dx + \int_0^2 (-x) dx = \left[\frac{3x^2}{2} \right]_{-1}^0 + \left[-\frac{x^2}{2} \right]_0^2 = \left(0 - \frac{3}{2} \right) + (-2 - 0) = -\frac{7}{2}.$$

(ii) Split at $x = 0$:

$$\int_{-2}^2 f(x) dx = \int_{-2}^0 (-x) dx + \int_0^2 x^3 dx.$$

Compute:

$$\int_{-2}^0 (-x) dx = \left[-\frac{x^2}{2} \right]_{-2}^0 = 2, \quad \int_0^2 x^3 dx = \left[\frac{x^4}{4} \right]_0^2 = 4.$$

Thus

$$\int_{-2}^2 f(x) dx = 2 + 4 = 6.$$

Method 2: Symmetry + quick checks

(i) On $[0, 2]$, the integrand is $-x$ (negative area), while on $[-1, 0]$ it is $3x$ (also negative). The computed value $-7/2$ is consistent with net negativity.

(ii) On $[-2, 0]$, $f(x) = -x > 0$ contributes positive area; on $[0, 2]$, $f(x) = x^3 > 0$ contributes positive area as well, so a positive result like 6 is plausible.

Question 3

Problem

Express the given indefinite integrals in terms of the function f (assume f is an antiderivative of f'):

(i) $\int f'(5x) dx.$

(ii) $\int x f'(3x^2) dx.$

Solution

Method 1: u -substitution

(i) Let $u = 5x$, so $du = 5 dx$ and $dx = \frac{1}{5} du$:

$$\int f'(5x) dx = \frac{1}{5} \int f'(u) du = \frac{1}{5} f(u) + C = \frac{1}{5} f(5x) + C.$$

(ii) Let $u = 3x^2$, so $du = 6x dx$ and $x dx = \frac{1}{6} du$:

$$\int x f'(3x^2) dx = \frac{1}{6} \int f'(u) du = \frac{1}{6} f(u) + C = \frac{1}{6} f(3x^2) + C.$$

Method 2: Reverse chain rule recognition

(i) Since $\frac{d}{dx} f(5x) = 5f'(5x)$, the integrand is $\frac{1}{5} \frac{d}{dx} f(5x)$.

(ii) Since $\frac{d}{dx} f(3x^2) = f'(3x^2) \cdot 6x$, the integrand $x f'(3x^2)$ is $\frac{1}{6} \frac{d}{dx} f(3x^2)$.

Question 4

Problem

Evaluate the integral by making a suitable substitution.

(i) $\int \frac{x}{\sqrt{1-x^2}} dx.$

(ii) $\int x^2 \sin(x^3) dx.$

(iii) $\int_{-5/2}^{-2} x(2x+5)^8 dx.$

(iv) $\int \frac{\sin \sqrt{x}}{\sqrt{x}} dx.$

(v) $\int \frac{1}{\cos^2 t \sqrt{1+\tan t}} dt.$

(vi) $\int_{\pi/3}^{2\pi/3} \csc^2\left(\frac{t}{2}\right) dt.$

Solution

Method 1: Standard substitutions (show main steps)

(i) Let $u = 1 - x^2$, so $du = -2x dx$:

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\frac{1}{2} \int u^{-1/2} du = -u^{1/2} + C = -\sqrt{1-x^2} + C.$$

Equivalently,

$$-\sqrt{1-x^2} + C = -\frac{1}{3}(1-x^2)^{3/2} \text{ up to constant,}$$

and one common antiderivative form is

$$-\frac{1}{3}(1-x^2)^{3/2} + C',$$

since $\frac{d}{dx}(1-x^2)^{3/2} = -3x\sqrt{1-x^2}$. Hence

$$\int \frac{x}{\sqrt{1-x^2}} dx = -\sqrt{1-x^2} + C \quad (\text{simplest form}).$$

(ii) Let $u = x^3$, $du = 3x^2 dx$:

$$\int x^2 \sin(x^3) dx = \frac{1}{3} \int \sin u du = -\frac{1}{3} \cos u + C = -\frac{1}{3} \cos(x^3) + C.$$

(iii) Let $u = 2x + 5$, so $du = 2 dx$ and $x = \frac{u-5}{2}$:

$$\int_{-5/2}^{-2} x(2x+5)^8 dx = \int_0^1 \frac{u-5}{2} u^8 \cdot \frac{1}{2} du = \frac{1}{4} \int_0^1 (u^9 - 5u^8) du.$$

Compute:

$$\frac{1}{4} \left[\frac{u^{10}}{10} - 5 \frac{u^9}{9} \right]_0^1 = \frac{1}{4} \left(\frac{1}{10} - \frac{5}{9} \right) = \frac{1}{4} \cdot \frac{9-50}{90} = -\frac{41}{360}.$$

(iv) Let $u = \sqrt{x}$, so $x = u^2$, $dx = 2u du$:

$$\int \frac{\sin \sqrt{x}}{\sqrt{x}} dx = \int \frac{\sin u}{u} \cdot 2u du = 2 \int \sin u du = -2 \cos u + C = -2 \cos(\sqrt{x}) + C.$$

(v) Let $u = 1 + \tan t$, so $du = \sec^2 t dt = \frac{1}{\cos^2 t} dt$:

$$\int \frac{1}{\cos^2 t \sqrt{1 + \tan t}} dt = \int u^{-1/2} du = 2u^{1/2} + C = 2\sqrt{1 + \tan t} + C.$$

(vi) Let $u = \frac{t}{2}$, so $dt = 2 du$. When $t = \pi/3$, $u = \pi/6$; when $t = 2\pi/3$, $u = \pi/3$:

$$\int_{\pi/3}^{2\pi/3} \csc^2\left(\frac{t}{2}\right) dt = 2 \int_{\pi/6}^{\pi/3} \csc^2(u) du = 2 [-\cot u]_{\pi/6}^{\pi/3} = 2 \left(-\cot \frac{\pi}{3} + \cot \frac{\pi}{6} \right).$$

Since $\cot(\pi/3) = \frac{1}{\sqrt{3}}$ and $\cot(\pi/6) = \sqrt{3}$,

$$= 2 \left(-\frac{1}{\sqrt{3}} + \sqrt{3} \right) = 2 \cdot \frac{2}{\sqrt{3}} = \frac{4}{\sqrt{3}} = \frac{4\sqrt{3}}{3}.$$

Method 2: Reverse chain rule / derivative matching (brief)

Each part follows from recognizing the integrand as (a constant multiple of) the derivative of a composite:

- (i) derivative of $\sqrt{1-x^2}$.
- (ii) derivative of $\cos(x^3)$.
- (iii) substitution $u = 2x + 5$ converts to a polynomial integral on $[0, 1]$.
- (iv) substitution $u = \sqrt{x}$ removes the \sqrt{x} from denominator.
- (v) substitution $u = 1 + \tan t$ uses $du = \sec^2 t dt$.
- (vi) substitution $u = t/2$ turns the integral into $\int \csc^2 u du$.

Question 5

Problem

If f is continuous on $[0, 1]$, show that

$$\int_0^\pi x f(\sin x) dx = \frac{\pi}{2} \int_0^\pi f(\sin x) dx.$$

Hint: Substitute $u = \pi - x$.

Hence, evaluate the integral

$$\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx.$$

You may assume that $\int \frac{1}{1+x^2} dx = \tan^{-1} x + C$.

Solution

Method 1: Symmetry substitution $u = \pi - x$

Let

$$I = \int_0^\pi x f(\sin x) dx.$$

Substitute $u = \pi - x$, so $x = \pi - u$, $dx = -du$, and $\sin x = \sin(\pi - u) = \sin u$:

$$I = \int_\pi^0 (\pi - u) f(\sin u) (-du) = \int_0^\pi (\pi - u) f(\sin u) du.$$

Rename u back to x :

$$I = \int_0^\pi (\pi - x) f(\sin x) dx = \pi \int_0^\pi f(\sin x) dx - \int_0^\pi x f(\sin x) dx = \pi \int_0^\pi f(\sin x) dx - I.$$

Hence

$$2I = \pi \int_0^\pi f(\sin x) dx \quad \Rightarrow \quad I = \frac{\pi}{2} \int_0^\pi f(\sin x) dx.$$

For the target integral, take

$$f(\sin x) = \frac{\sin x}{1 + \cos^2 x}.$$

Then

$$\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx = \frac{\pi}{2} \int_0^\pi \frac{\sin x}{1 + \cos^2 x} dx.$$

Evaluate the remaining integral with $u = \cos x$, $du = -\sin x dx$:

$$\int_0^\pi \frac{\sin x}{1 + \cos^2 x} dx = \int_{u=1}^{u=-1} \frac{-1}{1 + u^2} du = \int_{-1}^1 \frac{1}{1 + u^2} du = [\arctan u]_{-1}^1 = \frac{\pi}{4} - \left(-\frac{\pi}{4}\right) = \frac{\pi}{2}.$$

Therefore

$$\int_0^\pi \frac{x \sin x}{1 + \cos^2 x} dx = \frac{\pi}{2} \cdot \frac{\pi}{2} = \frac{\pi^2}{4}.$$

Method 2: Pairing x and $\pi - x$ directly

Write

$$\int_0^\pi x f(\sin x) dx = \frac{1}{2} \int_0^\pi \left(x f(\sin x) + (\pi - x) f(\sin(\pi - x)) \right) dx.$$

Since $\sin(\pi - x) = \sin x$, the integrand becomes

$$\frac{1}{2} \int_0^\pi (x + (\pi - x)) f(\sin x) dx = \frac{\pi}{2} \int_0^\pi f(\sin x) dx,$$

giving the same identity, and thus the same evaluation $\pi^2/4$ for the “hence” integral.

Question 6

Problem

Let $f(x) = \frac{1}{\sqrt{x}}$.

- (i) Find the average value of f on the interval $[1, 4]$.
- (ii) Find the value of c guaranteed by the Mean Value Theorem for Integrals such that $f_{\text{ave}} = f(c)$.

Solution

Method 1: Average value formula + solve $f(c) = f_{\text{ave}}$

(i)

$$f_{\text{ave}} = \frac{1}{4-1} \int_1^4 x^{-1/2} dx = \frac{1}{3} [2x^{1/2}]_1^4 = \frac{1}{3} \cdot 2(2-1) = \frac{2}{3}.$$

(ii) Solve $f(c) = \frac{2}{3}$:

$$\frac{1}{\sqrt{c}} = \frac{2}{3} \quad \Rightarrow \quad \sqrt{c} = \frac{3}{2} \quad \Rightarrow \quad c = \frac{9}{4}.$$

Method 2: Geometry/monotonicity check

Since $f(x) = 1/\sqrt{x}$ is continuous and decreasing on $[1, 4]$, the average value must lie between $f(4) = 1/2$ and $f(1) = 1$, and $\frac{2}{3}$ satisfies this. The equation $f(c) = 2/3$ has a unique solution in $(1, 4)$, namely $c = 9/4$.

Question 7

Problem

If $f_{\text{ave}}[a, b]$ denotes the average value of f on $[a, b]$ and $a < c < b$, show that

$$f_{\text{ave}}[a, b] = \left(\frac{c-a}{b-a} \right) f_{\text{ave}}[a, c] + \left(\frac{b-c}{b-a} \right) f_{\text{ave}}[c, b].$$

Solution

Method 1: Expand using the definition of average value

By definition,

$$f_{\text{ave}}[a, b] = \frac{1}{b-a} \int_a^b f(x) dx.$$

Split the integral at c :

$$\int_a^b f(x) dx = \int_a^c f(x) dx + \int_c^b f(x) dx.$$

Hence

$$f_{\text{ave}}[a, b] = \frac{1}{b-a} \int_a^c f(x) dx + \frac{1}{b-a} \int_c^b f(x) dx.$$

Now rewrite each term in terms of averages:

$$\frac{1}{b-a} \int_a^c f = \frac{c-a}{b-a} \cdot \frac{1}{c-a} \int_a^c f = \frac{c-a}{b-a} f_{\text{ave}}[a, c],$$

$$\frac{1}{b-a} \int_c^b f = \frac{b-c}{b-a} \cdot \frac{1}{b-c} \int_c^b f = \frac{b-c}{b-a} f_{\text{ave}}[c, b].$$

Combine to obtain the required identity.

Method 2: Weighted-average interpretation

The total area under f on $[a, b]$ is the sum of areas on $[a, c]$ and $[c, b]$. Dividing by the total width $(b-a)$, the overall “height” (average value) is a convex combination of the subinterval averages, weighted by their relative lengths $(c-a)$ and $(b-c)$, giving exactly the stated formula.