

MH2100 Calculus III

Midterm I (Solutions)

AY 2021/2022, Semester 1

Quantitative Research Society @NTU

September 2021

Overview

- **Coverage:** parametric curves and tangent lines; limits, continuity and partial derivatives; directional derivatives, tangent planes, the Chain Rule, implicit differentiation, and critical points.
- **Format:** three questions, 50 marks in total, 90 minutes (official paper: September 2021).
- **Historical permitted material:** the September 2021 paper permitted one two-sided A4 piece of paper with handwritten notes. Consult current course materials for present assessment rules.
- **Source:** MH2100 Midterm I, AY2021/2022 Semester 1. Questions are transcribed from the official paper; solutions and marking schemes are unofficial QRS@NTU material.

Instructions (verbatim)

1. DO NOT TURN OVER PAPER UNTIL INSTRUCTED.
2. This midterm paper contains THREE (3) questions.
3. Answer ALL questions. The marks for each question are indicated at the beginning of each question.
4. Candidates can write anywhere on this midterm paper.
5. ONE two-sided A4 piece of paper with handwritten notes is permitted.

Contents

Question 1	(17 Marks)	3
Solution		3
Method 1: Direct computation		3
Unofficial QRS marking scheme (17 marks)		4
Question 2	(16 Marks)	5
Solution		5
Method 1: Direct limits and definitions		5
Unofficial QRS marking scheme (16 marks)		6
Question 3	(17 Marks)	7
Solution		7
Method 1: Direct computation		7
Unofficial QRS marking scheme (17 marks)		8

Question 1**(17 Marks)****QUESTION 1.**

- (i) [3 marks] Find parametric equations for the line that is tangent to the curve parametrised by

$$\mathbf{r}(t) = \begin{pmatrix} 5 \sin t \\ t^3 \\ 4e^{3t} \end{pmatrix},$$

at the point $(0, 0, 4)$.

- (ii) [3 marks] The cartesian equation $y = ax + b$ describes the line tangent to the curve

$$\begin{cases} x(t) = \sin(2t), \\ y(t) = t \cos t \end{cases}$$

at the point defined by $t = \pi/2$. Find the values of a and b .

- (iii) [3 marks] At the point defined by $t = a$ the tangent vector of the curve parametrised by

$$\mathbf{r}(t) = \begin{pmatrix} e^{\sin t} \tan t \\ t^2 - 10t \\ 23 \end{pmatrix},$$

is orthogonal to the vector $\langle 0, 1, 2 \rangle$. Find the value of a .

- (iv) [3 marks] Find the directional derivative of the function

$$f(x, y, z) = x^3 + y^2 - 4z^2 + z \ln x$$

in the direction of the vector $\langle 1, 4, 1 \rangle$.

- (v) [5 marks] In which directions is the directional derivative of

$$f(x, y) = x^2y - y^8 + 2x$$

at $(6, 1)$ equal to zero? Write the unit vector for each direction.

Solution**Method 1: Direct computation**

- (i) $t^3 = 0 \implies t = 0$.

$$\mathbf{r}'(t) = \langle 5 \cos t, 3t^2, 12e^{3t} \rangle \implies \mathbf{r}'(0) = \langle 5, 0, 12 \rangle$$

$$\boxed{\mathbf{l}(s) = \langle 0, 0, 4 \rangle + s\langle 5, 0, 12 \rangle, \quad s \in \mathbb{R},} \quad \boxed{x = 5s, \quad y = 0, \quad z = 4 + 12s.}$$

- (ii)

$$(x, y)|_{t=\pi/2} = \left(\sin \pi, \frac{\pi}{2} \cos \frac{\pi}{2} \right) = (0, 0)$$

$$\frac{dy}{dx} = \frac{\cos t - t \sin t}{2 \cos 2t} \Big|_{t=\pi/2} = \frac{-\pi/2}{-2} = \frac{\pi}{4}$$

Line through the origin: $\boxed{a = \frac{\pi}{4}, \quad b = 0.}$

(iii)

$$\mathbf{r}'(t) \cdot \langle 0, 1, 2 \rangle = (2t - 10) + 2(0) = 2t - 10 = 0 \text{ at } t = a \implies \boxed{a = 5.}$$

$$(iv) \quad \mathbf{u} = \langle 1, 4, 1 \rangle / \sqrt{18}, \quad \nabla f = \left\langle 3x^2 + \frac{z}{x}, 2y, -8z + \ln x \right\rangle.$$

$$D_{\mathbf{u}}f = \nabla f \cdot \mathbf{u} \implies \boxed{D_{\mathbf{u}}f(x, y, z) = \frac{3x^2 + z/x + 8y - 8z + \ln x}{\sqrt{18}} \quad (x > 0).}$$

(v)

$$\begin{aligned} \nabla f &= \langle 2xy + 2, x^2 - 8y^7 \rangle \implies \nabla f(6, 1) = \langle 14, 28 \rangle = 14\langle 1, 2 \rangle \\ D_{\mathbf{u}}f(6, 1) = 0 &\iff u_1 + 2u_2 = 0, \quad u_1^2 + u_2^2 = 1 \\ &\implies \mathbf{u} = (-2u_2, u_2), \quad 5u_2^2 = 1 \implies u_2 = \pm \frac{1}{\sqrt{5}} \end{aligned}$$

$$\boxed{\mathbf{u} = \frac{1}{\sqrt{5}}\langle 2, -1 \rangle \quad \text{or} \quad \mathbf{u} = \frac{1}{\sqrt{5}}\langle -2, 1 \rangle.}$$

Unofficial QRS marking scheme (17 marks)

- (i) **(3 marks)** 1: identifies $t = 0$; 1: obtains $\mathbf{r}'(0) = \langle 5, 0, 12 \rangle$; 1: gives the line through $(0, 0, 4)$.
- (ii) **(3 marks)** 1: finds the point $(0, 0)$; 1: computes slope $\pi/4$; 1: gives $a = \pi/4, b = 0$.
- (iii) **(3 marks)** 1: differentiates the relevant components; 1: imposes orthogonality; 1: obtains $a = 5$.
- (iv) **(3 marks)** 1: computes the gradient; 1: normalizes the direction; 1: takes the correct dot product.
- (v) **(5 marks)** 2: computes $\nabla f(6, 1) = \langle 14, 28 \rangle$; 1: states orthogonality; 1: obtains one correct unit direction; 1: obtains the opposite direction.

Question 2**(16 Marks)****QUESTION 2.**

- (i) [8 marks] Evaluate each of the following limits or show that the limit does not exist. Justify your answers.

(a)

$$\lim_{(x,y) \rightarrow (5,0)} \frac{5x^2y + 10}{\sqrt{x^2 + y^2}}$$

(b)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{3y}{\sqrt{x^2 + y^2}}$$

(c)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2 + x^2y + xy^3 + 3y^2}{x^2 + y^2}$$

- (ii) [8 marks] Define

$$f(x, y) = \begin{cases} \frac{x^3y + y^3}{x^6 + y^2}; & \text{if } x^2 + y^2 \neq 0, \\ 0; & \text{if } x^2 + y^2 = 0. \end{cases}$$

- (a) Is f continuous at $(0, 0)$? Justify your answer.
 (b) Find $f_x(0, 0)$ and $f_y(0, 0)$.
 (c) Is f differentiable at $(0, 0)$? Justify your answer.

Solution**Method 1: Direct limits and definitions**

- (i) (a) Denominator $\rightarrow 5 \neq 0$ (quotient continuous at $(5, 0)$), so substitute:

$$\lim_{(x,y) \rightarrow (5,0)} \frac{5x^2y + 10}{\sqrt{x^2 + y^2}} = \frac{10}{5} = 2.$$

(b)

$$y = 0 : 0; \quad x = 0 : \frac{3y}{|y|} \rightarrow 3 \ (y \rightarrow 0^+), \rightarrow -3 \ (y \rightarrow 0^-)$$

the limit does not exist.

(c)

$$\begin{aligned} \frac{3x^2 + x^2y + xy^3 + 3y^2}{x^2 + y^2} &= 3 + \frac{x^2y}{x^2 + y^2} + \frac{xy^3}{x^2 + y^2} \\ \left| \frac{x^2y}{x^2 + y^2} \right| &\leq |y| \rightarrow 0, \quad \left| \frac{xy^3}{x^2 + y^2} \right| \leq \frac{r^4}{r^2} = r^2 \rightarrow 0 \quad (r = \sqrt{x^2 + y^2}) \\ &\implies \text{both} \rightarrow 0 \quad (\text{Squeeze Theorem}) \end{aligned}$$

$$\lim_{(x,y) \rightarrow (0,0)} \frac{3x^2 + x^2y + xy^3 + 3y^2}{x^2 + y^2} = 3.$$

(ii) (a)

$$x = 0 : f(0, y) = \frac{y^3}{y^2} = y \rightarrow 0$$

$$y = x^3 : f(x, x^3) = \frac{x^6 + x^9}{2x^6} = \frac{1 + x^3}{2} \rightarrow \frac{1}{2}$$

Path limits differ:

$$\boxed{f \text{ is not continuous at } (0, 0).}$$

(b)

$$f_x(0, 0) = \lim_{h \rightarrow 0} \frac{f(h, 0) - 0}{h} = 0$$

$$f_y(0, 0) = \lim_{k \rightarrow 0} \frac{f(0, k) - 0}{k} = \lim_{k \rightarrow 0} \frac{k}{k} = 1$$

$$\boxed{f_x(0, 0) = 0, \quad f_y(0, 0) = 1.}$$

(c) Differentiable $\implies$ continuous; (a) gives f not continuous at $(0, 0)$.

$$\boxed{f \text{ is not differentiable at } (0, 0).}$$

Unofficial QRS marking scheme (16 marks)(i) **(8 marks)**

- (a) **(2 marks)** 1: continuity/substitution justification; 1: final value 2.
- (b) **(3 marks)** 1: computes the $y = 0$ path; 1: uses $x = 0$ to obtain distinct one-sided values; 1: concludes nonexistence.
- (c) **(3 marks)** 1: separates the constant term; 1: bounds both remaining terms; 1: concludes 3.

(ii) **(8 marks)**

- (a) **(3 marks)** 1: path $x = 0$; 1: path $y = x^3$; 1: concludes non-continuity.
- (b) **(3 marks)** 1: correct difference quotient for f_x ; 1: correct difference quotient for f_y ; 1: states both values.
- (c) **(2 marks)**
- 1 mark: invokes differentiability implies continuity.
 - 1 mark: concludes non-differentiability.

Question 3**(17 Marks)****QUESTION 3.**

- (i) [3 marks] Find the directional derivative of the function

$$f(x, y) = \begin{cases} \frac{xy + y^2}{x^2 + y^2}; & \text{if } x^2 + y^2 \neq 0, \\ 0; & \text{if } x^2 + y^2 = 0, \end{cases}$$

in the direction of the vector $\langle 3, 4 \rangle$ at the point $(0, 0)$.

- (ii) [3 marks] Find an equation for the tangent plane to the surface given by the equation
- $z = 5x^2 - 10y^2$
- at the point
- $(2, 1, 10)$
- .

- (iii) [3 marks] Use the Chain Rule to find
- $\frac{\partial z}{\partial u}$
- when
- $u = 0$
- and
- $v = 2$
- for

$$z = \sin(xy) + \sin(y), \quad x = 2u^2 + 3v^2, \quad y = 2uv.$$

- (iv) [3 marks] Find
- $\frac{\partial z}{\partial y}$
- at the point
- $(\pi, 2\pi, \pi/2)$
- where
- z
- is defined implicitly by

$$3 \cos(x + 2y) + 3 \cos(x + 2z) + \cos(2y + z) = 0.$$

- (v) [5 marks] Let

$$f(x, y) = 27x^3 - 81x + 2y^2 + 8y - 17.$$

Find and classify all critical points of f . (For each critical point, determine whether or not it is a local max/min or a saddle point.)**Solution****Method 1: Direct computation**

- (i)
- $\mathbf{u} = \langle \frac{3}{5}, \frac{4}{5} \rangle$
- . For
- $t \neq 0$
- :

$$f(t\mathbf{u}) = \frac{(3t/5)(4t/5) + (4t/5)^2}{(3t/5)^2 + (4t/5)^2} = \frac{28}{25}$$

$$\frac{f(t\mathbf{u}) - f(0, 0)}{t} = \frac{28}{25t} \text{ is unbounded as } t \rightarrow 0$$

$$\boxed{D_{\mathbf{u}}f(0, 0) \text{ does not exist.}}$$

- (ii)
- $g = 5x^2 - 10y^2$
- :
- $g_x = 10x$
- ,
- $g_y = -20y \implies g_x(2, 1) = 20$
- ,
- $g_y(2, 1) = -20$
- .

$$z - 10 = 20(x - 2) - 20(y - 1) \implies \boxed{20x - 20y - z = 10.}$$

- (iii)

$$\frac{\partial z}{\partial u} = z_x x_u + z_y y_u, \quad z_x = y \cos(xy), \quad z_y = x \cos(xy) + \cos y, \quad x_u = 4u, \quad y_u = 2v$$

$$(u, v) = (0, 2) \implies (x, y) = (12, 0), \quad z_x = 0, \quad z_y = 12 + 1 = 13, \quad x_u = 0, \quad y_u = 4$$

$$\boxed{\left. \frac{\partial z}{\partial u} \right|_{(u,v)=(0,2)} = 0 \cdot 0 + 13 \cdot 4 = 52.}$$

(iv) $F = 3 \cos(x + 2y) + 3 \cos(x + 2z) + \cos(2y + z), \quad F_y + F_z z_y = 0.$

$$F_y = -6 \sin(x + 2y) - 2 \sin(2y + z)$$

$$F_z = -6 \sin(x + 2z) - \sin(2y + z)$$

At $(\pi, 2\pi, \pi/2)$: $\sin 5\pi = 0, \sin 2\pi = 0, \sin \frac{9\pi}{2} = 1 \implies F_y = -2, F_z = -1 \neq 0.$

$$\boxed{\frac{\partial z}{\partial y} = -\frac{F_y}{F_z} = -2.}$$

(v)

$$f_x = 81(x^2 - 1) = 0, \quad f_y = 4y + 8 = 0 \implies (1, -2), (-1, -2)$$

$$f_{xx} = 162x, \quad f_{xy} = 0, \quad f_{yy} = 4 \implies D = f_{xx}f_{yy} - f_{xy}^2 = 648x$$

$$(1, -2) : D = 648 > 0, \quad f_{xx} = 162 > 0 \implies \text{local min}$$

$$(-1, -2) : D = -648 < 0 \implies \text{saddle}$$

$$\boxed{(1, -2) \text{ is a local minimum; } (-1, -2) \text{ is a saddle point.}}$$

Unofficial QRS marking scheme (17 marks)

- (i) **(3 marks)** 1: normalizes the direction; 1: evaluates the function along that ray; 1: concludes that the directional derivative does not exist.
- (ii) **(3 marks)** 1: finds both partial derivatives; 1: evaluates the slopes at $(2, 1)$; 1: gives the correct plane.
- (iii) **(3 marks)** 1: writes the Chain Rule; 1: evaluates the intermediate variables and derivatives; 1: obtains 52.
- (iv) **(3 marks)** 1: forms F_y and F_z ; 1: evaluates them at the stated point; 1: obtains $z_y = -2$.
- (v) **(5 marks)**
 - 2 marks: computes and solves $f_x = f_y = 0$.
 - 1 mark: computes the Hessian entries and discriminant.
 - 1 mark: classifies $(1, -2)$.
 - 1 mark: classifies $(-1, -2)$.