

MH2500 PROBABILITY AND INTRODUCTION TO STATISTICS
NANYANG TECHNOLOGICAL UNIVERSITY

Review Questions with solutions

Part 3

These are review questions for continuous variable (Chapters 5, 6 in Ross' book).

.....

- (1) For some constant c , the random variable X has the probability density function

$$f(x) = \begin{cases} cx^4 & 0 < x < 2; \\ 0 & \text{otherwise.} \end{cases}$$

Find (a) $\mathbb{E}(X)$ and (b) $\text{Var}(X)$.

Solution: By $\int_0^2 cx^4 dx = 1$, we have $c = \frac{5}{32}$.

$$\mathbb{E}(X) = \int_0^2 x \left(\frac{5}{32} x^4 \right) dx = \left(\frac{5}{32} \right) \left(\frac{64}{6} \right) = \frac{5}{3}.$$

$$\mathbb{E}(X^2) = \int_0^2 x^2 \left(\frac{5}{32} x^4 \right) dx = \left(\frac{5}{32} \right) \left(\frac{128}{7} \right) = \frac{27}{7}.$$

This gives

$$\text{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}(X))^2 = \frac{27}{7} - \left(\frac{5}{3} \right)^2 = \frac{5}{63}.$$

- (2) The random variable X has the probability density function

$$f(x) = \begin{cases} ax + bx^2 & 0 < x < 1; \\ 0 & \text{otherwise.} \end{cases}$$

If $\mathbb{E}(X) = 0.6$, find (a) $P\{X < \frac{1}{2}\}$, and (b) $\text{Var}(X)$.

Solution: By $\int_0^1 (ax + bx^2) dx = 1$, we have $1 = \frac{1}{2}a + \frac{1}{3}b$. By

$$\mathbb{E}(X) = \int_0^1 x(ax + bx^2) dx = \frac{1}{3}a + \frac{1}{4}b,$$

we have

$$\frac{1}{3}a + \frac{1}{4}b = 0.6.$$

This gives $a = 3.6, b = -2.4$.

This gives

$$(a) P\{X < 1/2\} = \int_0^{1/2} (3.6x - 2.4x^2) dx = [1.8x^2 - 0.8x^3]_0^{1/2} = 0.35.$$

$$(b) \mathbb{E}(X^2) = \int_0^1 x^2 (3.6x - 2.4x^2) dx = 0.42. \text{ This gives}$$

$$\text{Var}(X) = \mathbb{E}(X^2) - (\mathbb{E}(X))^2 = 0.42 - 0.36 = 0.06.$$

- (3) The annual rainfall in Cleveland, Ohio is approximately a normal random variable with mean 40.2 inches and standard deviation 8.4 inches. What is the probability that

- (a) next year's rainfall will exceed 44 inches?
 (b) the yearly rainfalls in exactly 3 of the next 7 years will exceed 44 inches?

Assume that if A_i is the event that the rainfall exceeds 44 inches in year i (from now), then the events A_i , $i \geq 1$, are independent.

Solution: Let X be next year's rainfall and let

$$Z = (X - 40.2)/8.4.$$

- (a) The probability of next year's rainfall will exceed 44 inches is $P\{X > 44\} = P\{Z > 3.8/8.4\} \approx P\{Z > 0.4524\} \approx 0.3255$.
- (b) The probability of yearly rainfalls in exactly 3 of the next 7 years will exceed 44 inches is $\binom{7}{3}(0.3255)^3(0.6745)^4$.
- (4) A roulette wheel has 38 slots, numbered 0, 00, and 1 through 36. If you bet 1 on a specified number then you either win 35 if the roulette ball lands on that number or lose 1 if it does not. If you continually make such bets, approximate the probability that
- you are winning after 34 bets;
 - you are winning after 1000 bets;
 - you are winning after 100,000 bets.

Assume that each roll of the roulette ball is equally likely to land on any of the 38 numbers.

Solution: Let X denote the number of the first n bets that you win, then the amount that you will be winning after n bets is

$$35X - (n - X) = 36X - n.$$

Thus, we want to determine $p = P\{36X - n > 0\} = P\{X > n/36\}$ when X is a binomial random variable with parameters n and $p = 1/38$.

(a) $n = 34$:

$$\begin{aligned} p &= P\{X \geq 1\} \\ &= P\{X \geq 0.5\} \quad (\text{continuity correction}) \\ &= P\left\{\frac{X - \frac{34}{38}}{\sqrt{34(\frac{1}{38})(\frac{37}{38})}} \geq \frac{0.5 - \frac{34}{38}}{\sqrt{34(\frac{1}{38})(\frac{37}{38})}}\right\} \\ &= P\{Z \geq -0.4229\} \approx \Phi(0.4229) \approx 0.6638. \end{aligned}$$

(b) $n = 1000$:

$$\begin{aligned} p &= P\{X \geq 28\} \\ &= P\{X \geq 27.5\} \quad (\text{continuity correction}) \\ &= P\left\{\frac{X - \frac{1000}{38}}{\sqrt{1000(\frac{1}{38})(\frac{37}{38})}} \geq \frac{27.5 - \frac{1000}{38}}{\sqrt{1000(\frac{1}{38})(\frac{37}{38})}}\right\} \\ &\approx 1 - \Phi(0.2339) \approx 0.4075. \end{aligned}$$

(c) $n = 100,000$:

$$\begin{aligned} p &= P\{X \geq 2778\} \\ &= P\{X \geq 2777.5\} \quad (\text{continuity correction}) \\ &= P\left\{\frac{X - \frac{100000}{38}}{\sqrt{100000(\frac{1}{38})(\frac{37}{38})}} \geq \frac{2777.5 - \frac{100000}{38}}{\sqrt{100000(\frac{1}{38})(\frac{37}{38})}}\right\} \\ &\approx 1 - \Phi(0.2883) \approx 0.0020. \end{aligned}$$

- (5) Let X be the number of times that a fair coin that is flipped 40 times lands on heads. Find the probability that $X = 20$, by using the normal approximation.

Solution:

$$\begin{aligned} P\{X = 20\} &= P\{19.5 \leq X \leq 20.5\} \quad (\text{continuity correction}) \\ &= P\left\{\frac{19.5 - 20}{\sqrt{10}} \leq \frac{X - 20}{\sqrt{10}} \leq \frac{20.5 - 20}{\sqrt{10}}\right\} \\ &\approx P\{-0.16 \leq Z \leq 0.16\} \approx \Phi(0.16) - \Phi(-0.16) \approx 0.1272. \end{aligned}$$

- (6) To determine the effectiveness of a certain diet in reducing the amount of cholesterol in the bloodstream, 100 people are put on the diet. After they have been on the diet for a sufficient length of time, their cholesterol count will be taken. The nutritionist running this experiment has decided to endorse the diet if at least 65 percent of the people have a lower cholesterol count after going on the diet. What is the probability that the nutritionist endorses the new diet if, in fact, it has no effect on the cholesterol level?

Solution: Let us assume that if the diet has no effect on the cholesterol count, then, strictly by chance, each person's count will be lower than it was before the diet with probability $1/2$. Hence, if X is the number of people whose count is lowered, then the probability that the nutritionist will endorse the diet when it actually has no effect on the cholesterol count is

$$\begin{aligned} \sum_{i=65}^{100} \binom{100}{i} \left(\frac{1}{2}\right)^{100} &\approx P\{X \geq 64.5\} \quad (\text{continuity correction}) \\ &= P\left\{\frac{X - 100 \times \frac{1}{2}}{\sqrt{100 \times \frac{1}{2} \times \frac{1}{2}}} \geq \frac{64.5 - 100 \times \frac{1}{2}}{\sqrt{100 \times \frac{1}{2} \times \frac{1}{2}}}\right\} \\ &\approx P\{Z \geq 2.9\} = 1 - \Phi(2.9) \approx 0.0019. \end{aligned}$$

- (7) Suppose that the number of miles that a car can run before its battery wears out is exponentially distributed with an average value of 10,000 miles. If a person desires to take a 5000-mile trip, what is the probability that he or she will be able to complete the trip without having to replace the car battery? What can be said when the distribution is not exponential?

Solution: It follows by the memoryless property of the exponential distribution that the remaining lifetime (in thousands of miles) of the battery is exponential with parameter $\lambda = \frac{1}{10}$. Hence, the desired probability is

$$P\{\text{remaining lifetime} > 5\} = 1 - F(5) = e^{-5\lambda} = e^{-1/2} \approx 0.604.$$

If the lifetime distribution F is not exponential, then the relevant probability is

$$P\{\text{lifetime} > t + 5 \mid \text{lifetime} > t\} = \frac{1 - F(t + 5)}{1 - F(t)},$$

where t is the number of miles that the battery had been in use prior to the start of the trip.

Conclusion: if the distribution is not exponential, additional information is needed (namely, the value of t) before the desired probability can be calculated.

(8) The joint density function of X and Y is given by

$$f(x, y) = C(y - x)e^{-y}, \quad -y < x < y, 0 < y < \infty$$

and is equal to 0 outside the region.

- (a) Find C .
- (b) Find the density function of X .
- (c) Find the density function of Y .
- (d) Find $\mathbb{E}(X)$ and $\mathbb{E}(Y)$.

Solution:

(a) By

$$1 = \int_0^\infty \int_{-y}^y (C(y - x)e^{-y}) dx dy = C \int_0^\infty e^{-y} \left[\int_{-y}^y (y - x) dx \right] dy = C \int_0^\infty (e^{-y}(2y^2)) dy = 4C,$$

and we have $C = 1/4$.

We can use $\int_0^\infty (e^{-y}y^n) dy = n!$. *Prove it.*

- (b) We need to make sure that for $x > 0$, the joint density is nonzero only when $y > x$, and for $x < 0$, the joint density is nonzero only when $y > -x$. Thus, for $x > 0$, the density function of X is

$$\begin{aligned} f_X(x) &= \frac{1}{4} \int_x^\infty (y - x)e^{-y} dy \\ &= \frac{1}{4} \int_0^\infty ue^{-(x+u)} du \\ &= \frac{1}{4} e^{-x} \end{aligned}$$

and for $x < 0$, the density function of X is

$$\begin{aligned} f_X(x) &= \frac{1}{4} \int_{-x}^\infty (y - x)e^{-y} dy \\ &= \frac{1}{4} [-ye^{-y} - e^{-y} + xe^{-y}]_{-x}^\infty \\ &= \frac{1}{4} e^x (-2x + 1). \end{aligned}$$

- (c) The density function of Y is $f_Y(y) = \frac{1}{4} e^{-y} \int_{-y}^y ((y - x)e^{-y}) dx = \frac{1}{2} y^2 e^{-y}$.

(d)

$$\begin{aligned} \mathbb{E}(X) &= \frac{1}{4} \left[\int_0^\infty xe^{-x} dx + \int_{-\infty}^0 xe^x (-2x + 1) dx \right] \\ &= \frac{1}{4} \left[1 - \int_0^\infty (2y^2 e^{-y} + ye^{-y}) dy \right] \\ &= \frac{1}{4} [1 - 4 - 1] = -1 \\ \mathbb{E}(Y) &= \int_0^\infty y \left(\frac{1}{2} y^2 e^{-y} \right) dy = 3. \end{aligned}$$

(9) The joint density function of X and Y is given by

$$f(x, y) = xy, \quad 0 < x < 1, 0 < y < 2$$

and is equal to 0 outside the region.

- (a) Are X and Y independent?
- (b) Find the density function of X .
- (c) Find the density function of Y .

- (d) Find the joint distribution function.
- (e) Find $\mathbb{E}(Y)$.
- (f) Find $P\{X + Y > 1\}$.

Solution:

- (a) Yes. X and Y are independent. The joint density function factors as x and y .
- (b) For $0 < x < 1$, $f_X(x) = \int_0^2 (xy)dy = 2x$, and $f_X(x) = 0$ for $x \notin (0, 1)$.
- (c) For $0 < y < 2$, $f_Y(y) = \int_0^1 (xy)dx = y$, and $f_Y(y) = 0$ for $y \notin (0, 2)$.
- (d) $P\{X < x, Y < y\} = P\{X < x\}P\{Y < y\}$, where, for $x > 0$, $P\{X < x\} = \min\{1, x^2\}$ and for $y > 0$, $P\{Y < y\} = \min\{1, y^2/4\}$.
- (e) $\mathbb{E}(Y) = \int_0^2 (y \cdot y/2) dy = 4/3$.
- (f)

$$\begin{aligned} P\{X + Y > 1\} &= \int_0^1 \int_{1-x}^2 xy dy dx = \int_0^1 x \left(\int_{1-x}^2 y dy \right) dx = \frac{1}{2} \int_0^1 x (4 - (1-x)^2) dx \\ &= \frac{1}{2} \int_0^1 (3x + 2x^2 - x^3) dx = \frac{1}{2} (3/2 + 2/3 - 1/4) = 23/24. \end{aligned}$$

- (10) Let N be a geometric random variable with parameter p . Suppose that the conditional distribution of X given that $N = n$ is the gamma distribution with parameters n and λ . Find the conditional probability mass function of N given that $X = x$.

Solution:

$$\begin{aligned} P\{N = n \mid X = x\} &= f_{X|N}(x|n)P\{N = n\}/f_X(x) \\ &= C \frac{1}{(n-1)!} (\lambda x)^{n-1} (1-p)^{n-1} = C (\lambda(1-p)x)^{n-1} / (n-1)!. \end{aligned}$$

This shows that, conditional on $X = x$, $N - 1$ is a Poisson random variable with mean $\lambda(1-p)x$. That is, for $n \geq 1$,

$$\begin{aligned} P\{N = n \mid X = x\} &= P\{N - 1 = n - 1 \mid X = x\} \\ &= e^{-\lambda(1-p)x} (\lambda(1-p)x)^{n-1} / (n-1)!. \end{aligned}$$