

MH2100 Calculus III

Midterm I (Solutions)

AY 2023/2024, Semester 1

Quantitative Research Society @NTU

September 2023

Overview

- **Coverage:** parametric curves and tangent lines; limits and continuity; partial and directional derivatives; tangent planes, the Chain Rule, implicit differentiation, and critical points.
- **Historical source:** MH2100 Midterm I, AY2023/2024 Semester 1. Every question below is transcribed from the official paper. Solutions and marking schemes are unofficial QRS@NTU material.
- **Current administration (AY2026/2027):** Tuesday 6 October 2026, 16:30–18:20 (110 minutes), venue TBA. One double-sided typed A4 sheet is permitted; attachments and sticky notes are not.

Instructions to candidates (historical, verbatim)

1. DO NOT TURN OVER PAPER UNTIL INSTRUCTED.
2. This midterm paper contains THREE (3) questions.
3. Answer ALL questions. The marks for each question are indicated at the beginning of each question.
4. Candidates can write anywhere on this midterm paper.
5. ONE two-sided A4 piece of paper with handwritten notes is permitted.

Contents

Question 1	(17 Marks)	3
Solution		3
Method 1: Direct calculation		3
UNOFFICIAL QRS marking scheme (17 marks)		4
Question 2	(16 Marks)	5
Solution		5
Method 1: Direct calculation		5
UNOFFICIAL QRS marking scheme (16 marks)		6
Question 3	(17 Marks)	7
Solution		7
Method 1: Direct calculation		7
UNOFFICIAL QRS marking scheme (17 marks)		8

Question 1**(17 Marks)****QUESTION 1.**

- (i) [3 marks] Find parametric equations for the tangent line at the point when $t = 1$ to the curve with parametric equations

$$x = 3t^2, \quad y = 5t, \quad z = 7t^3.$$

- (ii) [3 marks] The cartesian equation $y = ax + b$ describes the line tangent to the curve

$$\begin{cases} x(t) = e^t, \\ y(t) = t^2 + 2t - 7 \end{cases}$$

at the point defined by $t = 0$. Find the values of a and b .

- (iii) [3 marks] Find all points (P, Q) on the following curve where the tangent line is vertical.

$$x = t^3 - 9t + 18, \quad y = \sin(t)$$

- (iv) [3 marks] At the point $(0, \pi, 1)$, find the directional derivative of the function $f(x, y, z) = \sin(xyz)$ in the direction of the vector $\langle 1, 2, 2 \rangle$.
- (v) [5 marks] Find all unit direction vectors $\mathbf{u} = \langle a, b \rangle$ in which the directional derivative of the function $f(x, y) = ye^{-xy}$ at the point $(0, 2)$ has the value 1.

Solution**Method 1: Direct calculation**

(i)

$$\begin{aligned} \mathbf{r}(1) &= \langle 3, 5, 7 \rangle, & \mathbf{r}'(t) &= \langle 6t, 5, 21t^2 \rangle \\ \mathbf{r}'(1) &= \langle 6, 5, 21 \rangle \implies \boxed{\mathbf{l}(s) = \langle 3, 5, 7 \rangle + s\langle 6, 5, 21 \rangle, \quad s \in \mathbb{R}.} \end{aligned}$$

(ii)

$$\begin{aligned} (x, y)|_{t=0} &= (1, -7), & a &= \frac{dy}{dx} = \frac{y'}{x'} = \frac{2t+2}{e^t} \Big|_{t=0} = 2 \\ -7 &= 2(1) + b \implies \boxed{a = 2, \quad b = -9.} \end{aligned}$$

(iii)

$$\begin{aligned} x'(t) = 3t^2 - 9 = 0 &\implies t = \pm\sqrt{3} & (x' = 0, y' \neq 0) \\ y'(\pm\sqrt{3}) = \cos(\pm\sqrt{3}) \neq 0 & & (\frac{\pi}{2} < \sqrt{3} < \frac{3\pi}{2}) \\ \boxed{(P, Q) = (18 - 6\sqrt{3}, \sin \sqrt{3}) \quad \text{or} \quad (18 + 6\sqrt{3}, -\sin \sqrt{3}).} \end{aligned}$$

(iv)

$$\begin{aligned} \nabla f &= \cos(xyz) \langle yz, xz, xy \rangle \implies \nabla f(0, \pi, 1) = \langle \pi, 0, 0 \rangle \\ \boxed{D_{\mathbf{u}}f(0, \pi, 1) &= \langle \pi, 0, 0 \rangle \cdot \frac{\langle 1, 2, 2 \rangle}{3} = \frac{\pi}{3}.} & (|\langle 1, 2, 2 \rangle| = 3) \end{aligned}$$

(v)

$$\begin{aligned}\nabla f &= e^{-xy} \langle -y^2, 1 - xy \rangle \implies \nabla f(0, 2) = \langle -4, 1 \rangle \\ \nabla f \cdot \mathbf{u} &= -4a + b = 1, \quad a^2 + b^2 = 1 \\ a^2 + (1 + 4a)^2 &= 1 \implies a(17a + 8) = 0 \quad (b = 1 + 4a)\end{aligned}$$

$$\mathbf{u} = \langle 0, 1 \rangle \quad \text{or} \quad \mathbf{u} = \left\langle -\frac{8}{17}, -\frac{15}{17} \right\rangle.$$

UNOFFICIAL QRS marking scheme (17 marks)

- (i) **(3 marks)** 1: point at $t = 1$; 1: tangent vector; 1: correct line.
- (ii) **(3 marks)** 1: point; 1: slope by dy/dx ; 1: both a and b .
- (iii) **(3 marks)** 1: vertical-tangent condition; 1: parameters and nonzero y' check; 1: both points.
- (iv) **(3 marks)** 1: gradient; 1: normalised direction; 1: dot product.
- (v) **(5 marks)** 1: partial derivatives; 1: gradient at the point; 1: directional-derivative equation; 1: unit-vector equation; 1: both vectors.

Question 2**(16 Marks)****QUESTION 2.**

- (i) [4 marks] Evaluate each of the following limits or show that the limit does not exist. Justify your answers.

(a)

$$\lim_{(x,y) \rightarrow (1,0)} \frac{\sin(xy)}{x^3y + (x-y)^4}$$

(b)

$$\lim_{(x,y) \rightarrow (0,0)} \frac{x^3y}{x^6 + y^2}$$

- (ii) [5 marks] Find all points (P, Q, R) of the level surface given by

$$x^3 + 3x^2y + 6yz - 3z^2 = 7$$

where the tangent plane has an equation of the form $Ax + Ay = B$ for some constants A and B .

- (iii) [7 marks] Define

$$f(x, y) = \begin{cases} \frac{y^3}{x^2 + y^2}; & \text{if } x^2 + y^2 \neq 0, \\ 0; & \text{if } x^2 + y^2 = 0. \end{cases}$$

- (a) Is f continuous at $(0, 0)$? Justify your answer.
 (b) Find $f_x(0, 0)$ and $f_y(0, 0)$.
 (c) Is f differentiable at $(0, 0)$? Justify your answer.

Solution**Method 1: Direct calculation**

- (i) (a)

$$\frac{\sin(xy)}{x^3y + (x-y)^4} \rightarrow \frac{\sin 0}{0+1} = \boxed{0}. \quad (\text{denominator} \rightarrow 1 \neq 0)$$

(b)

$$y = 0 : 0, \quad y = x^3 : \frac{x^6}{x^6 + x^6} = \frac{1}{2} \ (x \neq 0)$$

$$\therefore \boxed{\text{the limit does not exist.}}$$

- (ii)

$$F := x^3 + 3x^2y + 6yz - 3z^2$$

$$\nabla F = \langle 3x^2 + 6xy, 3x^2 + 6z, 6y - 6z \rangle \parallel \langle 1, 1, 0 \rangle \quad (Ax + Ay = B)$$

$$6y - 6z = 0 \implies z = y$$

$$3x^2 + 6xy = 3x^2 + 6y \implies y(x-1) = 0$$

$$y = z = 0 : x^3 = 7 \quad (F = 7)$$

$$x = 1, z = y : 1 + 3y + 3y^2 = 7 \implies y = 1 \text{ or } -2$$

$$\nabla F \neq \mathbf{0} : \text{planes } x + y = \sqrt[3]{7}, 2, -1$$

$$(P, Q, R) = (\sqrt[3]{7}, 0, 0), (1, 1, 1), \text{ or } (1, -2, -2).$$

(iii) (a)

$$|f(x, y)| = \frac{|y|^3}{x^2 + y^2} \leq |y| \rightarrow 0 = f(0, 0) \quad (\text{Squeeze})$$

$$f \text{ is continuous at } (0, 0).$$

(b)

$$\frac{f(h, 0)}{h} = 0, \quad \frac{f(0, h)}{h} = 1 \quad (h \neq 0)$$

$$f_x(0, 0) = 0, \quad f_y(0, 0) = 1.$$

(c)

$$L(x, y) = f_x(0, 0)x + f_y(0, 0)y = y \quad (\text{if differentiable})$$

$$\frac{f(t, t) - L(t, t)}{\sqrt{t^2 + t^2}} = \frac{t/2 - t}{\sqrt{2}|t|} = -\frac{t}{2\sqrt{2}|t|} \not\rightarrow 0$$

$$f \text{ is not differentiable at } (0, 0).$$

UNOFFICIAL QRS marking scheme (16 marks)

(i) (4 marks)

(a) (2 marks) 1: nonzero limiting denominator; 1: limit 0.

(b) (2 marks) 1: two valid paths; 1: nonexistence conclusion.

(ii) (5 marks) 1: gradient; 1: plane-normal conditions; 1: branch $y = 0$; 1: branch $x = 1$; 1: all three points.

(iii) (7 marks)

(a) (2 marks) 1: valid bound; 1: continuity conclusion.

(b) (2 marks) 1: f_x ; 1: f_y .

(c) (3 marks) 1: candidate derivative; 1: diagonal-path remainder; 1: non-differentiability conclusion.

Question 3**(17 Marks)****QUESTION 3.**

- (i) [3 marks] Find the second order partial derivative
- w_{xy}
- where

$$w = y \sin(3x^2 - 5y).$$

- (ii) [3 marks] Use the Chain Rule to find
- $\frac{\partial u}{\partial y}$
- at
- $(x, y, z) = (-1, 0, 1)$
- for the function

$$u(p, q, r) = \frac{p^2}{q + r},$$

where

$$p = x + 2y, \quad q = y + 2z, \quad r = x + y + z.$$

- (iii) [3 marks] Assume that
- z
- is uniquely determined in terms of
- x
- and
- y
- by the equation

$$z^3 - xy + yz + y^3 = 2.$$

Find the value of $\frac{\partial z}{\partial y}$ at the point $(1, 1, 1)$.

- (iv) [3 marks] In which directions is the directional derivative of

$$f(x, y) = \sin(xy) - x^2 + xy$$

at $(\pi, 0)$ equal to zero? Write the unit vector for each direction.

- (v) [5 marks] Let

$$f(x, y) = x^3 - y^2 + xy - \frac{x}{2}.$$

- (a) Find all critical points of
- $f(x, y)$
- .

- (b) For each critical point found in Part (a), determine whether or not it is a local max/min or a saddle point.

Solution**Method 1: Direct calculation**

- (i)

$$w_x = 6xy \cos(3x^2 - 5y)$$

$$\boxed{w_{xy} = 6x \cos(3x^2 - 5y) + 30xy \sin(3x^2 - 5y).} \quad (\text{product rule})$$

- (ii)

$$(p, q, r) = (-1, 2, 0), \quad (p_y, q_y, r_y) = (2, 1, 1)$$

$$u_p = \frac{2p}{q+r} = -1, \quad u_q = u_r = -\frac{p^2}{(q+r)^2} = -\frac{1}{4}$$

$$\frac{\partial u}{\partial y} = u_p p_y + u_q q_y + u_r r_y = -2 - \frac{1}{4} - \frac{1}{4} = \boxed{-\frac{5}{2}}$$

(iii)

$$3z^2 z_y - x + z + yz_y + 3y^2 = 0 \quad (\partial_y)$$

$$z_y = \frac{x - z - 3y^2}{3z^2 + y} = \frac{1 - 1 - 3}{3 + 1}$$

$$\boxed{\frac{\partial z}{\partial y} = -\frac{3}{4}.}$$

(iv)

$$f_x = y \cos(xy) - 2x + y, \quad f_y = x \cos(xy) + x$$

$$\nabla f(\pi, 0) = 2\pi \langle -1, 1 \rangle$$

$$\nabla f \cdot \langle a, b \rangle = 0 \iff -a + b = 0, \quad a^2 + b^2 = 1$$

$$\boxed{\mathbf{u} = \frac{1}{\sqrt{2}} \langle 1, 1 \rangle \quad \text{or} \quad \mathbf{u} = \frac{1}{\sqrt{2}} \langle -1, -1 \rangle.}$$

(v) (a)

$$f_x = 3x^2 + y - \frac{1}{2} = 0, \quad f_y = -2y + x = 0$$

$$y = \frac{x}{2} \implies 6x^2 + x - 1 = 0$$

$$\boxed{\left(\frac{1}{3}, \frac{1}{6}\right) \quad \text{and} \quad \left(-\frac{1}{2}, -\frac{1}{4}\right).}$$

(b)

$$f_{xx} = 6x, \quad f_{xy} = 1, \quad f_{yy} = -2$$

$$D = f_{xx}f_{yy} - f_{xy}^2 = -12x - 1$$

(Second Derivative Test)

$$\left(\frac{1}{3}, \frac{1}{6}\right): D = -5 < 0 \implies \text{saddle}$$

$$\left(-\frac{1}{2}, -\frac{1}{4}\right): D = 5 > 0, \quad f_{xx} = -3 < 0 \implies \text{local max}$$

$$\boxed{\left(\frac{1}{3}, \frac{1}{6}\right) \text{ saddle; } \left(-\frac{1}{2}, -\frac{1}{4}\right) \text{ local maximum.}}$$

UNOFFICIAL QRS marking scheme (17 marks)

- (i) **(3 marks)** 1: first partial; 1: product and Chain Rules; 1: correct w_{xy} .
- (ii) **(3 marks)** 1: Chain Rule; 1: intermediate values and derivatives; 1: value $-5/2$.
- (iii) **(3 marks)** 1: implicit differentiation; 1: isolates z_y ; 1: evaluates at the point.
- (iv) **(3 marks)** 1: gradient; 1: orthogonality equation; 1: both unit vectors.
- (v) **(5 marks)**
 - (a) **(2 marks)** 1: solves the first-derivative equations; 1: both critical points.
 - (b) **(3 marks)** 1: Hessian discriminant; 1: saddle classification; 1: maximum classification.