

Rising temperatures alter how and where mammals move

Stefano Mezzini^{1,2} Christen H. Fleming^{3,4} Karen E. Hodges^{1,2}

Siobhan Darlington^{1,2} Adam T. Ford^{1,2} TJ Gooliaff⁵

Kirk Safford⁶ Robert Serrouya^{1,2,7} Michael J. Noonan^{1,2,8}

¹ Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

² Department of Biology, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

³ Department of Biology, University of Central Florida, Orlando, Florida 32816, United States.

⁴ Smithsonian Conservation Biology Institute, National Zoological Park, 1500 Remount Rd., Front Royal, VA 22630, United States.

⁵ British Columbia Ministry of Forests, Lands, Natural Resource Operations and Rural Development, Penticton, BC, Canada.

⁶ British Columbia Ministry of Environment and Parks, Penticton, BC, Canada.

⁷ Wildlife Science Centre, Biodiversity Pathways, University of British Columbia Okanagan, Revelstoke, British Columbia, Canada.

⁸ Department of Computer Science, Math, Physics, and Statistics, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

Running Title: Temperature alters mammal movement behaviour

Article type: Research article

Words in abstract: 291

Words in main text: 7789

Figures: 9

Tables: 3

References: 155 (updated on 2025-10-01)

Appendices: 2

Key words: climate change, temperature, mammals, animal movement, movement behaviour, habitat selection, projections

Contact Information: *Corresponding author's telephone, and email details*

Abstract

Widespread warming during the last century has caused many terrestrial mammals to change how and where they move, with cascading effects on fitness and community dynamics. Previous studies have estimated the effects of temperature on mammalian movement behaviour, but few disentangled them from seasonal behaviour cycles. Consequently, it is still uncertain how mammals will adapt their movement behaviour throughout the 21st century. We address this gap by quantifying the effects of temperature on the movement rates and habitat selection of six large-bodied mammalian species (boreal and southern mountain caribou, cougars, elk, grizzly bears, mountain goats, and wolves) throughout western Canada between 1998 and 2023. We show that temperature significantly affected how and where these mammals moved, and we predict that climate change will impact mammals' movement behaviour throughout the 21st century. Projected responses to climate change suggested that rising temperatures will drive southern mountain caribou and mountain goats to move more, while cougars, elk, and wolves will move less. Boreal caribou and grizzly bears showed little change in projected yearly movement rates but clear shifts in phenology. We also predict that rising temperatures will reduce median habitat selection strength for four of the species, although cougars and elk are expected to show increased selection strength for higher altitudes. As mammals increasingly alter their movement rates and select against portions of their current ranges, changes in individuals' movement behaviour will impact encounter rates, including predator-prey dynamics and human-wildlife interactions. Conservation efforts should therefore account for future changes in movement behaviour as well as the consequences such changes may have on communities. Anticipating changes in mammalian movement behaviour will become crucial for effectively and proactively understanding community-level responses and selecting high-quality habitat for long-term conservation.

¹ 1 Introduction

² For most animals, movement is a rapid and low-cost response to a variety of stimuli (Nathan
³ *et al.*, 2008). Animals move to optimize resource use (e.g., food, water, shelter; Charnov,
⁴ 1976; Kacelnik *et al.*, 1992; Merkle *et al.*, 2016; Mezzini *et al.*, 2025), optimize habitat use
⁵ (Winter *et al.*, 2024), and avoid predators and competitors (Tórrez-Herrera *et al.*, 2020; Pe-
⁶ terson *et al.*, 2021; Bartashevich *et al.*, 2024; Tan *et al.*, 2024). How and where animals move
⁷ thus also determines encounter rates with resources (Mezzini *et al.*, 2025), humans (Johnson
⁸ *et al.*, 2018; Weststrate *et al.*, 2024), predators, prey, and competitors (Martinez-Garcia *et*
⁹ *al.*, 2020; Glass *et al.*, 2021; Brivio *et al.*, 2024). However, ambient temperature affects
¹⁰ animals' movement rates by altering the energetic cost of movement and thermoregulation
¹¹ (McNab, 1970; Taylor *et al.*, 1982; Brown *et al.*, 2004; Fuller *et al.*, 2016; Jahn & Seebacher,
¹² 2022), and extreme temperatures can impact animals' movement rates (Dyer *et al.*, 2023),
¹³ phenology (McLellan & McLellan, 2015; Pigeon *et al.*, 2016; Johnson *et al.*, 2018; Carbeck
¹⁴ *et al.*, 2022), and life history (Newediuk *et al.*, 2024), and even cause physiological damage
¹⁵ and death (Hetem *et al.*, 2014; Powers *et al.*, 2017; Ratnayake *et al.*, 2019; Schmidt *et al.*,
¹⁶ 2020; Schwerdt *et al.*, 2024). Animals may search for a short-term buffer from extreme
¹⁷ temperatures by moving to thermal refugia (Hannah *et al.*, 2014; Elmore *et al.*, 2017; Attias
¹⁸ *et al.*, 2018; Arechavala-Lopez *et al.*, 2019; Gulland *et al.*, 2022), and shelter may even be
¹⁹ preferred over forage abundance (Hall *et al.*, 2016). However, climate change during the last
²⁰ century (IPCC, 2023) has caused animals to alter how and where they move as they respond
²¹ to hotter temperatures and increasingly severe and frequent extreme-heat events (Bunnell
²² *et al.*, 2011; Thompson *et al.*, 2022; IPCC, 2023).

²³ Mammals are particularly susceptible to adverse effects from excessive heat (Sherwood
²⁴ & Huber, 2010). While extreme cold often leads to reduced movement, provided that in-
²⁵ dividuals can find refuge and take advantage of their body heat (Berger *et al.*, 2018; Hou
²⁶ *et al.*, 2020; Glass *et al.*, 2021), extreme heat can often result in more severe physiological

27 damage in a shorter span of time (Jessen, 2001; Sherwood & Huber, 2010; Mota-Rojas *et*
28 *al.*, 2021; Newediuk *et al.*, 2024). Mammals' homeothermy can cause them to have difficulty
29 dissipating heat, particularly when ambient temperatures are near or above body tempera-
30 ture. Larger-bodied mammals are more likely to alter their movement behaviour with rising
31 temperatures, since they tend to move long distances (Merkle *et al.*, 2016), are particularly
32 sensitive to overheating (Fuller *et al.*, 2016; Alston *et al.*, 2020; Verzuh *et al.*, 2023), and
33 their movement rates are thus strongly limited by excessive heat (Dyer *et al.*, 2023).

34 Global warming during the last century has resulted in milder and shorter winters, hotter
35 and longer summers, and a greater risk of extreme-heat events (IPCC, 2023), with widespread
36 changes in mammals' phenologies, circadian rhythms, fitness, and life histories (Botero *et al.*,
37 2015; McLellan & McLellan, 2015; Pigeon *et al.*, 2016; Wells *et al.*, 2022; Newediuk *et al.*,
38 2024). However, behavioural responses to heat vary substantially across species depending
39 on body size, vagility, and activity times (McCain & King, 2014; Williams & Blois, 2018).
40 For example, warming temperatures have allowed deer (*Odocoileus* spp.) to shift northward,
41 which has led to increased density of wolves (*Canis lupus*) and higher predation on caribou
42 (*Rangifer tarandus caribou*) (Latham *et al.*, 2011; Barber *et al.*, 2018; Dickie *et al.*, 2024).
43 Preparing for and responding to future changes will require a detailed understanding of
44 not only how and where mammals move throughout the thermal landscape but also
45 the cascade of effects that such changes will have at the population and community levels
46 (Cunningham *et al.*, 2021).

47 As heat stress intensifies throughout the 21st century, mammals will continue to suffer
48 impacts on their fitness, movement rates, and habitat selection (Deb *et al.*, 2020; Woo-
49 Durand *et al.*, 2020). It remains unclear how or whether species will be able to respond to
50 climate change in the current century (Deb *et al.*, 2020; Woo-Durand *et al.*, 2020; Verzuh
51 *et al.*, 2023), especially if populations fail to adapt (Botero *et al.*, 2015; Sawyer *et al.*, 2019)
52 or are physiologically incapable to do so (Sherwood & Huber, 2010; Botero *et al.*, 2015;
53 Williams & Blois, 2018). Recent work has documented the effects of climate change on

54 mammals' ranges (Leclerc *et al.*, 2021), thermoregulation (Mota-Rojas *et al.*, 2021), and
55 movement behaviour (McCain, 2019; Cunningham *et al.*, 2021; Melin *et al.*, 2023), but there
56 remains a need for fine-scale estimates of the effects of temperature on mammals' movement
57 behaviour and quantitative predictions of future changes in movement behaviour.

58 In this paper, we quantify the effects of proximal air temperature on mammalian move-
59 ment behaviour, namely movement rates and habitat selection of six large-bodied terrestrial
60 mammal species in western Canada (boreal and southern mountain caribou, cougars: *Puma*
61 *concolor*, elk: *Cervus canadensis*, grizzly bears: *Ursus arctos horribilis*, mountain goats:
62 *Oreamnos americanus*, and wolves; Table 1). Western Canada is currently experiencing
63 accelerating and widespread but spatially heterogeneous warming (Turner & Clifton, 2009;
64 Kienzle, 2018; Dierauer *et al.*, 2021), phenological shifts (plants: Post & Forchhammer,
65 2008; Tysor, 2025; ice: Basu *et al.*, 2024), and more frequent and intense extreme heat
66 events (Zhang *et al.*, 2023). Decision makers must understand which areas will best sustain
67 and protect biodiversity not only for current conditions but future decades, too. Failing to
68 understand how climate change will affect mammalian movement behaviour will hinder our
69 ability to respond to the current climate crisis and make decisions that are viable in the
70 long term. Furthermore, understanding how temperature affects individual species is the
71 first step towards understanding how temperature impacts the community dynamics (Hegel
72 *et al.*, 2010; Cunningham *et al.*, 2021; Carbeck *et al.*, 2022) and phenology (Bunnell *et al.*,
73 2011; Duncan *et al.*, 2012; Merkle *et al.*, 2016; Slatyer *et al.*, 2022; Gerlich *et al.*, 2025), and
74 how these changes will impact mammals in the coming decades (Sobie *et al.*, 2021; Eum *et*
75 *al.*, 2023; Weststrate *et al.*, 2024).

Table 1: Summary statistics of each species' set of GPS data, namely: the start and end of each set of GPS telemetry data, the number of GPS fixes (after data cleaning), the median sampling interval (Δt ; stratified by animal), the number of animals (Animals), and the number of animals with finite speed estimates (Has speed).

Species	Start	End	Fixes	Δt (hours)	Animals	Has speed
Caribou (boreal)	2011-03-02	2018-01-03	200,561	13.00	146	131

Species	Start	End	Fixes	Δt (hours)	Animals	Has speed
Caribou (southern mountain)	1998-03-20	2009-06-06	27,921	6.00	22	20
Cougars	2006-02-04	2021-07-12	80,650	2.00	29	29
Elk	2007-01-13	2013-11-19	875,853	2.00	169	169
Grizzly bears	2004-09-29	2009-09-07	39,021	1.00	18	18
Mountain goats	2019-06-24	2023-10-04	65,452	6.25	11	10
Wolves	2013-01-26	2017-08-29	202,433	0.25	39	39

76 2 Methods

77 2.1 Datasets used

78 In this study, we estimate how temperature affected mammals' probability of moving,
 79 speed when moving, and habitat selection by leveraging three datasets: (1) a multi-species
 80 collection of 25 years of GPS telemetry data throughout a large spatial range of western
 81 Canada (Fig. 1), (2) historical, hourly air temperature data from the ERA5 reanalyzed
 82 dataset (Hersbach *et al.*, 2023), and (3) temporally static rasters of percent forest cover,
 83 elevation, and distance from water. We then forecast each species' movement behaviour
 84 throughout the 21st century using monthly climate change projections under four Shared
 85 Socioeconomic Pathways (SSPs; Riahi *et al.*, 2017; Mahony *et al.*, 2022): SSP 1-2.6 ("Sus-
 86 tainability – Taking the Green Road"; low challenges to mitigation and adaptation), SSP
 87 2-4.5 ("Middle of the Road: medium challenges to mitigation and adaptation"), SSP 3-7.0
 88 ("Regional Rivalry – A Rocky Road"; high challenges to mitigation and adaptation), and
 89 SSP 5-8.5 ("Fossil-fueled Development – Taking the Highway"; high challenges to mitigation,
 90 low challenges to adaptation).

91 We acknowledge that our models of habitat selection ignore many important variables of
 92 habitat selection (e.g., forest type, structure, and age; terrain slope and aspect; prey avail-
 93 ability; population density). We limited our models to only use three simple and temporally
 94 static rasters to produce simple and easily interpretable estimates and forecasts under the
 95 assumption of temporally invariant landscapes.




Figure 1: GPS telemetry data for the six species in this study (dots) over a digital elevation model for the study area. The inset in the bottom left shows the telemetry data within the north american continent. Both maps use the BC Albers Equal Area Conic projection (EPSG:3005).

96 **2.1.1 GPS telemetry data**

97 Boreal caribou and wolf telemetry data were acquired from a data repository of the British
98 Columbia Oil and Gas Research and Innovation Society (BC ORGIS) that was collected as
99 part of the boreal caribou radio-collaring program of the BC OGRIS Research and Effec-
100 tiveness Monitoring Board (REMB; BC OGRIS, 2018). Southern mountain caribou data
101 were obtained from Ford *et al.* (2023). Cougar data are from Darlington *et al.* (2025) and
102 (**serrouya_ref?**). Elk data from the work of Ciuti *et al.* (2012) were downloaded from
103 Movebank (Kays *et al.*, 2022). Finally, grizzly bear telemetry data are from (**ref?**), while
104 mountain goat data are from (**ref?**).

105 From the full set of telemetry data, we removed 2,396 problematic GPS locations (0.16%,
106 including collar calibration data) after inspecting diagnostic plots of (1) distance from the
107 median location, (2) straight-line displacement between locations, (3) turning angle, and
108 (4) time interval between consecutive points. Particular attention was paid to points with
109 large turning angles ($\gtrapprox 170^\circ$) and excessively high straight-line displacement, especially if
110 antecedent and subsequent points indicated relatively stationary behaviour (Appendix A).

111 **2.1.2 Historical temperature data and climate change projections**

112 Rasters of hourly air temperature data were downloaded from the ERA5 database (Hers-
113 bach *et al.*, 2023) from the European Center for Medium-range Weather Forecasting server
114 (ECMWF; www.ecmwf.int; <https://cds.climate.copernicus.eu>). Proximal air temperature
115 was estimated for each location by extracting the value from the corresponding raster cell
116 from the temporally nearest raster using the `extract()` function from the `terra` package
117 (v. 1.7-46, Hijmans, 2023) for `R` (R Core Team, 2024).

118 To obtain projected average monthly temperatures from 2025 to 2100 at a 0.08° spatial
119 resolution, we used the `climatenetR` package (v. 1.0, Burnett, 2023) for `R` and ClimateNA
120 v. 7.4.2 (Wang *et al.*, 2016; Mahony *et al.*, 2022). However, the climate projections only
121 included estimates of future monthly averages, a scale substantially coarser than that of our

tracking data (Table 1). To estimate the distribution of temperatures at an hourly scale within a month, we assumed values to be approximately normally distributed with mean μ_T and variance σ_T^2 . We estimated μ_T using the ClimateNA projections, while we estimated σ_T^2 by modeling the observed variance in historical weather data for western Canada from 1998 to 2023 (inclusive). For each location $\langle x, y \rangle$ and month m of each year (e.g., the observed variance at coordinates $\langle -119.40, 49.94 \rangle$ in April 2005), we calculated the variance in historical temperature data, for a total of $(2024 - 1998) \times 12 = 312$ observations per location. We then modeled σ_T^2 a Generalized Additive Model (GAM) for Location and Scale (GAMLS, Rigby & Stasinopoulos, 2005; Stasinopoulos & Rigby, 2007; section 7.9 in Wood, 2017) fitted with the `mgcv` package for R (v. 1.9-1, Wood, 2017). The linear predictor for the location (i.e., the mean) included smooth terms of: each year's estimated within-pixel monthly mean temperature ($\hat{\mu}_T$), month (m), a two-dimensional smooth of spatial coordinates $\langle x, y \rangle$, and a tensor product interaction term of month and space to allow for seasonal terms to vary smoothly over space. The linear predictor for the scale term, which governs the mean-variance relationship (table 3.1 on p. 104 of Wood, 2017), included smooth terms of the monthly mean temperature, month, and space. We did not include smooth terms of year to avoid unrealistic projections when extrapolating beyond past 2023. The complete model for the distribution of projected temperature was thus

$$\left\{ \begin{array}{l} T \sim \text{Normal}(\mu_T, \sigma_T^2) \\ \mu_T \approx \hat{\mu}_T \text{ as projected by ClimateNA} \\ \sigma_T^2 \sim \text{Gamma}(\mu_{\sigma_T^2}, \nu_{\sigma_T^2}) \\ \mathbb{E}(\sigma_T^2) = \mu_{\sigma_T^2} \\ \mathbb{V}(\sigma_T^2) = (\mu_{\sigma_T^2})^2 / \nu_{\sigma_T^2} \\ \log(\mu_{\sigma^2}) = \beta_{L,0} + f_{L,1}(\mu_T) + f_{L,2}(m) + f_{L,3}(x, y) + f_{L,4}(x, y, m) \\ \log(\nu_{\sigma^2}) = \beta_{S,0} + f_{S,1}(\mu_T) + f_{S,2}(m) + f_{S,3}(x, y) \end{array} \right. , \quad (1)$$

where $\mu_{\sigma_T^2}$ and $\nu_{\sigma_T^2}$ indicate the location and scale parameters of the gamma distribution of σ_T^2 , and together they determine the variance of σ_T^2 , indicated as $\mathbb{V}(\sigma_T^2)$. Functions $f_{L,j}$ and $f_{S,j}$ indicate the j^{th} smooth functions for the location and scale parameters, respectively. To ensure the smooth functions of month, $f_{L,2}(m)$ and $f_{S,2}(m)$, joined smoothly between December and January, the terms use cyclic cubic spline bases. The spatial terms used two-dimensional Duchon splines, a generalization of thin-plate regression splines (p. 221 of Wood (2017)). The smoothness parameters were estimated via REstricted Maximum Likelihood (REML; p. 83 of Wood, 2017). See the Data Availability Statement for additional information on the code used to fit the model.

We simulated hourly variation in future years by assuming hourly temperature followed a normal distribution with mean specified by the monthly `climatenar` climate projections and variance as specified by the gamma GMLS. For each month within each year from 2025 to 2100, we simulated hourly weather by including temperatures from the 0.1 to the 0.9 quantiles by increments of 0.1, and we weighted each quantile proportionally to the normalized Gaussian probability density for each quantile.

2.1.3 Habitat rasters

We estimated percent forest cover and distance from water using the temporally static rasters created by Tuanmu & Jetz (2014). We calculated total forest cover by summing the four rasters of evergreen/deciduous needle-leaf trees, evergreen broadleaf trees, deciduous broadleaf trees, and mixed/other trees (raster classes 1-4, respectively). We converted the raster of percent cover of open water (class 12) to a binary raster of pixels with water (cover > 0%) or without water (cover = 0%) and then calculated each pixel's distance from the nearest pixel with water using the `distance()` function from the `terra` package. Finally, we obtained two digital elevation models using the `get_elev_raster()` function from the `elevatr` package (v. 0.99.0, Hollister *et al.*, 2023). We used a raster with a zoom of 6 (a resolution of 0.009°) for model fitting and one with a zoom of 3 (a resolution of 0.08°)

166 for downloading climate change projections via `climatenetR`. All final rasters and code are
167 available on GitHub (see the Data Availability Statement). For ease of interpretability and
168 comparison with current conditions, we assumed resources would remain constant through
169 to 2100, although we recognize that the spatial distribution of forest cover and open water
170 will change throughout the 21st century.

171 **2.2 Estimating mammals' instantaneous speeds**

172 We modeled each animal's movement using continuous-time movement models (Fleming
173 *et al.*, 2014) via the `ctmm` package (v. 1.2.0, Fleming & Calabrese, 2023) for R. We then
174 estimated each mammal's instantaneous speed at each observed location by applying the
175 `ctmm::speeds()` function on all models with finite speed estimates (415 of 433; Fleming *et*
176 *al.*, 2014; Noonan *et al.*, 2019a). The remaining 18 movement models had sampling rates that
177 were too coarse, relative to the animals' directional persistence, to reconstruct the animals'
178 movement trajectories (Noonan *et al.*, 2019a; Nathan *et al.*, 2022; DeNicola *et al.*, 2025).
179 The models were for one mountain goat, 15 boreal caribou, and two southern mountain
180 caribou (Table 1).

181 Since `ctmm`'s movement models assume a single moving state with stochastic but non-zero
182 speed (Calabrese *et al.*, 2016; Noonan *et al.*, 2019a), we corrected data-informed speeds so
183 that the minimum instantaneous speed could be 0. We performed this correction by sub-
184 tracting each model's mean speed while assuming speeds were χ^2 -distributed. The function
185 we used is available on GitHub (see the Data Availability Statement). To avoid artifacts
186 due to excessively small, non-zero speeds, we determined whether an animal was moving or
187 not using a *k*-means algorithm with 2 clusters for each species' distribution of detrended
188 speeds. When the algorithm clearly failed to discriminate between states, we estimated the
189 split point using the inflection points in histograms of the detrended speeds (Fig. B1).

190 **2.3 Estimating the effects of temperature on mammals' movement behaviour**

191 Ambient temperature is only one of the many drivers of mammalian movement behaviour
192 (Fig. 2). Many species alter their movement rates (e.g., movement frequency and speed)
193 daily or seasonally in response to factors such as solar time, photoperiod, forage availability,
194 reproductive cycles, and predator avoidance. Similarly, ambient temperature also fluctuates
195 throughout the day and across seasons. Therefore, estimating the effects of temperature
196 on movement rates requires accounting for how mammals' response to temperature changes
197 with time of day and day of year (Fig. 2A; Péron *et al.*, 2017; Buderman *et al.*, 2018; Leclerc
198 *et al.*, 2021; Xu *et al.*, 2021). Similarly, mammals' selection strength for resources depends
199 on ambient temperature, since higher temperatures can promote a selection for refuge from
200 heat (e.g., thicker forest cover, higher elevation, proximity to water; Attias *et al.*, 2018;
201 Giroux *et al.*, 2023).

202 To assess the importance of including temperature as an explicit covariate (as opposed
203 to including its effects with time of day and day of year), we fit models with and without
204 smooth effects of temperature and compared the fits of the two sets of models via analyses
205 of deviance (a form of generalized likelihood ratio tests) following the methods of Section
206 3.3 in Wood (2017). See Appendix B for additional information.

207 **2.3.1 Effects of temperature on movement rates**

208 We estimated the effects of temperature on mammals' instantaneous movement state (mov-
209 ing or not) and speed when moving using two Hierarchical Generalized Additive Models
210 (HGAMs; Pedersen *et al.*, 2019 and Appendix B) with the `mgcv` package for R. The first
211 HGAM estimated the probability that an animal was moving, $P(M)$, with a binomial fam-
212 ily of distributions and logit link function. The second HGAM estimated an animal's speed
213 when moving with a gamma family of distributions and log link function. We fit the models
214 with fast Restricted Maximum Likelihood ('fREML') and discretized covariates (`discrete =`
215 `TRUE`) to optimize computational efficiency with no appreciable losses to model performance




Figure 2: Directed Acyclical Graphs (DAGs) assumed for inferring the causal effects of temperature (red) on each species' movement behaviour. (A) Ambient temperature affects mammals' movement rates (i.e. probability of moving, speed when moving, and their product: hourly distance traveled). The effects of temperature on mammals' movement rates depend on circadian rhythm and phenology, since animals may respond to temperatures differently at different times of day and or days of year. Additionally, both temperature and phenology depend on time of day and day of year. Finally, circadian rhythm changes with seasonal phenology due to changes in photoperiod (e.g., the time of twilight changes throughout the year, affecting crepuscular activity). (B) Similarly, habitat selection depends on the availability and selection of habitat variables (forest cover, elevation, and distance from water), but the selection strength for each variable is conditional on temperature. For example, an animal may select for more densely-forested areas at extreme temperatures. The resource selection functions for (B) also included marginal smooths of temperature to account for sampling biases across seasons.

216 (Appendix B; Wood *et al.*, 2015, 2017; Li & Wood, 2020). Together, the binomial HGAM
 217 and the gamma HGAM inform us on an animal's long-term average speed, since it is the
 218 product of the probability of moving and its average speed when moving.

219 The HGAMs (equations (2) and (3)) included fixed-effect intercepts for each species
 220 (β_s), random intercepts for each animal (Z_a), and species-level **by** smooths that allowed
 221 independent smoothness parameters for each species (model I in Figure 4 of Pedersen *et*
 222 *al.*, 2019). The **by** smooths accounted for trends in time of day (in Pacific Daylight Time;
 223 `tod_pdt`), day of year (`doy`), and temperature (`temp_c`). To account for the cyclicity of time
 224 of day and day of year, the smooth terms used cyclic cubic splines (p. 202 of Wood, 2017).
 225 The models also had three tensor product interaction terms by each species: (1) day of year
 226 and time of day, (2) temperature and time of day, and (3) temperature and day of year.
 227 These three terms accounted for smooth changes in: (1) daily behaviour across day of year,
 228 (2) the response to temperature over time of day (e.g., changes in nocturnality), and (3)

229 the response to temperature over day of year (e.g., the timing of molting, migration, and
 230 hibernation). Finally, two smooth terms of log-transformed sampling interval (dt ; hours)
 231 corrected for biases in speed estimates arising from irregular GPS sampling intervals, since
 232 longer intervals result in lower speed estimates (Nathan *et al.*, 2022; DeNicola *et al.*, 2025).
 233 A global smooth term of $\log(\text{dt})$ accounted for the overall effect of sampling interval, while
 234 a factor-smooth interaction term ($\text{bs} = \text{'fs'}$) of $\log(\text{dt})$ and species accounted for species-
 235 level deviations from the global term while assuming a common smoothness parameter across
 236 species (model GS in Figure 4 of Pedersen *et al.*, 2019). Formally, the model for movement
 237 state M , with $M = 0$ indicating no movement and $M = 1$ indicating movement, was

$$\left\{ \begin{array}{l} M \sim \text{Bin}(p) \\ \mathbb{E}(M) = p \\ \mathbb{V}(M) = p(1-p) \\ \log(p) = \beta_s + Z_a + f_{1,s}(\text{tod_pdt}) + f_{2,s}(\text{doy}) + f_{3,s}(\text{temp_c}) + \\ f_{4,s}(\text{doy}, \text{tod_pdt}) + f_{5,s}(\text{temp_c}, \text{tod_pdt}) + f_{6,s}(\text{temp_c}, \text{doy}) + \\ f_7(\log(\text{dt})) + f_{8,s}(\log(\text{dt})) \end{array} \right. , \quad (2)$$

238 while the model for movement speed when moving (i.e., $M = 1$, indicated with S) was

$$\left\{ \begin{array}{l} S \sim \text{Gamma}(\mu_S, \nu_S) \\ \mathbb{E}(S) = \mu_S \\ \mathbb{V}(S) = \mu_S^2 / \nu_S \\ \log(\mu_S) = \beta_s + Z_a + f_{1,s}(\text{tod_pdt}) + f_{2,s}(\text{doy}) + f_{3,s}(\text{temp_c}) + \\ f_{4,s}(\text{doy}, \text{tod_pdt}) + f_{5,s}(\text{temp_c}, \text{tod_pdt}) + f_{6,s}(\text{temp_c}, \text{doy}) + \\ f_7(\log(\text{dt})) + f_{8,s}(\log(\text{dt})) \end{array} \right. . \quad (3)$$

239 In both models, β_s indicates a fixed intercept for species s , Z_a indicates a Gaussian random
240 effect for animal a (of species s), $f_{j,s}$ indicates the j^{th} smooth function for species s , and
241 functions with two variables indicate tensor product interaction terms. The model code used
242 to fit the models is available in Appendix B.

243 **2.3.2 Effects of temperature on habitat selection**

244 We estimated the effects of temperature on each species' selection for percent forest
245 cover (`forest_perc`), elevation (`elevation_m`, in meters), and distance from water
246 (`dist_water_m`, in meters) by fitting a Hierarchical Resource Selection Function (HRSF) for
247 each species (McCabe *et al.*, 2021). We fit each HRSF using an HGAM with a Poisson family
248 of distributions and log link function (Appendix B; Aarts *et al.*, 2008). After removing
249 non-resident individuals (Table B1), we accounted for the spatiotemporal autocorrelation
250 in the telemetry locations by weighting each point based on the telemetry's Autocorrelated
251 Kernel Density Estimate (Fleming & Calabrese, 2017; Noonan *et al.*, 2019b; Alston *et al.*,
252 2022) to produce estimates of second-order habitat selection (Johnson, 1980). Quadrature
253 points were used to approximate the likelihood function of a Poisson point process through
254 Monte-Carlo Markov chain-based integration (Aarts *et al.*, 2008; Fithian & Hastie, 2013;
255 Baddeley *et al.*, 2015) and were determined using the raster cells in each animal's 99.9%
256 AKDE percentile, with each raster cell having a weight of 1. The number of quadrature
257 locations greatly outnumbered the number of observed locations (Fig. B12), especially after
258 accounting for the AKDE weights (Fig. B13).

259 Each species' model had the same structure:

$$\left\{ \begin{array}{l} O \sim \text{Pois}(\lambda) \\ \mathbb{E}(O) = \mathbb{V}(O) = \lambda \\ \log(\lambda) = f_1(\text{forest_perc}) + f_2(\text{elevation_m}) + f_3(\text{dist_water_m}) + \\ Z_a + f_{4,a}(\text{forest_perc}) + f_{5,a}(\text{elevation_m}) + f_{6,a}(\text{dist_water_m}) + \\ f_7(\text{forest_perc}, \text{temp_c}) + f_8(\text{elevation_m}, \text{temp_c}) + \\ f_9(\text{dist_water_m}, \text{temp_c}) + f_{10}(\text{temp_c}) + f_{11,a}(\text{temp_c}) \end{array} \right. , \quad (4)$$

where O indicates whether an animal was observed ($O = 1$) or not ($O = 0$), and the species-level indices are omitted for readability, but each term in the model can be assumed to be species-specific. Smooth effects of percent forest cover (`forest_perc`), elevation (`elevation_m`, in meters), and distance to water (`dist_water_m`, in meters) accounted for the species-level selection strength for each resource. A Gaussian random effect for each individual animal (Z_a) corrected for uneven sampling across individuals, while factor-smooth interaction terms for each animal ($f_{j,a}$) accounted for animal-level resource selection (i.e., individual-level deviations from the species-level estimate; Jeltsch *et al.*, 2025). Tensor product interaction terms of the three resources and temperature (`temp_c`) estimated the smooth change in resource selection at different temperatures. Finally, marginal smooth terms of temperature and factor-smooth interaction terms of temperature and animal accounted for species- and individual-level sampling biases at different temperatures (e.g., sampling more during warm periods).

3 Results

Of the GPS fixes with finite speed estimates, 2.6% had temperatures lower than -20°C , while 6.5% had temperatures above 20°C , and temperature ranges differed across species (Table 2, Fig. B2). Sampling interval affected estimates of probability as well of as speed when

moving (Fig. B8). All species' estimated probability of moving and speed when moving decreased with sampling intervals above 1 hour, except for cougars' speed, although the estimated trends were highly uncertain (Fig. B8). Consequently, we present all results while predicting specifically for one-hour sampling intervals. At 0°C, species differed in estimated mean probabilities of moving ($\hat{P}(M = 1)$; range: 0.05 – 0.31), mean speed when moving ($\hat{\mathbb{E}}(S|M = 1)$; range: 0.42 – 2.67 km/h), and mean overall speed (i.e., $\hat{P}(M) \times \hat{\mathbb{E}}(S|M = 1)$, range: 0.04 – 0.61 km/h; Table 2). Grizzly bears had the lowest movement frequency ($\hat{P}(M) \approx 0.05$), while wolves and cougars moved most often ($\hat{P}(M) \geq 0.22$). Mountain goats and southern mountain caribou moved the slowest ($\hat{\mathbb{E}}(S|M = 1) \approx 0.43$ km/h), while wolves had the highest mean speed when moving ($\hat{\mathbb{E}}(S|M = 1) \approx 2.67$ km/h). Consequently, at 0°C, wolves traveled an average of 0.22×2.67 km/h ≈ 0.6 km/h; 2.5 to 16.7 times further than other species.

Table 2: Summary statistics for each species' GPS fixes with finite speed estimates, namely: the number fixes after data cleaning (n), the percentage of fixes with temperature (T) below -20°C and above 20°C , the estimated mean probability of moving ($\hat{P}(M = 1)$), the mean speed when moving ($\hat{\mathbb{E}}(S|M = 1)$; km/h), and the mean hourly distance travelled ($\hat{P}(M = 1) \times \hat{\mathbb{E}}(S|M = 1) = \hat{\mathbb{E}}(D)$; km/h), for a sampling interval of 1 hour and a temperature of $T = 0^{\circ}\text{C}$.

Species	n	T < -20°C (%)	T > +20°C (%)	$\hat{P}(M = 1 T)$	$\hat{\mathbb{E}}(S M = 1, T)$	$\hat{\mathbb{E}}(D T)$
Caribou (boreal)	187,679	6.8	7.9	0.18	0.73	0.13
Caribou (s. mountain)	26,518	1.3	3.4	0.11	0.42	0.05
Cougars	80,621	0.7	6.9	0.31	0.76	0.24
Elk	875,682	2.4	4.9	0.17	0.57	0.10
Grizzly bears	39,001	0.0	8.4	0.05	0.72	0.04
Mountain goats	65,219	0.7	2.8	0.13	0.42	0.06
Wolves	202,386	1.7	13.0	0.22	2.67	0.60
Total	1,477,106	2.6	6.5			

Across all species, Relative Selection Strength (RSS) was weakest for forest cover and strongest for elevation. At temperatures near 0°C, boreal caribou selected for forest cover between 50% and 75%, elevations near 500 m, and distances from water < 10 km, while southern mountain caribou selected for dense forest cover, elevations near 2 km, and distances from water $\lesssim 5$ km. Cougars selected for dense forest cover ($> 75\%$), an elevation of ~ 1

km, and distances from water < 7.5 km. Elk selected for intermediate forest cover (\approx 50%), elevations between 1 and 2 km, and distances from water of 10-15 km. Grizzly bears selected for relatively sparse forest cover (25-50%), elevation between 1 and 2 km, and distances from water < 3 km. Mountain goats selected for sparse forest cover (< 25%), elevations near 1.5 km, and distances from water < 5 km. Finally, wolves selected for forest cover (\gtrapprox 50%), elevations near 1 km, and distances from water < 5 km. There was relatively strong agreement between models with and without temperature (Figs. B3, and B14), but including temperature always resulted in better fits (all p-values $< 2.2 \times 10^{-16}$; all $\Delta AIC \geq 342$; Appendix B). Accounting for temperature even resulted in a simpler HRSF for mountain goats (lower model degrees of freedom due to simpler smooth terms) that fit the data better (lower deviance), and hence no test was necessary to show that the inclusion of temperature produced a better fit.

3.1 Effects of temperature on movement rates

Species' changes in movement rates to temperature varied in both direction and magnitude (Figs. 3, B4-B6), even after accounting for differences in daily and seasonal activity (e.g., sleeping, migration, hibernation; Figs. B4-B6). Smooth interaction terms were well-behaved and indicated clear shifts in activity over time of day and day of year for all species. The models had good in-sample prediction (Fig. B7) and explained reasonably high proportions of the deviance [79.3% for the gamma model and 10.7% for the binomial model, which is relatively high for a binomial model with binary responses]. All species altered their daily and seasonal movement behaviour to changes in temperature (Fig. 3). The response was most visible in cougars. In late spring (June 1st), they moved from evening to early morning if hourly temperatures were below 20°C, but if temperatures were above 20°C they moved mostly between 3:00 and 6:00 AM. Throughout the year, they tended to move more when it was colder, but what they perceived as "colder" was relative to the time of year. Overall, uncertainty around the estimated effects was generally higher at extreme temperatures due

320 to lower data availability (Figs. B4A, B5A, and B6A).




Figure 3: **Temperature is a strong driver of how far and when mammals travel.** The fill indicates the effects of temperature on the relative change in hourly distance travelled (probability of moving times mean speed when moving) over time of day on June 1st (**A**) and day of year at 12:00 PDT (**B**). Predictions in the surface plots extend to 10% of the range away from each datum. The color bar is on the log₂ scale to help visualize patterns in doubling, and values are capped to $2^{\pm 2}$ for ease of readability.

321 3.2 Effects of temperature on habitat selection

322 Species' RSS was generally strongest for elevation and weakest for forest cover, but RSS
 323 depended significantly on temperature for all species (all p-values < 2.2×10^{-16} ; Fig. 4).
 324 Changes in RSS with temperature were strongest for elevation and generally weakest for
 325 distance from water, but there were no common trends across all species for any of the
 326 three resources. All species except cougars exhibited clear temperature-dependent shifts in
 327 their preference for forest cover. At higher temperatures, wolves relaxed their preference
 328 for forested areas, while mountain goats relaxed their preference for open areas (cover <
 329 50%). As temperatures warmed, elk and boreal caribou shifted towards more forest cover
 330 closer to 50%, while southern mountain caribou and grizzly bears selected for areas with
 331 50% forest cover or less. All species shifted elevationally with temperature, although boreal
 332 caribou had a much sharper change in RSS across elevation. Elk, mountain goats, and cougars

moved to higher elevations with temperature, while wolves, southern mountain caribou, and grizzly bears moved to lower elevations. Wolves, elk, and southern mountain caribou moved closer to water with temperature, while mountain goats, cougars, and grizzly bears moved somewhat further away from water but remained mainly within 5-10 km of water. As with movement rates, estimated RSS was generally most uncertain at extreme temperatures, for which data were scarcer (Fig. B15).




Figure 4: Mammals' habitat selection depends on ambient temperature. Estimated relative selection strength (RSS) for forest cover (%), elevation (km), and distance from water (km) as a function of temperature. The grey contours indicate the extent of each species' observed data. RSS values were re-centered and capped to between 0.25 and 4 ($2^{\pm 2}$) to improve readability.

Fig. 5 presents the predicted habitat selection for boreal caribou and wolves. Overall, both species selected for the same low-elevation area (< 500 m; Fig. 5A-B), but caribou generally avoided the river system that wolves were often found at, and their selection strength varied with temperature. As wolves' habitat selection strengthened at warmer temperatures, encounters at 20°C were approximately four times more frequent than at -20°C, independent of changes in the species' movement rates (Fig. 5C).

3.3 Predicted changes in movement behaviour during the 21st century

Predicted changes in movement rates with future climate change varied across species in both magnitude and direction, but worse SSPs always corresponded to greater absolute changes




Figure 5: Temperature affects boreal caribou's and wolves' habitat selection and, consequently, their encounter rates. Panel (A) shows the rasters of the three habitat variables used in the hierarchical resource selection functions, while panel (B) shows the relative selection strength (RSS) for boreal caribou and wolves, as a function of temperature. RSS values were re-centered and capped to between 0.25 and 4 ($2^{\pm 2}$) to improve readability. Finally, panel (C) shows the (scaled) product of the RSS values in panel (B) as a proxy for encounter rates between individuals of the two species.

(Fig. 6). Additionally, species that were predicted to move less often did not necessarily have lower speeds when moving, and vice versa (Figs. B9 and B10). Estimated changes in average yearly distance traveled were negligible for boreal caribou and grizzly bears, although both species showed seasonal changes in seasonal movement rates. As temperatures warmed, boreal caribou were predicted to move more in winter, spring, and fall but less in summer (Fig. 3), while grizzly bears were predicted to show a clear shift towards earlier emergence from hibernation (Fig. B4) and greater movement earlier in the year but less movement in early fall. Southern mountain caribou and mountain goats are projected to travel further by 2100, although the estimates for southern mountain caribou varied greatly over space (Fig. 7). Cougars, elk, and wolves were projected to move less by 2100, with cougars showing as much as a 24% decrease in mean yearly distance travelled.

Absolute relative changes in predicted yearly distance travelled were small under the best-case SSP (0-4% change in 2100 relative to 2025). Under the worst-case SSP, absolute changes by 2100 (relative to 2025) ranged from ~2% (grizzly bears) to ~24% (cougars).

362 Projected changes in 2100 varied spatially due to spatial heterogeneity in climate change
 363 projections (Fig. 7). Again, absolute changes were generally greatest under worse SSPs, but
 364 the direction of change at each location also varied across SSPs (most visible in cougars).




Figure 6: Species are predicted to alter their movement rates differently in response to climate change, and worse climate-change scenarios will result in the greatest change. Bold lines indicate the median change in yearly distance travelled (probability of moving times speed when moving) due to predicted changes in temperature within each species' study area. Shaded areas indicate the 95th and 5th percentiles. Changes are relative to the mean predicted distance travelled at each location in 2025, averaged across the four Shared Socioeconomic Pathways (SSPs). Values > 1 indicate an increase, while values < 1 indicate a decrease. The projections only account for changes in movement frequency and speed, and they ignore changes in physiology or movement costs.

365 Median RSS was projected to decrease over time within each species' observed range, but,
 366 again, changes were stronger under worse SSPs (Fig. 8). Decreases were most pronounced
 367 in areas with the lowest RSS and most severe for elk, mountain goats, cougars, and southern
 368 mountain caribou. Changes for boreal wolves and caribou were negligible. Elk, cougars,
 369 and grizzly bears were predicted to increase their selection strength for top-RSS areas, and
 370 elk, mountain goats, cougars, and southern mountain caribou were predicted to further
 371 decrease their selection for areas with low RSS. Unsurprisingly, the predicted change in
 372 RSS between 2025 and 2100 also varied spatially for all species (Fig. 9). Overall, RSS
 373 decreased throughout most of each species' current range, although elk, cougars, and bears




Figure 7: Climate change is predicted to impact each species' movement rates differently, but changes will also vary spatially. The color of each pixel indicates the predicted changes in distance traveled in 2100 at that pixel, relative to the value in 2025, averaged across all four scenarios. Values < 1 indicate a decrease, and values > 1 indicate an increase. For ease of readability, the color bar is on the \log_2 scale to help visualize patterns in doubling). The predictions only account for the predicted temperature throughout the areas and ignore environmental factors such as terrain slope, soil type, and forest cover. All maps extend to each species' study area (Fig. 1) and are shown in the BC Albers Equal Area Conic projection (EPSG:3005).

³⁷⁴ were predicted to increase their selection for higher-altitude habitats. Still, none of the
³⁷⁵ species were projected to increase RSS throughout their habitat (10).




Figure 8: **Species are predicted to alter their habitat selection differently in response to climate change, and worse climate-change scenarios will result in the greatest change.** Bold lines indicate the change in median RSS due to predicted changes in temperature within each species' study area. Shaded areas indicate the 95th and 5th percentiles in change in RSS. Values > 1 indicate an increase, while values < 1 indicate a decrease. Changes are relative to each location's mean RSS in 2025, averaged across the four Shared Socioeconomic Pathways (SSPs).

³⁷⁶ 4 Discussion

³⁷⁷ We have demonstrated that temperature is an important driver of how and where large
³⁷⁸ mammals move, and that it affects mammals' seasonal and daily movement behaviour in
³⁷⁹ complex and nonlinear ways. However, predicting mammals' responses to climate change
³⁸⁰ remains a challenge, as habitats are warming rapidly and mammals are exposed to increas-
³⁸¹ ingly novel and extreme conditions (Diffenbaugh & Field, 2013; Botero *et al.*, 2015; IPCC,
³⁸² 2023). We leveraged the flexibility and interpretability of HGAMs to estimate mammals'
³⁸³ nonlinear responses to temperature without imposing rigid assumptions about the shape of
³⁸⁴ the responses (Simpson, 2018, 2025). In particular, tensor product interaction terms were



Figure 9: Climate change will impact each species' relative selection strength (RSS) differently. The color scale indicates the predicted changes in RSS in 2100, relative to each location's average RSS in 2025 across all four scenarios (i.e., not relative to other pixels). Values < 1 indicate a decrease and values > 1 indicate an increase. For ease of readability, the color bar is on the \log_2 scale to help visualize patterns in doubling, and values are capped to 0.8 and 1.25 ($\approx 2^{\pm 0.322}$; original data ranged 0.71 to 1.93). All maps extend to each species' study area and are shown in the BC Albers Equal Area Conic projection (EPSG:3005).




Figure 10: Climate change is predicted to cause species to decrease their selection strength for their current habitats, and worse climate change scenarios will have the greatest impact. The density plots indicate each species’ change in RSS in 2100 for their current habitat (Fig. 8), relative to each location’s RSS in 2025. Values < 1 indicate a decrease and values > 1 indicate an increase. For ease of readability, values > 1.3 were changed to 1.3 (original maximum: 1.93).

385 crucial for estimating smooth, nonlinear changes in daily and seasonal trends with tem-
 386 perature with data from multiple individuals. By allowing interaction terms to be smooth
 387 bivariate functions, we were able to quantify changes in movement behaviour without the
 388 need to discretize time into windows (e.g., day/night and seasons). We were thus able to
 389 present results using relatively simple and interpretable surface plots (Figs. 3 and 4) that
 390 conveyed nuanced, continuous-time estimates of change. This framework was essential for
 391 estimating how mammals behave at extreme temperatures. In fact, despite the recent in-
 392 crease in frequency of extreme heat events in western Canada, including a severe heat wave
 393 in 2021 (Kienzle, 2018; Thompson *et al.*, 2022; Zhang *et al.*, 2023), such events remain rela-
 394 tively rare, so estimating mammals’ nonlinear responses required the flexibility provided by
 395 HGAMs (Wood, 2017). Additionally, the hierarchical approach allowed us to leverage data
 396 from multiple individuals in a single model while accounting for differences in behaviour and
 397 data availability across individuals (Pedersen *et al.*, 2019; Muff *et al.*, 2020; McCabe *et al.*,

398 2021; Jeltsch *et al.*, 2025; Silva *et al.*, 2025). Still, GAMs' flexibility and complexity can be
399 a frustrating limitation if used too generously, so we suggest choosing each smooth term's
400 basis size (k) carefully. While penalized splines and REML can help avoid unruly smooth
401 terms, the penalty terms can struggle to restrict model complexity when datasets are as
402 large as those presented here (Wood, 2017; Wood *et al.*, 2017).

403 The effects of warming temperatures on movement rates varied across species, seasons,
404 and space due to differences in species' behavior and physiology as well as their habitats'
405 phenology (e.g., the timing of green-up). For boreal caribou, hotter temperatures decreased
406 movement rates summer but increased them otherwise, mirroring previous studies (Stien *et*
407 *al.*, 2012; Leclerc *et al.*, 2021; Lessard *et al.*, 2025). The increase in movement rates with
408 temperature during the cold months is likely partly due to shallower snow depth, which
409 results in greater mobility (Pedersen *et al.*, 2021), but warmer temperatures during snowy
410 seasons also increase the risk of rain-on-snow events, which limit forage availability and
411 increase time spent foraging (Stien *et al.*, 2012; Berger *et al.*, 2018; Mezzini *et al.*, 2025).
412 The reduced movement rates during warmer summers are likely mainly due to more frequent
413 resting and use of thermal refugia, given ungulates' elevated risk of hyperthermia (Alston
414 *et al.*, 2020; Verzuh *et al.*, 2023). Therefore, although we predict that boreal caribou's
415 yearly average movement rates will remain approximately unchanged throughout the 21st
416 century, the projections do not show the changes in seasonal movement phenology. This is an
417 important caveat, since the effects of extreme temperatures are exacerbated by phenological
418 mismatches with seasonal photoperiod (Walker *et al.*, 2019), including the timing of molting
419 and reproduction. Earlier growth seasons in boreal and arctic regions have resulted in lower
420 calf birth and survival (Post & Forchhammer, 2008), while the lengthening of the growth
421 season has allowed moose (*Alces alces*) and deer to encroach on boreal caribou habitat and
422 increase the density of coyotes (*Canis latrans*), cougars, and wolves (Barber *et al.*, 2018;
423 DeMars *et al.*, 2023), whose movement behaviour also depends on ambient temperature.

424 Boreal wolves responded to temperature similarly to boreal caribou, and habitat selec-

425 tion strength was not projected to change noticeably by 2100, but wolves were predicted
426 to move less in future decades. In contrast, cougars showed markedly different seasonal
427 responses to temperature, with warmer temperatures always resulting in less travel, which
428 resulted in strong predicted decreases in movement rates over the 21st century. However,
429 spatial heterogeneity in habitat, climate, and cougar behaviour may cause boreal individuals
430 to respond to temperature differently from those in our study. Still, rising temperatures
431 clearly affect more than each individual species' movement behaviour; they have complex
432 and interconnected effects on entire communities and their trophic structures.

433 Cunningham *et al.* (2021) discuss the importance of adopting a community-level per-
434 spective when assessing the effects of climate change on animals' movement behaviour and
435 phenology. For example, as temperatures warm and both boreal caribou and wolves in-
436 crease their movement rates in fall, winter, and spring, they also increase their encounter
437 rates (Martinez-Garcia *et al.*, 2020), which will likely increase predation risk when caribou
438 are pregnant or with young calves. Additionally, while both species saw reduced movement
439 rates during hot summer days, boreal wolves' use of anthropogenic linear features (e.g., roads,
440 seismic lines) may allow them to reduce the total thermal costs of movement by moving for
441 shorter periods while increasing the chances of encountering heat-stressed prey (Whittington
442 *et al.*, 2011; Dickie *et al.*, 2017; Dickie *et al.*, 2022). Furthermore, caribou that attempt to
443 reduce predation risk from wolves by avoiding wolf habitat may still risk increasing predation
444 pressure from bears, cougars, and coyote (Leblond *et al.*, 2016; DeMars *et al.*, 2023; Labadie
445 *et al.*, 2023).

446 Considerations about changes in trophic interactions leads to another important caveat
447 about the results we present. The estimated changes in movement behaviour cannot be fully
448 attributed to physiological responses to temperature alone, since other aspects of habitats'
449 seasonal phenology are (nonlinearly) correlated with temperature. For example, mountain
450 goats' lower movement rates at temperatures near 0°C in winter may be in part due to
451 increased movement difficulty, since melting snow and rain-on-snow events increase the en-

452 energetic costs of movement (White, 2025) and encounter rates with predators (Sullender *et*
453 *al.*, 2023). However, milder temperatures can also increase plant growth, which allows goats
454 to spend more time foraging within and less time travelling across patches (Charnov, 1976;
455 White *et al.*, 2025). Therefore, figure 3 illustrates an estimate of the total effects of temper-
456 ature on each species' movement rates, but it does not explicitly account for energetic costs.
457 Similarly, species' changes in habitat selection do not explicitly account for any changes in
458 physiological or energetic costs. Additionally, forest cover, elevation, and distance from water
459 are often correlated, so the estimated effects should be interpreted carefully, and they may
460 not be applicable outside the study areas used here. For example, both mountain goats and
461 elk selected for higher elevation as temperatures warmed, but our models did not account
462 for differences in forage availability at different elevations.

463 Climate change has also had extensive impacts on plant growth and phenology (Cleland *et*
464 *al.*, 2007; Denny, 2019; Tysor, 2025) with consequences on herbivores' behaviour and fitness
465 (Post & Forchhammer, 2008; Aikens *et al.*, 2017). Additionally, plants' limited ability to
466 disperse to and establish in new habitats is often a limiting factor for herbivores' ability to
467 escape hotter temperatures by moving to higher altitudes or latitudes, and the advance of
468 herbivores can further limit plants' ability to establish (Speed *et al.*, 2012; Diffenbaugh &
469 Field, 2013). Consequently, mammals that move to higher elevations may be forced to spend
470 more time foraging (Mezzini *et al.*, 2025), which will increase both their energetic expenditure
471 and encounter rates with predators and competitors (Kohl *et al.*, 2019; Martinez-Garcia *et*
472 *al.*, 2020; Tórrez-Herrera *et al.*, 2020; Smith *et al.*, 2023). If local peaks become unsuitable
473 habitat, and moving to nearby higher peaks requires traveling across low-quality and high-
474 risk habitat (Figure 9), habitat fragmentation may prevent movement across ranges (White
475 *et al.*, 2025).

476 Growing evidence suggests that mammals' exposure, sensitivity, and adaptability to cli-
477 mate change will limit their ability to adapt to change (Ameca y Juárez *et al.*, 2012; Diffen-
478 baugh & Field, 2013). Exposure to warming temperatures is widespread throughout western

479 Canada (Bunnell *et al.*, 2011; Kienzle, 2018; Eum *et al.*, 2023), and changes in temperature
480 and phenology may be too rapid for many species to respond to (Diffenbaugh & Field, 2013).
481 Species' adaptability is further challenged by the difficulty in responding to concurrent, mul-
482 tifarious, and interconnected changes (Diffenbaugh & Field, 2013; Polazzo *et al.*, 2024). In
483 the future, landscapes may thus become unable to support species and population sizes they
484 supported historically, and species may begin selecting against historical ranges (Williams
485 & Blois, 2018). Therefore, it is crucial for conservationists to distinguish between a species'
486 ideal environmental space and its best available geographic space (*sensu* Matthiopoulos,
487 2022).

488 Accelerating multifarious change in species' movement behaviour and energetic costs re-
489 quire conservation measures that are proactive and anticipate future changes. Polazzo *et al.*
490 (2024) present a framework for estimating species' responses to multifarious change using
491 GAMs that is particularly useful when estimating species' (nonlinear) responses to inter-
492 actions between environmental variables (e.g., changes in both temperature and limiting
493 nutrients). By accounting for the effects of multiple variables at once, they demonstrate
494 how a species' response to one variable can depend on other concurrent changes. Estimat-
495 ing responses to multifarious change is crucial in assessing mammals' responses to hotter
496 temperatures, altered phenology, and novel community dynamics.

497 Protected areas must support populations not only in present conditions but also as
498 species' needs and behaviour change over the decades (Simmons *et al.*, 2021; Livingstone *et*
499 *al.*, 2023). Recent evidence has questioned whether current protected areas will be effective
500 in the long term (Loarie *et al.*, 2009; Diffenbaugh & Field, 2013; Williams & Blois, 2018),
501 and increasingly more experts have been calling for proactive and long-term conservation.
502 By identifying and actively protecting climate change refugia and corridors, we may be able
503 to reduce the effects of climate change on mammalian movement behaviour and community-
504 level responses and ensure that protected areas may be viable for decades to come (Michalak
505 *et al.*, 2018; Graham *et al.*, 2019; Stralberg *et al.*, 2020; Hua *et al.*, 2022).

506 Future work should develop models that account for covariates that are specific to the
507 study species and area, such as forest age and disturbance (DeMars *et al.*, 2023), terrain
508 ruggedness (White & Gregovich, 2017), anthropogenic linear features (Dickie *et al.*, 2017),
509 and growing human presence in wild spaces (Sih *et al.*, 2011; Rice, 2022), including the
510 growing risk of human-wildlife conflict (Abrahms *et al.*, 2023; Weststrate *et al.*, 2024). In
511 particular, forests in western Canada are highly dynamic over time (Zhang *et al.*, 2015), and
512 fires size and burn severity have increased in western Canada in recent decades (Whitman *et*
513 *al.*, 2022; Parisien *et al.*, 2023; Wang *et al.*, 2025). Thus, while our use of a static raster of
514 forest cover allowed for simpler models and climate change projections, future studies should
515 account for mammals' responses to temporal forest dynamics, including the impacts of fire
516 (Whitman *et al.*, 2022; Parisien *et al.*, 2023; Wang *et al.*, 2025), logging (Lochhead *et al.*,
517 2022) and other extractive industries (Dickie *et al.*, 2017, 2024). However, the inclusion of
518 temporally dynamic variables will require that any future projections of species' behaviour
519 depend on temporally dynamic estimates of how such variables will change over time with
520 changes in climate and anthropogenic activity.

521 We have demonstrated that temperature has complex and nonlinear effects on how and
522 where mammals move. Rising temperatures have impacted mammals' daily and seasonal
523 movement cycles, and changes are expected to intensify throughout the 21st century. Quan-
524 tifying how temperature affects mammalian behavior and fitness is a first step towards
525 developing an understanding of the effects of climate change on landscape dynamics and the
526 effectiveness of protected areas in the future. By leveraging the flexibility and statistical
527 power of Hierarchical Generalized Additive Models, we have provided researchers with mal-
528 leable and data-driven methods for assessing species' responses to climate change. We hope
529 that future work will develop more species-specific models with the intent of developing a
530 community-level framework for quantifying and predicting how communities will respond to
531 climate change.

⁵³² **Author contributions**

⁵³³ SM performed the data cleaning, ran the analyses, and wrote the manuscript. CHF wrote
⁵³⁴ code for estimating instantaneous movement speeds. MJN conceived of the project idea and
⁵³⁵ supervised SM throughout the project. All other authors contributed telemetry data and
⁵³⁶ reviewed the interpretation of the results for their species of interest. KH and SD contributed
⁵³⁷ substantially to manuscript review. All authors contributed to reviewing the manuscript.

538 **References**

- 539 Aarts G, MacKenzie M, McConnell B, Fedak M, Matthiopoulos J (2008) [Estimating space-](#)
540 [use and habitat preference from wildlife telemetry data](#). *Ecography*, **31**, 140–160.
- 541 Abrahms B, Carter NH, Clark-Wolf TJ et al. (2023) [Climate change as a global amplifier of](#)
542 [human–wildlife conflict](#). *Nature Climate Change*, **13**, 224–234.
- 543 Aikens EO, Kauffman MJ, Merkle JA, Dwinnell SPH, Fralick GL, Monteith KL (2017) [The](#)
544 [greenscape shapes surfing of resource waves in a large migratory herbivore](#) (ed Nathan
545 R). *Ecology Letters*, **20**, 741–750.
- 546 Alston JM, Joyce MJ, Merkle JA, Moen RA (2020) [Temperature shapes movement and](#)
547 [habitat selection by a heat-sensitive ungulate](#). *Landscape Ecology*, **35**, 1961–1973.
- 548 Alston JM, Fleming CH, Kays R et al. (2022) [Mitigating pseudoreplication and bias in](#)
549 [resource selection functions with autocorrelation-informed weighting](#). *Methods in Ecology*
550 [and Evolution](#), 2041–210X.14025.
- 551 Ameca y Juárez EI, Mace GM, Cowlishaw G, Pettorelli N (2012) [Natural population die-offs:](#)
552 [Causes and consequences for terrestrial mammals](#). *Trends in Ecology & Evolution*, **27**,
553 272–277.
- 554 Arechavala-Lopez P, Minguito-Frutos M, Follana-Berná G, Palmer M (2019) [Common octo-](#)
555 [pus settled in human-altered mediterranean coastal waters: From individual home range](#)
556 [to population dynamics](#) (ed Durif C). *ICES Journal of Marine Science*, **76**, 585–597.
- 557 Attias N, Oliveira-Santos LGR, Fagan WF, Mourão G (2018) [Effects of air temperature on](#)
558 [habitat selection and activity patterns of two tropical imperfect homeotherms](#). *Animal*
559 [Behaviour](#), **140**, 129–140.
- 560 Baddeley A, Rubak E, Turner R (2015) [Spatial point patterns: Methodology and applications](#)
561 [with r](#), 0th edn. Chapman; Hall/CRC.
- 562 Barber QE, Parisien M, Whitman E et al. (2018) [Potential impacts of climate change on](#)
563 [the habitat of boreal woodland caribou](#). *Ecosphere*, **9**, e02472.

- 564 Bartashevich P, Herbert-Read JE, Hansen MJ, Dhellemmes F, Domenici P, Krause J, Ro-
565 manczuk P (2024) Collective anti-predator escape manoeuvres through optimal attack
566 and avoidance strategies. *Communications Biology*, **7**, 1586.
- 567 Basu A, Culpepper J, Blagrave K, Sharma S (2024) Phenological shifts in lake ice cover
568 across the northern hemisphere: A glimpse into the past, present, and the future of lake
569 ice phenology. *Water Resources Research*, **60**, e2023WR036392.
- 570 BC OGRIS (2018) Boreal caribou and wolf telemetry data.
- 571 Berger J, Hartway C, Gruzdev A, Johnson M (2018) Climate degradation and extreme icing
572 events constrain life in cold-adapted mammals. *Scientific Reports*, **8**, 1156.
- 573 Botero CA, Weissing FJ, Wright J, Rubenstein DR (2015) Evolutionary tipping points in
574 the capacity to adapt to environmental change. *Proceedings of the National Academy of
575 Sciences*, **112**, 184–189.
- 576 Brivio F, Apollonio M, Anderwald P, Filli F, Bassano B, Bertolucci C, Grignolio S (2024)
577 Seeking temporal refugia to heat stress: Increasing nocturnal activity despite predation
578 risk. *Proceedings of the Royal Society B: Biological Sciences*, **291**, 20231587.
- 579 Brown JH, Gillooly JF, Allen AP, Savage VM, West GB (2004) Toward a metabolic theory
580 of ecology. *Ecology*, **85**, 1771–1789.
- 581 Buderman FE, Hooten MB, Alldredge MW, Hanks EM, Ivan JS (2018) Time-varying preda-
582 tory behavior is primary predictor of fine-scale movement of wildland-urban cougars.
583 *Movement Ecology*, **6**, 22.
- 584 Bunnell FL, Kremsater LL, Wells RW (2011) Global weirding in british columbia: Climate
585 change and the habitat of terrestrial vertebrates. *Journal of Ecosystems and Manage-
586 ment*, **12**.
- 587 Burnett M (2023) climatenaR: Tools to access ClimateNA data.
- 588 Calabrese JM, Fleming CH, Gurarie E (2016) Ctmm: An r package for analyzing animal
589 relocation data as a continuous-time stochastic process (ed Freckleton R). *Methods in
590 Ecology and Evolution*, **7**, 1124–1132.

- 591 Carbeck K, Wang T, Reid JM, Arcese P (2022) Adaptation to climate change through
592 seasonal migration revealed by climatic versus demographic niche models. *Global Change
593 Biology*, **28**, 4260–4275.
- 594 Charnov EL (1976) Optimal foraging, the marginal value theorem. *Theoretical Population
595 Biology*, **9**, 129–136.
- 596 Ciuti S, Muhly TB, Paton DG, McDevitt AD, Musiani M, Boyce MS (2012) Human selection
597 of elk behavioural traits in a landscape of fear. *Proceedings of the Royal Society B:
598 Biological Sciences*, **279**, 4407–4416.
- 599 Cleland E, Chuine I, Menzel A, Mooney H, Schwartz M (2007) Shifting plant phenology in
600 response to global change. *Trends in Ecology & Evolution*, **22**, 357–365.
- 601 Cunningham SJ, Gardner JL, Martin RO (2021) Opportunity costs and the response of
602 birds and mammals to climate warming. *Frontiers in Ecology and the Environment*, **19**,
603 300–307.
- 604 Darlington S, Gooliaff T, Stent P, Hodges K, Ford AT (2025) Southern BC cougar project.
605 Movebank study 1620039803.
- 606 Deb JC, Forbes G, MacLean DA (2020) Modelling the spatial distribution of selected north
607 american woodland mammals under future climate scenarios. *Mammal Review*, **50**, 440–
608 452.
- 609 DeMars CA, Johnson CJ, Dickie M et al. (2023) Incorporating mechanism into conservation
610 actions in an age of multiple and emerging threats: The case of boreal caribou. *Ecosphere*,
611 **14**, e4627.
- 612 DeNicola VL, Mezzini S, Cagnacci F, Fleming CH (2025) Are your data too coarse for speed
613 estimation? Diffusion rates as an alternative measure of animal movement.
- 614 Denny M (2019) Performance in a variable world: Using jensen's inequality to scale up from
615 individuals to populations (ed Domenici P). *Conservation Physiology*, **7**, coz053.
- 616 Dickie M, Serrouya R, McNay RS, Boutin S (2017) Faster and farther: Wolf movement on
617 linear features and implications for hunting behaviour (ed Du Toit J). *Journal of Applied*

- 618 *Ecology*, **54**, 253–263.
- 619 Dickie M, Serrouya R, Avgar T et al. (2022) Resource exploitation efficiency collapses the
620 home range of an apex predator. *Ecology*.
- 621 Dickie M, Serrouya R, Becker M et al. (2024) Habitat alteration or climate: What drives
622 the densities of an invading ungulate? *Global Change Biology*, **30**, e17286.
- 623 Dierauer JR, Allen DM, Whitfield PH (2021) Climate change impacts on snow and stream-
624 flow drought regimes in four ecoregions of british columbia. *Canadian Water Resources
625 Journal / Revue canadienne des ressources hydriques*, **46**, 168–193.
- 626 Diffenbaugh NS, Field CB (2013) Changes in ecologically critical terrestrial climate condi-
627 tions. *Science*, **341**, 486–492.
- 628 Duncan C, Chauvenet ALM, McRae LM, Pettorelli N (2012) Predicting the future impact
629 of droughts on ungulate populations in arid and semi-arid environments (ed Seebacher
630 F). *PLoS ONE*, **7**, e51490.
- 631 Dyer A, Brose U, Berti E, Rosenbaum B, Hirt MR (2023) The travel speeds of large animals
632 are limited by their heat-dissipation capacities (ed Hedenström A). *PLOS Biology*, **21**,
633 e3001820.
- 634 Elmore RD, Carroll JM, Tanner EP, Hovick TJ, Grisham BA, Fuhlendorf SD, Windels
635 SK (2017) Implications of the thermal environment for terrestrial wildlife management.
636 *Wildlife Society Bulletin*, **41**, 183–193.
- 637 Eum H-I, Fajard B, Tang T, Gupta A (2023) Potential changes in climate indices in alberta
638 under projected global warming of 1.5–5 °c. *Journal of Hydrology: Regional Studies*, **47**,
639 101390.
- 640 Fithian W, Hastie T (2013) Finite-sample equivalence in statistical models for presence-only
641 data. *The Annals of Applied Statistics*, **7**.
- 642 Fleming CH, Calabrese JM (2017) A new kernel density estimator for accurate home-range
643 and species-range area estimation (ed Dray S). *Methods in Ecology and Evolution*, **8**,
644 571–579.

- 645 Fleming CH, Calabrese JM (2023) **Ctmm: Continuous-time movement modeling**.
- 646 Fleming CH, Calabrese JM, Mueller T, Olson KA, Leimgruber P, Fagan WF (2014) **From**
647 **fine-scale foraging to home ranges: A semivariance approach to identifying movement**
648 **modes across spatiotemporal scales.** *The American Naturalist*, **183**, E154–E167.
- 649 Ford AT, Noonan MJ, Bollefer K, Gill R, Legebokow C, Serrouya R (2023) **The effects of**
650 **maternal penning on the movement ecology of mountain caribou.** *Animal Conservation*,
651 **26**, 72–85.
- 652 Fuller A, Mitchell D, Maloney SK, Hetem RS (2016) **Towards a mechanistic understanding**
653 **of the responses of large terrestrial mammals to heat and aridity associated with climate**
654 **change.** *Climate Change Responses*, **3**, 10.
- 655 Gerlich HS, Loboda S, Simpson GL, Savage J, Schmidt NM, Holmstrup M, Høye TT (2025)
656 **Species' traits modulate rapid changes in flight time in high-arctic muscid flies under**
657 **climate change.** *Proceedings of the Royal Society B: Biological Sciences*, **292**.
- 658 Giroux A, Ortega Z, Attias N, Desbiez ALJ, Valle D, Börger L, Rodrigues Oliveira-Santos
659 LG (2023) **Activity modulation and selection for forests help giant anteaters to cope with**
660 **temperature changes.** *Animal Behaviour*, **201**, 191–209.
- 661 Glass TW, Breed GA, Robards MD, Williams CT, Kielland K (2021) **Trade-off between pre-**
662 **dation risk and behavioural thermoregulation drives resting behaviour in a cold-adapted**
663 **mesocarnivore.** *Animal Behaviour*, **175**, 163–174.
- 664 Graham V, Baumgartner JB, Beaumont LJ, Esperón-Rodríguez M, Grech A (2019) **Prioritizing**
665 **the protection of climate refugia: Designing a climate-ready protected area network.**
666 *Journal of Environmental Planning and Management*, **62**, 2588–2606.
- 667 Gulland FMD, Baker JD, Howe M et al. (2022) **A review of climate change effects on marine**
668 **mammals in united states waters: Past predictions, observed impacts, current research**
669 **and conservation imperatives.** *Climate Change Ecology*, **3**, 100054.
- 670 Hall LE, Chalfoun AD, Beever EA, Loosen AE (2016) **Microrefuges and the occurrence of**
671 **thermal specialists: Implications for wildlife persistence amidst changing temperatures.**

- 672 *Climate Change Responses*, **3**, 8.
- 673 Hannah L, Flint L, Syphard AD, Moritz MA, Buckley LB, McCullough IM (2014) Fine-
674 grain modeling of species' response to climate change: Holdouts, stepping-stones, and
675 microrefugia. *Trends in Ecology & Evolution*, **29**, 390–397.
- 676 Hegel TM, Mysterud A, Huettmann F, Stenseth NC (2010) Interacting effect of wolves and
677 climate on recruitment in a northern mountain caribou population. *Oikos*, **119**, 1453–
678 1461.
- 679 Hersbach H, Bell B, Berrisford P et al. (2023) ERA5 hourly data on single levels from 1940
680 to present.
- 681 Hetem RS, Fuller A, Maloney SK, Mitchell D (2014) Responses of large mammals to climate
682 change. *Temperature*, **1**, 115–127.
- 683 Hijmans RJ (2023) Terra: Spatial data analysis.
- 684 Hollister J, Shah T, Nowosad J, Robitaille AL, Beck MW, Johnson M (2023) Elevatr: Access
685 elevation data from various APIs.
- 686 Hou R, Chapman CA, Jay O, Guo S, Li B, Raubenheimer D (2020) Cold and hungry:
687 Combined effects of low temperature and resource scarcity on an edge-of-range temperate
688 primate, the golden snub-nose monkey. *Ecography*, **43**, 1672–1682.
- 689 Hua F, Bruijnzeel LA, Meli P et al. (2022) The biodiversity and ecosystem service contrib-
690 utions and trade-offs of forest restoration approaches. *Science*, **376**, 839–844.
- 691 IPCC (2023) Climate change 2021 – the physical science basis: Working group i contribution
692 to the sixth assessment report of the intergovernmental panel on climate change, 1st edn.
693 Cambridge University Press.
- 694 Jahn M, Seebacher F (2022) Variations in cost of transport and their ecological consequences:
695 A review. *Journal of Experimental Biology*, **225**, jeb243646.
- 696 Jeltsch F, Roeleke M, Abdelfattah A et al. (2025) The need for an individual-based global
697 change ecology. *Individual-based Ecology*, **1**, 1–18.
- 698 Jessen C (2001) Temperature regulation in humans and other mammals. Springer Berlin

- 699 Heidelberg, Berlin, Heidelberg.
- 700 Johnson DH (1980) The comparison of usage and availability measurements for evaluating
701 resource preference. *Ecology*, **61**, 65–71.
- 702 Johnson HE, Lewis DL, Verzuh TL, Wallace CF, Much RM, Willmarth LK, Breck SW (2018)
703 Human development and climate affect hibernation in a large carnivore with implications
704 for human–carnivore conflicts (ed Bieber C). *Journal of Applied Ecology*, **55**, 663–672.
- 705 Kacelnik A, Krebs JR, Bernstein C (1992) The ideal free distribution and predator-prey
706 populations. *Trends in Ecology & Evolution*, **7**, 50–55.
- 707 Kays R, Davidson SC, Berger M et al. (2022) The movebank system for studying global
708 animal movement and demography. *Methods in Ecology and Evolution*, **13**, 419–431.
- 709 Kienzle SW (2018) Has it become warmer in alberta? Mapping temperature changes for the
710 period 1950–2010 across alberta, canada. *Canadian Geographies / Géographies canadi-
711 ennes*, **62**, 144–162.
- 712 Kohl MT, Ruth TK, Metz MC, Stahler DR, Smith DW, White PJ, MacNulty DR (2019) Do
713 prey select for vacant hunting domains to minimize a multi-predator threat? (ed Gaillard
714 J). *Ecology Letters*, **22**, 1724–1733.
- 715 Labadie G, Hardy C, Boulanger Y, Vanlandeghem V, Hebblewhite M, Fortin D (2023) Global
716 change risks a threatened species due to alteration of predator–prey dynamics. *Ecosphere*,
717 **14**, e4485.
- 718 Latham ADM, Latham MC, Mccutchen NA, Boutin S (2011) Invading white-tailed deer
719 change wolf–caribou dynamics in northeastern alberta. *The Journal of Wildlife Manage-
720 ment*, **75**, 204–212.
- 721 Leblond M, Dussault C, Ouellet J, St-Laurent M (2016) Caribou avoiding wolves face in-
722 creased predation by bears – caught between scylla and charybdis (ed Singh N). *Journal
723 of Applied Ecology*, **53**, 1078–1087.
- 724 Leclerc M, Leblond M, Le Corre M, Dussault C, Côté SD (2021) Determinants of migra-
725 tion trajectory and movement rate in a long-distance terrestrial mammal (ed Reyna R).

- 726 *Journal of Mammalogy*, **102**, 1342–1352.
- 727 Lessard É, Johnson CJ, St-Laurent M-H (2025) Local weather interacts with human distur-
728 bances to shape the behaviour of boreal caribou across a large climate gradient. *Biodi-
729 versity and Conservation*, **34**, 1115–1138.
- 730 Li Z, Wood SN (2020) Faster model matrix crossproducts for large generalized linear models
731 with discretized covariates. *Statistics and Computing*, **30**, 19–25.
- 732 Livingstone SW, Hughes J, Pither R, Fortin M-J (2023) Integrative adaptive management
733 to address interactions between biological invasions and protected area connectivity: A
734 canadian perspective (ed Smol JP). *FACETS*, **8**, 1–12.
- 735 Loarie SR, Duffy PB, Hamilton H, Asner GP, Field CB, Ackerly DD (2009) The velocity of
736 climate change. *Nature*, **462**, 1052–1055.
- 737 Lochhead KD, Kleynhans EJ, Muhly TB (2022) Linking woodland caribou abundance to
738 forestry disturbance in southern british columbia, canada. *The Journal of Wildlife Man-
739 agement*, **86**, e22149.
- 740 Mahony CR, Wang T, Hamann A, Cannon AJ (2022) A global climate model ensemble
741 for downscaled monthly climate normals over north america. *International Journal of
742 Climatology*, **42**, 5871–5891.
- 743 Martinez-Garcia R, Fleming CH, Seppelt R, Fagan WF, Calabrese JM (2020) How range res-
744 idency and long-range perception change encounter rates. *Journal of Theoretical Biology*,
745 **498**, 110267.
- 746 Matthiopoulos J (2022) Defining, estimating, and understanding the fundamental niches of
747 complex animals in heterogeneous environments. *Ecological Monographs*, **92**, e1545.
- 748 McCabe JD, Clare JD, Miller TA et al. (2021) Resource selection functions based on hierar-
749 chical generalized additive models provide new insights into individual animal variation
750 and species distributions. *Ecography*, **44**, 1756–1768.
- 751 McCain CM (2019) Assessing the risks to united states and canadian mammals caused
752 by climate change using a trait-mediated model (ed Heske E). *Journal of Mammalogy*,

- 753 gyz155.
- 754 McCain CM, King SRB (2014) Body size and activity times mediate mammalian responses
755 to climate change. *Global Change Biology*, **20**, 1760–1769.
- 756 McLellan ML, McLellan BN (2015) Effect of season and high ambient temperature on activity
757 levels and patterns of grizzly bears (*ursus arctos*) (ed Crocker DE). *PLOS ONE*, **10**,
758 e0117734.
- 759 McNab BK (1970) Body weight and the energetics of temperature regulation. *Journal of*
760 *Experimental Biology*, **53**, 329–348.
- 761 Melin M, Matala J, Mehtätalo L, Pusenius J, Packalen T (2023) The effect of snow depth
762 on movement rates of GPS-collared moose. *European Journal of Wildlife Research*, **69**,
763 21.
- 764 Merkle JA, Monteith KL, Aikens EO et al. (2016) Large herbivores surf waves of green-up
765 during spring. *Proceedings of the Royal Society B: Biological Sciences*, **283**, 20160456.
- 766 Mezzini S, Fleming CH, Medici EP, Noonan MJ (2025) How resource abundance and resource
767 stochasticity affect organisms' range sizes. *Movement Ecology*, **13**, 20.
- 768 Michalak JL, Lawler JJ, Roberts DR, Carroll C (2018) Distribution and protection of climatic
769 refugia in north america. *Conservation Biology*, **32**, 1414–1425.
- 770 Mota-Rojas D, Titto CG, Orihuela A et al. (2021) Physiological and behavioral mechanisms
771 of thermoregulation in mammals. *Animals*, **11**, 1733.
- 772 Muff S, Signer J, Fieberg J (2020) Accounting for individual-specific variation in habi-
773 tation-selection studies: Efficient estimation of mixed-effects models using bayesian or fre-
774 quentist computation (ed Wal EV). *Journal of Animal Ecology*, **89**, 80–92.
- 775 Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, Smouse PE (2008) A
776 movement ecology paradigm for unifying organismal movement research. *Proceedings of*
777 *the National Academy of Sciences*, **105**, 19052–19059.
- 778 Nathan R, Monk CT, Arlinghaus R et al. (2022) Big-data approaches lead to an increased
779 understanding of the ecology of animal movement. *Science*, **375**, eabg1780.

- 780 Newediuk L, Richardson ES, Biddlecombe BA et al. (2024) Climate change, age acceleration,
781 and the erosion of fitness in polar bears.
- 782 Noonan MJ, Fleming CH, Akre TS et al. (2019a) Scale-insensitive estimation of speed and
783 distance traveled from animal tracking data. *Movement Ecology*, **7**, 35.
- 784 Noonan MJ, Tucker MA, Fleming CH et al. (2019b) A comprehensive analysis of autocor-
785 relation and bias in home range estimation. *Ecological Monographs*, **89**, e01344.
- 786 Parisien M-A, Barber QE, Bourbonnais ML et al. (2023) Abrupt, climate-induced increase in
787 wildfires in british columbia since the mid-2000s. *Communications Earth & Environment*,
788 **4**, 309.
- 789 Pedersen EJ, Miller DL, Simpson GL, Ross N (2019) Hierarchical generalized additive models
790 in ecology: An introduction with mgcv. *PeerJ*, **7**, e6876.
- 791 Pedersen SH, Bentzen TW, Reinking AK et al. (2021) Quantifying effects of snow depth on
792 caribou winter range selection and movement in arctic alaska. *Movement Ecology*, **9**, 48.
- 793 Péron G, Fleming CH, De Paula RC, Mitchell N, Strohbach M, Leimgruber P, Calabrese JM
794 (2017) Periodic continuous-time movement models uncover behavioral changes of wild
795 canids along anthropization gradients. *Ecological Monographs*, **87**, 442–456.
- 796 Peterson AN, Soto AP, McHenry MJ (2021) Pursuit and evasion strategies in the predator–
797 prey interactions of fishes. *Integrative and Comparative Biology*, **61**, 668–680.
- 798 Pigeon KE, Stenhouse G, Côté SD (2016) Drivers of hibernation: Linking food and weather
799 to denning behaviour of grizzly bears. *Behavioral Ecology and Sociobiology*, **70**, 1745–
800 1754.
- 801 Polazzo F, Limberger R, Pennekamp F, Ross SRP-J, Simpson GL, Petchey OL (2024) Mea-
802 suring the response diversity of ecological communities experiencing multifarious envi-
803 ronmental change. *Global Change Biology*, **30**, e17594.
- 804 Post E, Forchhammer MC (2008) Climate change reduces reproductive success of an arctic
805 herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society B:*
806 *Biological Sciences*, **363**, 2367–2373.

- 807 Powers DR, Langland KM, Wethington SM, Powers SD, Graham CH, Tobalske BW (2017)
808 Hovering in the heat: Effects of environmental temperature on heat regulation in foraging
809 hummingbirds. *Royal Society Open Science*, **4**, 171056.
- 810 R Core Team (2024) *R: A language and environment for statistical computing*. R Foundation
811 for Statistical Computing, Vienna, Austria.
- 812 Ratnayake HU, Kearney MR, Govekar P, Karoly D, Welbergen JA (2019) Forecasting wildlife
813 die-offs from extreme heat events. *Animal Conservation*, **22**, 386–395.
- 814 Riahi K, Vuuren DP van, Kriegler E et al. (2017) The shared socioeconomic pathways and
815 their energy, land use, and greenhouse gas emissions implications: An overview. *Global
816 Environmental Change*, **42**, 153–168.
- 817 Rice WL (2022) The conspicuous consumption of wilderness, or leisure lost in the wilderness.
818 *World Leisure Journal*, **64**, 451–468.
- 819 Rigby RA, Stasinopoulos DM (2005) Generalized additive models for location, scale and
820 shape (with discussion). *Journal of the Royal Statistical Society: Series C (Applied
821 Statistics)*, **54**, 507–554.
- 822 Sawyer H, Merkle JA, Middleton AD, Dwinnell SPH, Monteith KL (2019) Migratory plastic-
823 ity is not ubiquitous among large herbivores (ed Chapman J). *Journal of Animal Ecology*,
824 **88**, 450–460.
- 825 Schmidt NM, Grøndahl C, Evans AL et al. (2020) On the interplay between hypothermia
826 and reproduction in a high arctic ungulate. *Scientific Reports*, **10**, 1514.
- 827 Schwerdt L, De Villalobos AE, Ferretti N (2024) Ecological niche modelling and thermal pa-
828 rameters to assess the prevalence of an endemic tarantula: The endurance of grammostola
829 vachoni schiapelli & gerschman, 1961. *Journal of Insect Conservation*.
- 830 Sherwood SC, Huber M (2010) An adaptability limit to climate change due to heat stress.
831 *Proceedings of the National Academy of Sciences*, **107**, 9552–9555.
- 832 Sih A, Ferrari MCO, Harris DJ (2011) Evolution and behavioural responses to human-
833 induced rapid environmental change: Behaviour and evolution. *Evolutionary Applica-*

- 834 *tions*, **4**, 367–387.
- 835 Silva I, Fleming CH, Noonan MJ, Fagan WF, Calabrese JM (2025) Too few, too many, or
836 just right? Optimizing sample sizes for population-level inferences in animal tracking
837 projects.
- 838 Simmons BA, Nolte C, McGowan J (2021) Tough questions for the “30 × 30” conservation
839 agenda. *Frontiers in Ecology and the Environment*, **19**, 322–323.
- 840 Simpson GL (2018) Modelling palaeoecological time series using generalised additive models.
841 *Frontiers in Ecology and Evolution*, **6**, 149.
- 842 Simpson GL (2025) Method: Using generalized additive models in the animal sciences.
- 843 Slatyer RA, Umbers KDL, Arnold PA (2022) Ecological responses to variation in seasonal
844 snow cover. *Conservation Biology*, **36**, e13727.
- 845 Smith BJ, MacNulty DR, Stahler DR, Smith DW, Avgar T (2023) Density-dependent habitat
846 selection alters drivers of population distribution in northern yellowstone elk. *Ecology*
847 *Letters*, **26**, 245–256.
- 848 Sobie SR, Zwiers FW, Curry CL (2021) Climate model projections for canada: A comparison
849 of CMIP5 and CMIP6. *Atmosphere-Ocean*, **59**, 269–284.
- 850 Speed JDM, Austrheim G, Hester AJ, Mysterud A (2012) Elevational advance of alpine plant
851 communities is buffered by herbivory (ed Ejrnaes R). *Journal of Vegetation Science*, **23**,
852 617–625.
- 853 Stasinopoulos MD, Rigby RA (2007) Generalized additive models for location scale and
854 shape (GAMLSS) in r. *Journal of Statistical Software*, **23**.
- 855 Stien A, Ims RA, Albon SD et al. (2012) Congruent responses to weather variability in high
856 arctic herbivores. *Biology Letters*, **8**, 1002–1005.
- 857 Stralberg D, Carroll C, Nielsen SE (2020) Toward a climate-informed north american pro-
858 tected areas network: Incorporating climate-change refugia and corridors in conservation
859 planning. *Conservation Letters*, **13**, e12712.
- 860 Sullender BK, Cunningham CX, Lundquist JD, Prugh LR (2023) Defining the danger zone:

- 861 Critical snow properties for predator–prey interactions. *Oikos*, **2023**, e09925.
- 862 Tan M, Zhang S, Stevens M, Li D, Tan EJ (2024) Antipredator defences in motion: Animals
863 reduce predation risks by concealing or misleading motion signals. *Biological Reviews*,
864 **99**, 778–796.
- 865 Taylor CR, Heglund NC, Maloiy GMO (1982) Energetics and mechanics of terrestrial loco-
866 motion. I. Metabolic energy consumption as a function of speed and body size in birds
867 and mammals. *Journal of Experimental Biology*, **97**, 1–21.
- 868 Thompson V, Kennedy-Asser AT, Vosper E et al. (2022) The 2021 western north america
869 heat wave among the most extreme events ever recorded globally. *Science Advances*, **8**,
870 eabm6860.
- 871 Tórrez-Herrera LL, Davis GH, Crofoot MC (2020) Do monkeys avoid areas of home range
872 overlap because they are dangerous? A test of the risk hypothesis in white-faced capuchin
873 monkeys (*cebus capucinus*). *International Journal of Primatology*, **41**, 246–264.
- 874 Tuanmu M, Jetz W (2014) A global 1-km consensus land-cover product for biodiversity and
875 ecosystem modelling. *Global Ecology and Biogeography*, **23**, 1031–1045.
- 876 Turner NJ, Clifton H (2009) “It’s so different today”: Climate change and indigenous lifeways
877 in british columbia, canada. *Global Environmental Change*, **19**, 180–190.
- 878 Tysor CS (2025) Countergradient variation in lodgepole pine flowering phenology.
- 879 Verzuh TL, Rogers SA, Mathewson PD et al. (2023) Behavioural responses of a large,
880 heat-sensitive mammal to climatic variation at multiple spatial scales. *Journal of Animal
881 Ecology*, **92**, 619–634.
- 882 Walker WH, Meléndez-Fernández OH, Nelson RJ, Reiter RJ (2019) Global climate change
883 and invariable photoperiods: A mismatch that jeopardizes animal fitness. *Ecology and
884 Evolution*, **9**, 10044–10054.
- 885 Wang T, Hamann A, Spittlehouse D, Carroll C (2016) Locally downscaled and spatially
886 customizable climate data for historical and future periods for north america (ed Álvarez
887 I). *PLOS ONE*, **11**, e0156720.

- 888 Wang W, Wang X, Flannigan MD et al. (2025) Canadian forests are more conducive to
889 high-severity fires in recent decades. *Science*, **387**, 91–97.
- 890 Wells CP, Barbier R, Nelson S, Kanaziz R, Aubry LM (2022) Life history consequences of
891 climate change in hibernating mammals: A review. *Ecography*, **2022**, e06056.
- 892 Weststrate DK, Chhen A, Mezzini S, Safford K, Noonan MJ (2024) How climate change
893 and population growth will shape attendance and human-wildlife interactions at british
894 columbia parks. *Journal of Sustainable Tourism*, 1–15.
- 895 White KS (2025) Population ecology of mountain goats in relation to climate, weather and
896 snow avalanches. PhD thesis, University of Victoria.
- 897 White KS, Gregovich DP (2017) Mountain goat resource selection in relation to min-
898 ing-related disturbance. *Wildlife Biology*, **2017**, 1–12.
- 899 White KS, Cadsand B, Côté SD et al. (2025) Mountain sentinels in a changing world:
900 Review and conservation implications of weather and climate effects on mountain goats
901 (*Oreamnos americanus*). *Global Ecology and Conservation*, **57**, e03364.
- 902 Whitman E, Parks SA, Holsinger LM, Parisien M-A (2022) Climate-induced fire regime
903 amplification in alberta, canada. *Environmental Research Letters*, **17**, 055003.
- 904 Whittington J, Hebblewhite M, DeCesare NJ, Neufeld L, Bradley M, Wilmshurst J, Musiani
905 M (2011) Caribou encounters with wolves increase near roads and trails: A time-to-event
906 approach. *Journal of Applied Ecology*, **48**, 1535–1542.
- 907 Williams JE, Blois JL (2018) Range shifts in response to past and future climate change:
908 Can climate velocities and species' dispersal capabilities explain variation in mammalian
909 range shifts? *Journal of Biogeography*, **45**, 2175–2189.
- 910 Winter VA, Smith BJ, Berger DJ et al. (2024) Forecasting animal distribution through in-
911 dividual habitat selection: Insights for population inference and transferable predictions.
912 *Ecography*, e07225.
- 913 Wood SN (2017) Generalized additive models: An introduction with r, Second edition edn.
914 CRC Press/Taylor & Francis Group, Boca Raton.

- 915 Wood SN, Goude Y, Shaw S (2015) Generalized additive models for large data sets. *Journal*
916 *of the Royal Statistical Society: Series C (Applied Statistics)*, **64**, 139–155.
- 917 Wood SN, Li Z, Shaddick G, Augustin NH (2017) Generalized additive models for gigadata:
918 Modeling the u.k. Black smoke network daily data. *Journal of the American Statistical
919 Association*, **112**, 1199–1210.
- 920 Woo-Durand C, Matte J-M, Cuddihy G, McGourdji CL, Venter O, Grant JWA (2020) In-
921 creasing importance of climate change and other threats to at-risk species in canada.
922 *Environmental Reviews*, **28**, 449–456.
- 923 Xu W, Barker K, Shawler A et al. (2021) The plasticity of ungulate migration in a changing
924 world. *Ecology*, **102**, e03293.
- 925 Zhang J, Huang S, He F (2015) Half-century evidence from western canada shows forest
926 dynamics are primarily driven by competition followed by climate. *Proceedings of the
927 National Academy of Sciences*, **112**, 4009–4014.
- 928 Zhang X, Zhou T, Zhang W et al. (2023) Increased impact of heat domes on 2021-like heat
929 extremes in north america under global warming. *Nature Communications*, **14**, 1690.