

How resource abundance and stochasticity affect organisms' space-use requirements

Article type: Major article

Words in abstract: 199

Words in main text: 5265

Figures: 6

Tables: 0

References: 95

Appendices: 3

Key words: energetics, energetic landscape, environmental stochasticity, space use, spatial needs, ctmm

To do:

- make sure appendices flow (added stuff from main text)
- check number of references (updated on 2024-03-18)

1 Abstract

2 The amount of space organisms use is thought to be tightly linked to the availability of
3 resources within their habitats, such that organisms living in productive habitats generally
4 require less space than those in resource-poor habitats. This hypothesis has widespread em-
5 pirical support, but existing studies have focused primarily on responses to the mean amount
6 of resources, while responses to the variance around the mean are still largely unknown. This
7 is not a trivial oversight. Organisms adjust to variable environmental conditions, so failing
8 to consider the effects of resource (un)predictability can result in a limited understanding
9 of organisms' space-use requirements, challenging ecological theory and applied conservation
10 alike. In this study, we leverage the available literature to provide a unifying framework and
11 hypotheses for the effect of mean and variance in resources on organisms' space use. Next,
12 we use simulated movement data to demonstrate how the combined effects of mean and
13 variance in resource abundance interact to shape predictable patterns in space use. Finally,
14 we use real-world tracking data on a lowland tapir (*Tapirus terrestris*) from the Brazilian
15 Cerrado to show how this framework can be applied to better understand the movement
16 ecology of free-ranging animals.

¹⁷ **Introduction**

¹⁸ The amount of resources an organism is able to access is a strong determinant of its odds
¹⁹ of survival and reproduction. Resource limitations can cause individuals to experience a
²⁰ negative energetic balance, which can then result in lower fitness (Le Bot et al. 2019; Hou
²¹ et al. 2020), altered physiology (Wessling et al. 2018; Le Bot et al. 2019; Rocha et al. 2021;
²² Dai Pra et al. 2022), lower chance of reproduction (Douglas and Pearce-Higgins 2014; Le
²³ Bot et al. 2019; Schmidt et al. 2020; Stefanescu et al. 2021), and even death (Foley et
²⁴ al. 2008; Berger et al. 2018), along with changes to community structure (Burson et al.
²⁵ 2018; Ghislandi et al. 2018; Haney and Siepielski 2018; Riotte-Lambert and Matthiopoulos
²⁶ 2020). Thus, many organisms adapt their behaviors in response to changes in local resource
²⁷ abundance to ensure their needs are met. Movement represents one of the most readily
²⁸ available behaviors that species can adjust to optimize their resource use (Charnov 1976;
²⁹ Kacelnik et al. 1992).

³⁰ The relationship between organisms' movement and resource abundance has been of
³¹ interest to biologists for nearly a century. Burt (1943) considered the search for food as
³² the primary driver for movement within an organism's home range. Three decades after,
³³ Southwood (1977) suggested that change in resource abundance is a strong determinant of
³⁴ how organisms decide where to live and when to reproduce. Two years later, Harestad and
³⁵ Bunnel (1979) proposed that the simplest relationship between resource abundance and an
³⁶ organism's home-range size is

$$H = C/R, \quad (1)$$

³⁷ where H is the organism's home-range size, C is the organism's resource consumption (kcal
³⁸ day^{-1}), and R is the resources the organism can access ($\text{kcal day}^{-1} \text{ unit area}^{-1}$). Harestad
³⁹ and Bunnel's model is simple to conceptualize and allows for testable predictions, but it fails
⁴⁰ to account for the nonlinear relationship required to avoid negative home range estimates

41 (e.g., fig. 2 in the work of Bista et al. 2022). Many researchers have since demonstrated that
42 organisms adapt their home ranges in response to resources abundance, but studies often fail
43 to build upon previous work. Instead, results are typically reported as independent, novel
44 findings. Thus, there remains a need for a clear hypothesis for the effects of resource abun-
45 dance on organisms' space-use requirements. Additionally, while much work has been done
46 on estimating organisms' responses to average resource abundance, there is little information
47 on how they respond to unpredictable changes in resources (but see: Stephens and Charnov
48 1982; Rizzuto et al. 2021).

49 Here, we refer to a location's average amount of resources as "resource abundance",
50 while we use the phrase "resource stochasticity" to indicate the variability in resources after
51 accounting for changes in the mean. We argue that, on its own, a habitat's long-term resource
52 abundance is not sufficient to assess the habitat's quality, nor make predictions about how
53 much space an organism might use. For instance, herbivores in a grassland with relatively
54 low but constant forage availability (low mean and variance) will adopt different behaviors
55 and adaptations from herbivores in a desert with equally scarce forage but rare, sudden, and
56 strong pulses of resources (low mean and high stochasticity). In the grassland, an animal
57 may require a large but constant home range as it moves between locations in search of food
58 (Teitelbaum and Mueller 2019), while an animal in the desert may switch between dispersal
59 in search for high-resource patches and short-term range residency whinith patches (Charnov
60 1976). Although it may be possible for both habitats to have the same long-term average
61 resource abundance, the differences in resource unpredictability result in different selection of
62 movement and life history strategies. Although resource unpredictability (and seasonality)
63 may ecrease organisms' fitness and a landscape's energetic balance (Chevin et al. 2010),
64 there is still limited empirical evidence to support this hypothesis (but see: Herfindal et al.
65 2005; Nilsen et al. 2005; Rizzuto et al. 2021).

66 Here, we illustrate how organisms' space use depends on both the abundance and unpre-
67 dictability of resources. First, we set the theoretical background necessary for the successive

68 sections by introducing key concepts and notation. Next, we provide a review of the effects
69 of resource abundance on organisms' space use while suggesting a unifying hypothesis. Af-
70 terwards, we provide a review of the effects of resource stochasticity on organisms' space use
71 while suggesting a second unifying hypothesis. Subsequently, we demonstrate the power of
72 these two hypotheses using quantitative, simulated responses to changes in resource abun-
73 dance and unpredictability. Finally, we demonstrate how this framework can be used in
74 practice to describe the movement ecology of a lowland tapir (*Tapirus terrestris*) from the
75 Brazilian Cerrado (Medici et al. 2022). Results from the simulations and empirical example
76 are presented using a fully transparent approach that allows researchers to replicate the work
77 and apply the methods to their own tracking data.

78 Resources as a random variable

79 Resources are often unpredictable (and difficult to quantify), since they depend on various
80 factors which cannot be accounted for easily, including climate (Lindstedt and Boyce 1985;
81 Morellet et al. 2013; Schmidt et al. 2020), weather (Morellet et al. 2013; Fjelldal et al.
82 2021), competitive pressure (Rich et al. 2012; Tórrez-Herrera et al. 2020), and differences
83 in energetics at the individual (Schmidt et al. 2020) and species level (Jetz et al. 2004).
84 Thus, let the random variable R indicate the amount of resources at a given point in space
85 and time. Quantifying resources as a numerical random variable, as opposed to using *ad hoc*
86 qualitative descriptions, provides us with the capacity to leverage techniques from probability
87 theory and statistics. We assume readers are familiar with fundamental statistical concepts
88 (e.g., random variables, probability distributions, expected value and variance of a random
89 variable), but we define and explain the necessary concepts in Appendix A.

90 Effects of resource abundance, $E(R)$

91 While organisms' needs vary greatly between taxonomic groups, some needs are essential
92 for most species for survival and reproduction. All heterotrophic organisms require sources

of chemical energy (i.e., food), water, and various limiting nutrients to survive, grow, and reproduce (Harvey and Clutton-Brock 1981; Baldwin and Bywater 1984; Reich 2001). Failing to acquire sufficient resources can result in lower fitness, behavioral changes, physiological changes or damage, inability to reproduce, and death. Motile organisms can move to new locations in search of resources (Charnov 1976; Brown et al. 1999), but they must also account for a higher metabolism and movement costs (Taylor et al. 1982). The hypothesis that space-use requirements decrease with resource abundance, $E(R)$, is commonly accepted and well supported, but many studies assume a linear relationship (e.g., Harestad and Bunnel 1979; Relyea et al. 2000; McClintic et al. 2014; Rizzuto et al. 2021; Bista et al. 2022; Bradsworth et al. 2022). This is problematic because a linear relationship can lead to questionable estimates and problematic biases, such as negative or excessively small home ranges (e.g., fig. 2 in the work of Bista et al. 2022). Conceptually, the relationship between space-use requirements and $E(R)$ must be nonlinear, since there are limits to how much space an organism is able to explore in its lifetime and a minimum amount of space it requires to survive (which is necessarily greater than zero – see: Lucherini and Lovari 1996; Herfindal et al. 2005; Nilsen et al. 2005; Simcharoen et al. 2014; Watson 2020, and contrast them to the estimates based on linear models listed above).

Fig. 1 illustrates our first of two hypotheses: When $E(R)$ is high, we expect organisms' spatial requirements to be relatively small and near the smallest amount of space required to survive (e.g., Relyea et al. 2000; Herfindal et al. 2005; Nilsen et al. 2005). At lower values of $E(R)$, we expect organisms' spatial requirements to increase nonlinearly, since low values of $E(R)$ force organisms to expand their home ranges (Lucherini and Lovari 1996; Relyea et al. 2000; Herfindal et al. 2005; Nilsen et al. 2005; Bista et al. 2022), migrate to better locations (Samarra et al. 2017; Middleton et al. 2018; Geremia et al. 2019), or move nomadically (Singh et al. 2012; Polansky et al. 2015; Nandintsetseg et al. 2019; Teitelbaum and Mueller 2019). It is unclear when organisms switch from range residency to migration or nomadism (or vice-versa), but understanding the connection between these




Figure 1: Hypothesized space-use requirements of an organism as a function of resource abundance ($E(R)$). We expect low values of $E(R)$ to result in high space-use requirements as organisms are forced to explore large areas to collect the resources they require to survive, whether they be range-resident or nomadic. As $E(R)$ increases, space-use requirements should decrease nonlinearly until they reach the minimum amount of space required by the organism to survive. Note that the relationship between $E(R)$ and space-use requirements cannot be linear because it would require space-use requirements to be negative for high values of $E(R)$.

types of movement is important for quantifying the effect of resource abundance on organisms' space-use requirements and when an organism may choose to migrate or disperse rather than remaining range-resident (mammals: Teitelbaum et al. 2015; moose, *Alces alces*: Singh et al. 2012; eagles, *Haliaeetus leucocephalus*: Wheat et al. 2017; Poessel et al. 2022; lesser flamingos, *Phoeniconaias minor*: Pretorius et al. 2020). Still, species-level changes in movement behavior are more likely to occur over evolutionary timescales than over an organism's lifespan, since larger ranges require greater vagility, which, in turn, is facilitated by morphological features such as hinged joints and elongated limbs (Andersson 2004; Jetz et al. 2004; Samuels et al. 2013; Hirt et al. 2017).

Consider the scenario where $E(R) = \mu(t)$ changes over time but $\text{Var}(R) = \sigma^2 > 0$ is constant over time and space (fig. A1). Since R is random, an organism that moves in the landscape will not find the same R at different time points or throughout the habitat. However, if $\mu(t)$ changes repetitively and regularly over time (e.g. peaks each spring and is

¹³³ lowest in winter), an organism may learn to predict times of high or low $E(R)$ (Samarra et
¹³⁴ al. 2017; Abrahms et al. 2019; Geremia et al. 2019; e.g., Falcón-Cortés et al. 2021) but will
¹³⁵ still not be able to predict R exactly.

¹³⁶ **Effects of resource stochasticity, $\text{Var}(R)$**

¹³⁷ The example illustrated in the section above can occasionally be a useful simplification of
¹³⁸ relatively stable environments or when data is scarce, but it likely not realistic, since $\text{Var}(R)$
¹³⁹ often also changes over time. Periods of scarcity (e.g., winter, droughts) not only have
¹⁴⁰ lower $E(R)$ but also have narrower ranges of R . In contrast, periods of high $E(R)$ (e.g.,
¹⁴¹ spring green-up) also often tend to have higher $\text{Var}(R)$ because R tends to vary across wider
¹⁴² ranges (e.g., due to high spatial variability, or high R followed by low R after grazing).
¹⁴³ When both $E(R)$ and $\text{Var}(R)$ change over time (fig. A2), predicting R becomes more com-
¹⁴⁴ plex, since disentangling changes in $E(R)$ and $\text{Var}(R)$ is not simple (Steixner-Kumar and
¹⁴⁵ Gläscher 2020). Statistically, this is because the more change one attributes to $\mu(t)$ (i.e.,
¹⁴⁶ the wigglier it is), the smaller $\sigma^2(t)$ becomes. Conversely, the smoother $\mu(t)$ is, the larger
¹⁴⁷ $\sigma^2(t)$ becomes. Biologically, this is important because an organism's perception scale (Levin
¹⁴⁸ 1992) determines whether it sees a change in R as a trend in $E(R)$ or as a stochastic event
¹⁴⁹ (i.e., due to $\text{Var}(R)$). An organism's perception of changes in R will also depend strongly
¹⁵⁰ on its cognitive capacities and memory (Foley et al. 2008; Fagan et al. 2013; Mueller et
¹⁵¹ al. 2013; Abrahms et al. 2019; Falcón-Cortés et al. 2021). Whether, an organism is able to
¹⁵² predict trends in $\sigma^2(t)$ or not, environmental variability is thought to reduce a landscape's
¹⁵³ energetic balance (Chevin et al. 2010), which, in turn, decreases organisms' fitness (e.g.,
¹⁵⁴ Berger et al. 2018) and increases their space-use requirements. While this is true for both
¹⁵⁵ predictable and unpredictable stochasticity, extreme and rare events are more likely to have
¹⁵⁶ a stronger effect due to their unpredictability and magnitude (Logares and Nuñez 2012;
¹⁵⁷ Anderson et al. 2017). A few recent studies support these hypotheses (Chevin et al. 2010;
¹⁵⁸ Morellet et al. 2013; Nandintsetseg et al. 2019; Riotte-Lambert and Matthiopoulos 2020),

159 but many of them are limited in geographic and taxonomic scales, so the extent to which
160 these preliminary findings can be generalized is still limited. Thus, there remains a need for
161 developing a more complete understanding of how organisms' space-use requirements change
162 with environmental stochasticity.

163 Similarly to $E(R)$, we hypothesize $\text{Var}(R)$ to have a nonlinear effect on organisms' space-
164 use requirements. When $\text{Var}(R)$ is low enough that R is relatively predictable, we do not
165 expect changes in $\text{Var}(R)$ to have a noticeable effect. But as resources become increasingly
166 unpredictable, we expect space-use requirements to increase progressively faster (fig. 2) be-
167 cause: (1) as $\text{Var}(R)$ increases, the chances of finding low R increase superlinearly, (2)
168 stochastic environments tend to be less productive (Chevin et al. 2010), (3) stochasticity
169 reduces an organism's ability to specialize and reduce competition for R (Levins 1974), and
170 (4) the added movement required to search for food increases organisms' energetic require-
171 ments. If resources remain highly unpredictable over long periods of time (e.g., multiple
172 lifespans), organisms may evolve or develop new behaviors (such as nomadism, dispersal,
173 and migration) or adaptations (such as increased fat storage or food caching). If changes in
174 $\sigma^2(t)$ are sufficiently predictable, organisms may learn to anticipate and prepare for periods
175 of greater stochasticity by pre-emptively caching food, migrating, or relying on alternative
176 food sources during stochastic events (e.g., the seasonal arrival of competitors).

177 Interactive effects of $E(R)$ and $\text{Var}(R)$

178 We have provided the case for why both $E(R)$ and $\text{Var}(R)$ affect organisms' space-use require-
179 ments, but we presented the two parameters as independent drivers of movement. However,
180 organisms may respond to changes in $\text{Var}(R)$ more when resources are scarce than when they
181 are abundant. Consequently, models estimating the effects of R on organism's movement
182 should account for not only the marginal effects of $E(R)$ and $\text{Var}(R)$ but also their interac-
183 tive effects. A highly unpredictable habitat may be very inhospitable if resources are poor,
184 but $\text{Var}(R)$ may have little effect if resources are stochastic but always abundant. Thus, we




Figure 2: Hypothesized space-use requirements of an organism as a function of resource stochasticity ($\text{Var}(R)$). We expect low values of $\text{Var}(R)$ to result in low space-use requirements as organisms are able to depend on relatively predictable resources. As $\text{Var}(R)$ increases, space-use requirements should increase nonlinearly, whether this results in an expansion of the home range (in the case of range-resident organisms) or a switch to dispersal, nomadism, or migration. Note that the relationship between $\text{Var}(R)$ and space-use requirements cannot be linear because it would require space-use requirements to be negative for low values of $\text{Var}(R)$.

185 expect $\text{Var}(R)$ to have a stronger effect on space-use requirements when $E(R)$ is low, and
 186 less of an effect when $E(R)$ is high. We explore this more in the following section.

187 **Simulating responses to $E(R)$ and $\text{Var}(R)$**

188 To support our hypotheses of how organisms' space use is affected by $E(R)$, $\text{Var}(R)$, and the
 189 interaction effect of $E(R)$ and $\text{Var}(R)$, we present the results from a series of quantitative
 190 simulations. To start, we used the `ctmm` package (Fleming and Calabrese 2021) for `R` (`R`
 191 Core Team 2023) to generate 200 tracks (see Appendix B for sensitivity analyses) from an
 192 Integrated Ornstein-Uhlenbeck movement model (IOU model, see Gurarie et al. 2017). The
 193 IOU model's correlated velocity produces realistic tracks with directional persistence despite
 194 the tracks being discrete samples of continuous-time movement processes. However, unlike
 195 Ornstein-Uhlenbeck (OU) and Ornstein-Uhlenbeck Foraging (OUF) models, IOU models do
 196 not produce spatially stationary movement, so the organism is not range-resident. Con-

197 sequentially, each track is spatially unrestricted and can be interpreted as purely exploratory
198 movement.

199 Each of the 200 tracks were placed on a raster with common starting point $\langle 0, 0 \rangle$ and
200 sufficient time between tracks to be independent of one another (other than the starting
201 point). Each time the track moved to a new cell, the organism collected resources R sampled
202 from a Gamma distribution. The mean and variance of the distribution were defined by
203 deterministic functions $\mu(t)$ and $\sigma^2(t)$ (orange and blue lines in fig. 3). The value of t
204 was constant within each set of 200 tracks, so the time spent moving by the organism in
205 each track did not affect the distribution R was sampled from. Tracks were truncated once
206 the organism reached satiety, and the organism was given enough time to return to $\langle 0, 0 \rangle$
207 independently from the following track. Finally, as each set of 200 truncated was now
208 spatially restricted and centered around $\langle 0, 0 \rangle$, the set of tracks was modeled to calculate
209 the 95% utilization distribution from an OUF model and Autocorrelated Kernel Density
210 Estimate. We designed the simulations presented here to estimate the effects of $E(R)$ and
211 $\text{Var}(R)$ in simplistic environments. Consequently, we made the following four assumptions:
212 (1) $E(R)$ and $\text{Var}(R)$ are constant over space and short periods of time (i.e., within each set
213 of 200 tracks, but not over t). (2) Organisms move randomly in the environment with no
214 energetic costs. (3) Organisms have no perceptive range or memory, so they are unable to
215 detect, learn, or predict where resources are abundant (high $E(R)$) or reliable (low $\text{Var}(R)$)
216 over time. (4) Resources do not deplete, and there is no competition or predator avoidance.
217 Additional information is provided in Appendix B.

218 Fig. 3 shows how simulated space-use requirements responded to changes in $\mu(t)$ and
219 $\sigma^2(t)$ in scenarios where both functions can remain constant, increase linearly, oscillate
220 cyclically, drift stochastically, or change erratically. The top row shows how space-use re-
221 quirements vary for different trends in $\mu(t)$ while $\text{Var}(R)$ remains constant (like in fig. A1).
222 As $E(R)$ increases at a constant slope (linear $\mu(t)$) H decreases nonlinearly, with larger
223 changes when $E(R)$ is low, until it approaches the minimum size required by the organism.




Figure 3: Simulated home range sizes, H , of an organism living in habitats where the mean and variance in resources are constant, linearly increasing, cyclical, drifting, or erratic over time (and homogenous over space). Note how H decreases nonlinearly as $\mu(t)$ increases and increases nonlinearily as $\sigma^2(t)$ increases. Additionally, the variance in H is higher when $\mu(t)$ is lower or $\sigma^2(t)$ is higher, and changes in $\sigma^2(t)$ have greater impacts when $\mu(t)$ is low.

224 Although less evident, note how the noise in the lines also decreases as $E(R)$ increases.

225 The leftmost column of fig. 3 illustrates the effects of $\text{Var}(R)$ on the simulated space-use
 226 requirements while $E(R)$ remains constant. Overall, both mean H and the variance around
 227 it increase with $\sigma^2(t)$ (most visible with constant $E(R)$ and linear $\text{Var}(R)$). This is because,
 228 similarly to resource-poor periods, times of greater stochasticity require the organism to
 229 move over larger areas for longer periods of time. This results in a greater uncertainty in
 230 how much time and space the organism will require to reach satiety, or indeed whether
 231 an organism living in highly stochastic environments can even reach satiety within a finite
 232 amount of time.

233 The remaining panels in fig. 3 illustrate $E(R)$ and $\text{Var}(R)$ jointly affect H and how
 234 complex the effects can be. Since $E(R)$ and $\text{Var}(R)$ have opposite effects on H , disentangling
 235 the effects can be particularly difficult when both parameters change in a correlated manner
 236 (e.g., linear $E(R)$ and $\text{Var}(R)$). When both $E(R)$ and $\text{Var}(R)$ increase linearly, H initially

increases since the effect of $\text{Var}(R)$ is stronger, but then decreases as the effect of $E(R)$ begins to dominate. Difficulties in disentangling the two effects are explored in greater depth in the case study in the following section.

Although the temporal trends in fig. 3 are complex and the effects of $E(R)$ and $\text{Var}(R)$ can be hard to disentangle, two simple relationships emerge when H is shown as a function of either $E(R)$ or $\text{Var}(R)$, rather than time (panels A and B of fig. 4). The estimated relationships follow the hypotheses we presented in figs. 1 and 2, although we found that the effect of $\text{Var}(R)$ at average $E(R)$ was linear with a slight sublinear saturation as $\text{Var}(R)$ increased. However, notice that the effect of $\text{Var}(R)$ on $E(H)$ depends strongly on $E(R)$ (panel C). The increase in $E(H)$ over $\text{Var}(R)$ is strongly nonlinear at low values of $E(R)$, but it is not affected by $\text{Var}(R)$ much when $E(R)$ is very low (and $E(H)$ is consequently very high).




Figure 4: Effects of $E(R)$ and $\text{Var}(R)$ on simulated spatial requirements with 95% Bayesian credible intervals for the mean. While the effect of $\text{Var}(R)$ is sublinear in panel B, the effect of $\text{Var}(R)$ depends strongly on $E(R)$ (see panel C). Credible intervals were calculated using 10,000 samples from the posterior distribution. The relationships were estimated using a Generalized Additive Model for Location and Scale with a Gamma location-scale family of distributions (`mgcv::gammals`). Details on the model structure are provided in Appendix B.

As expected by the changes in the spread of the points in panels A and B of figure 4, the variance in H also depends on $E(R)$ and $\text{Var}(R)$ (figure 5A-B). Here, the effects of the two variables are more pronounced, even for low values of $E(R)$ or high values of $\text{Var}(R)$. Consequences of these effects are explored in the discussion section.




Figure 5: Effects of $E(R)$ and $Var(R)$ on the variance in simulated spatial requirements, with 95% Bayesian credible intervals for the estimated variance. Credible intervals were calculated using 10,000 samples from the posterior distribution. Additional details on the model structure are provided in Appendix B.

253 A case study on a lowland tapir in the Brazilian Cerrado

254 The simulations in the section above mostly follow the hypotheses we presented in the
 255 introduction, but they are based on assumptions that are often not met in real natural en-
 256 viroments. Organisms live in spatiotemporally heterogeneous and dynamic environments,
 257 which the development of perceptual ranges, navigation, and memory. Together, these abil-
 258 ities result in selective space use that depends on resource availability (Kacelnik et al. 1992)
 259 and resource depletion (Charnov 1976).

260 In this section, we test the hypotheses illustrated in the introduction using empirical
 261 tracking data on a lowland tapir from the Brazilian Cerrado along with estimates of $E(R)$ and
 262 $Var(R)$ using the Normalized Difference Vegetation Index (NDVI, see Pettorelli et al. 2011),
 263 a remote-sensed measure of landscape greenness, as a proxy of forage abundance. Appendix
 264 C contains additional information on how NDVI was modeled along with details on the
 265 continuous-time movement models (Noonan et al. 2019a; Fleming and Calabrese 2021) and
 266 autocorrelated kernel density estimation (Noonan et al. 2019b; Alston et al. 2022; Silva et
 267 al. 2022) used to quantify the tapir's space-use requirements. Through this example, we aim
 268 to provide support for the two hypotheses using empirical data that do not depend on as
 269 strong assumptions as those required by the simulations in the section above. Concurrently,

we also hope to show the potential complications that arise with empirical data.

Fig. 6 illustrates how a tapir in the Brazilian Cerrado (data from the individual labelled as “Anna” in the dataset from Medici et al. 2022) adapts its spatial needs to changes in $E(R)$ and $\text{Var}(R)$. Panels A and B show the changes in seven-day average mean and variance in NDVI, respectively, experienced by the tapir during the tracking period. The mean and variance in NDVI were estimated using a Generalized Additive Model for Location and Scale (GAMLS, theory: Rigby and Stasinopoulos 2005; Stasinopoulos and Rigby 2007; examples: Bjoernsdahl et al. 2022; Mariën et al. 2022; Gushulak et al. 2024) with a Beta family of distributions (NDVI values ranged from 0.3534 to 0.9475). Panel C shows the changes in the tapir’s 7-day home range over time. Note how the tapir uses more space during periods of lower NDVI (e.g., August 2017) and less space during periods with high NDVI (January 2018). Additionally, when resources are scarce and highly unpredictable (August 2018), the tapir uses up to 5 times more space than when resources are abundant and predictable (e.g., January 2018). Finally, panels D and E show the estimated (marginal) effects of $\mu(t)$ and $\sigma^2(t)$ on the tapir’s space use. Since $\mu(t)$ and $\sigma^2(t)$ are strongly correlated (panel F) and (spatio)temporally autocorrelated (panels A and B), the effects of R on H should be modeled carefully. To avoid over-fitting the model, we constrained the smooth effects of $\mu(t)$ and $\sigma^2(t)$ using low number of knots ($k = 3$). Additional information is provided in appendix C. Overall, the results presented in fig. 6 support our hypotheses, as the tapir’s 7-day home range decreases nonlinearly with $\mu(t)$ and increases nonlinearly with $\sigma^2(t)$. However, due to the strong nonlinear correlation between the two predictors, the marginal effects do not follow the data closely, particularly in the case of $\mu(t)$. We provide comments about these results in the related section in the discussion below.

Discussion

This paper presents two hypotheses for describing the effects of resource abundance and stochasticity on organisms’ space-use requirements. We support the hypotheses using quan-




Figure 6: Seven-day home-range size of a lowland tapir (*Tapirus terrestris*) in response to changes in mean and variance in resources. (A) Trends in resource abundance over time, $\mu(t)$, estimated as the average mean NDVI at the locations visited by the tapir during a seven-day period. (B) Variance in resources over time, $\sigma^2(t)$, estimated as the average variance in NDVI at the locations visited by the tapir during a seven-day period. (C) Estimated seven-day home range based on the 95% utilization quantiles. (D, E) Estimated marginal effects of $\mu(t)$ and $\sigma^2(t)$ on home-range size. The model accounted for the marginal effects of $\mu(t)$ and $\sigma^2(t)$ and their interaction on mean space-use requirements and the variance around them. (F) The effect of $\mu(t)$ does not follow the data closely because $E(R)$ and $\text{Var}(R)$ are highly correlated. Consequently, while estimating the effects of $E(R)$ and $\text{Var}(R)$ via separate models would allow result in a closer fit, the estimated effects would be inappropriate because they do not disentangle the effects of $E(R)$ and $\text{Var}(R)$. See Appendix C for additional information. The tapir movement data corresponds to the individual named “Anna” from the Cerrado sample of Medici *et al.* (2022).

296 titative simulations and an empirical case study. Each of the two methods allows us to high-
297 light the strengths and limitations of simulation-based and data-driven approaches, which
298 we discuss below.

299 **Strengths and limitations of the simulation-based approach**

300 The simulations we present here provide a simple and direct estimate of the effects of re-
301 source abundance and stochasticity on organisms' space-use requirements. The combination
302 of simplicity (few drivers of movement) and realism (continuous-time movement models)
303 allowed us to estimate the effects of $E(R)$ and $\text{Var}(R)$ on organisms' space-use requirements
304 using simple causal models of inference without the confounding variables that are often
305 present in empirical data (e.g., predator avoidance, territoriality, competition, landscape
306 connectivity). Additionally, since our sample size was virtually unlimited, we were able to
307 simulate sufficient data to ensure the simulations were stable and the resulting estimated
308 coefficients were well estimated (see Appendix B).

309 Avoiding confounding variables rather than attempting to account for them (as is required
310 with empirical data) allowed us to develop theory easily while providing a baseline model
311 against which one can contrast empirical examples. In figures B6, we present the Directed
312 Acyclical Graph (DAG, see McElreath 2016) we used to infer causal the mechanisms of
313 changes in H and estimate the direct, partial effects of $E(R)$ and $\text{Var}(R)$ on H (contrast
314 this with figure C5 and the section below).

315 By quantifying the deviations from the simple theoretical models, one can then assess the
316 importance of other drivers movement and behavior. Still, the simplicity of the underlying
317 model and its (often invalid) assumptions often limit the applicability of simulated models in
318 realistic scenarios. Responses to $E(R)$ and $\text{Var}(R)$ in real ecosystems likely depend on many
319 other drivers of movement, including competition, movement costs, and predation, which
320 may affect the effects of $E(R)$ and $\text{Var}(R)$ on both $E(H)$ and $\text{Var}(H)$ (Jetz et al. 2004;
321 Noonan et al. 2023). And while it is possible to account for all of these factors through

322 simulations, empirical data often can provide evidence for these effects without the need for
323 complex model structures.

324 **Strengths and limitations of the empirical approach**

325 In the absence of direct measurements of R , empirical models estimating the effects of R on
326 H must depend on proxies of R , such as NDVI. However, using proxies of R rather than
327 direct measurements introduces sources of bias. While R and NDVI are correlated (e.g.,
328 Phillips et al. 2008; Merkle et al. 2016, 2019; Middleton et al. 2018; Geremia et al. 2019;
329 Seigle-Ferrand et al. 2021), the relationship between the two can be weak (Gautam et al.
330 2019) and nonlinear (**ref?**). This is because remote-sensed proxies of R and they do not
331 measure resource availability directly, so they introduce two sources of bias: ecosystem-level
332 biases (indicated as Z in the DAG in figure C5) and satellite-level confounding variables (S
333 in figure C5).

334 Examples of ecosystem-level biases are the effects of competition, predation, habitat
335 connectivity, and movement costs, all of which can depend on habitat quality, and, conse-
336 quently, be correlated nonlinearly to R and *NDVI* (Jetz et al. 2004; Prox and Farine 2020).

337 Resource-rich patches can attract larger amounts of competitors (Kacelnik et al. 1992) and
338 predators, which may increase pressures from competition (**ref?**) and predation (Brown et
339 al. 1999). While such pressures may increase avoidance behaviors, avoidance may have a
340 complex effect on H , since it can result in both an expansion of the home range (Jetz et al.
341 2004; Prox and Farine 2020) or a contraction of an animal's movement, since larger home
342 ranges can be harder to defend and also result in higher movement costs (Grant 1993; Jetz
343 et al. 2004). Additionally, persistent territorial defense from competitors may prevent or-
344 ganisms from using space freely and as necessary (theoretical framework: **despotic_ref?**;
345 examples: Rich et al. 2012; Bengsen et al. 2016; Tórrez-Herrera et al. 2020).

346 Satellite-level confounds include information loss due to coarse spatiotemporal resolution
347 (**ref?**), satellite-level error (Tian et al. 2015; Fan and Liu 2016; Huang et al. 2021), and

348 other limitations of remote sensing (e.g., inability to quantify specific resources, competitive
349 pressure, and predator avoidance). However, nonlinear models such as Generalized Additive
350 Models (Wood 2017) can help account for preferences for intermediate values of NDVI (e.g.,
351 young grass rather than mature grasslands, see **ref?**). In the case study we present here,
352 sampling limitations resulted in a strong nonlinear correlation between $E(R)$ and $\text{Var}(R)$ that
353 prevented us from estimating nonlinear preferences in mean NDVI (i.e., the proxy for $E(R)$),
354 since estimating the divers of movement and disentangling them from potential confounds
355 in a stochastic environemtn is often complex or impossible (e.g., Wang et al. 2019). A
356 tracking time series of two or more years or additional individuals would have provided
357 better estimates, but that was beyond the scope of this empirical example.

358 **here:** need a few lines to tie this back to the general

359 Conclusion

360 needs a few more lines at the end

361 The work presented here provides a unifying framework for viewing movement as a response
362 to resource abundance and stochasticity. We provide realistic and flexible hypotheses of
363 the effects of $E(R)$ and $\text{Var}(R)$ on organisms' space-use requirements and movement behav-
364 ior. We demonstrate that organisms' space-use requirements respond nonlinearly to both
365 resource abundance and stochasticity, and we demonstrate the importance of accounting
366 for $\text{Var}(R)$ besides $E(R)$. We also provide a moving-window approach that allows one to
367 model movement continuously rather than imposing arbitrary thresholds between different
368 periods or types of behaviors, including temporal changes in space-use requirements and
369 shifts between range-residency and nomadism. The methods presented here can be applied
370 to assess both current and future quality of protected areas while also estimating the amount
371 of space organisms currently need and will need in the future under different climate change
372 scenarios.

³⁷³ **Conflict of interest**

³⁷⁴ The authors declare there are no conflicts of interest.

375 **References**

- 376 Abrahms, B., E. L. Hazen, E. O. Aikens, M. S. Savoca, J. A. Goldbogen, S. J. Bograd,
377 M. G. Jacox, et al. 2019. Memory and resource tracking drive blue whale migrations.
378 Proceedings of the National Academy of Sciences 116:5582–5587.
- 379 Alston, J. M., C. H. Fleming, R. Kays, J. P. Streicher, C. T. Downs, T. Ramesh, B. Reineking,
380 et al. 2022. Mitigating pseudoreplication and bias in resource selection functions with
381 autocorrelation-informed weighting. Methods in Ecology and Evolution 2041–210X.14025.
- 382 Anderson, S. C., T. A. Branch, A. B. Cooper, and N. K. Dulvy. 2017. Black-swan events in
383 animal populations. Proceedings of the National Academy of Sciences 114:3252–3257.
- 384 Andersson, K. 2004. Elbow-joint morphology as a guide to forearm function and foraging
385 behaviour in mammalian carnivores. Zoological Journal of the Linnean Society 142:91–
386 104.
- 387 Baldwin, R., and A. Bywater. 1984. Nutritional energetics of animals. Annual review of
388 nutrition 4:101–114.
- 389 Bengsen, A. J., D. Algar, G. Ballard, T. Buckmaster, S. Comer, P. J. S. Fleming, J. A. Friend,
390 et al. 2016. Feral cat home-range size varies predictably with landscape productivity and
391 population density. Journal of Zoology 298:112–120.
- 392 Berger, J., C. Hartway, A. Gruzdev, and M. Johnson. 2018. Climate degradation and
393 extreme icing events constrain life in cold-adapted mammals. Scientific Reports 8:1156.
- 394 Bista, D., G. S. Baxter, N. J. Hudson, S. T. Lama, and P. J. Murray. 2022. Effect of
395 disturbances and habitat fragmentation on an arboreal habitat specialist mammal using
396 GPS telemetry: A case of the red panda. Landscape Ecology 37:795–809.
- 397 Bjorndahl, J. A., C. A. C. Gushulak, S. Mezzini, G. L. Simpson, H. A. Haig, P. R. Leavitt,
398 and K. Finlay. 2022. Abrupt changes in the physical and biological structure of endorheic
399 upland lakes due to 8-m lake-level variation during the 20th century. Limnology and
400 Oceanography 67:1022–1039.

- 401 Bradsworth, N., J. G. White, A. R. Rendall, N. Carter, D. A. Whisson, and R. Cooke.
402 2022. Using thresholds to determine priorities for apex predator conservation in an urban
403 landscape. *Landscape and Urban Planning* 228:104559.
- 404 Brown, J. S., J. W. Laundre, and M. Gurung. 1999. The ecology of fear: Optimal foraging,
405 game theory, and trophic interactions. *Journal of Mammalogy* 80:385–399.
- 406 Burson, A., M. Stomp, E. Greenwell, J. Grosse, and J. Huisman. 2018. Competition for nu-
407 trients and light: Testing advances in resource competition with a natural phytoplankton
408 community. *Ecology* 99:1108–1118.
- 409 Burt, W. H. 1943. Territoriality and home range concepts as applied to mammals. *Journal*
410 of *Mammalogy* 24:346.
- 411 Charnov, E. L. 1976. Optimal foraging, the marginal value theorem. *Theoretical Population*
412 *Biology* 9:129–136.
- 413 Chevin, L.-M., R. Lande, and G. M. Mace. 2010. Adaptation, plasticity, and extinction in a
414 changing environment: Towards a predictive theory. (J. G. Kingsolver, ed.) *PLoS Biology*
415 8:e1000357.
- 416 Dai Pra, R., S. M. Mohr, D. K. Merriman, S. N. Bagriantsev, and E. O. Gracheva. 2022.
417 Ground squirrels initiate sexual maturation during hibernation. *Current Biology* 32:1822–
418 1828.e4.
- 419 Douglas, D. J. T., and J. W. Pearce-Higgins. 2014. Relative importance of prey abundance
420 and habitat structure as drivers of shorebird breeding success and abundance: Drivers of
421 shorebird breeding success and abundance. *Animal Conservation* 17:535–543.
- 422 Fagan, W. F., M. A. Lewis, M. Auger-Méthé, T. Avgar, S. Benhamou, G. Breed, L. LaDage,
423 et al. 2013. Spatial memory and animal movement. (J. Clobert, ed.) *Ecology Letters*
424 16:1316–1329.
- 425 Falcón-Cortés, A., D. Boyer, E. Merrill, J. L. Frair, and J. M. Morales. 2021. Hierarchical,
426 memory-based movement models for translocated elk (*cervus canadensis*). *Frontiers in*
427 *Ecology and Evolution* 9:702925.

- ⁴²⁸ Fan, X., and Y. Liu. 2016. A global study of NDVI difference among moderate-resolution
⁴²⁹ satellite sensors. *ISPRS Journal of Photogrammetry and Remote Sensing* 121:177–191.
- ⁴³⁰ Fjelldal, M. A., J. Wright, and C. Stawski. 2021. Nightly torpor use in response to weather
⁴³¹ conditions and individual state in an insectivorous bat. *Oecologia* 197:129–142.
- ⁴³² Fleming, C. H., and J. M. Calabrese. 2021. Ctmm: Continuous-time movement modeling.
- ⁴³³ Foley, C., N. Pettorelli, and L. Foley. 2008. Severe drought and calf survival in elephants.
⁴³⁴ *Biology Letters* 4:541–544.
- ⁴³⁵ Gautam, H., E. Arulmalar, M. R. Kulkarni, and T. N. C. Vidya. 2019. NDVI is not
⁴³⁶ reliable as a surrogate of forage abundance for a large herbivore in tropical forest habitat.
⁴³⁷ *Biotropica* 51:443–456.
- ⁴³⁸ Geremia, C., J. A. Merkle, D. R. Eacker, R. L. Wallen, P. J. White, M. Hebblewhite, and M.
⁴³⁹ J. Kauffman. 2019. Migrating bison engineer the green wave. *Proceedings of the National
440 Academy of Sciences* 116:25707–25713.
- ⁴⁴¹ Ghislandi, P. G., S. Pekár, M. Matzke, S. Schulte-Döinghaus, T. Bilde, and C. Tuni. 2018.
⁴⁴² Resource availability, mating opportunity and sexual selection intensity influence the ex-
⁴⁴³ pression of male alternative reproductive tactics. *Journal of Evolutionary Biology* 31:1035–
⁴⁴⁴ 1046.
- ⁴⁴⁵ Grant, J. W. A. 1993. Whether or not to defend? The influence of resource distribution.
⁴⁴⁶ *Marine Behaviour and Physiology* 23:137–153.
- ⁴⁴⁷ Gurarie, E., C. H. Fleming, W. F. Fagan, K. L. Laidre, J. Hernández-Pliego, and O.
⁴⁴⁸ Ovaskainen. 2017. Correlated velocity models as a fundamental unit of animal move-
⁴⁴⁹ ment: Synthesis and applications. *Movement Ecology* 5:13.
- ⁴⁵⁰ Gushulak, C. A. C., S. Mezzini, K. E. Moir, G. L. Simpson, L. Bunting, B. Wissel, D. R.
⁴⁵¹ Engstrom, et al. 2024. Impacts of a century of land-use change on the eutrophication
⁴⁵² of large, shallow, prairie lake manitoba in relation to adjacent lake winnipeg (manitoba,
⁴⁵³ canada). *Freshwater Biology* 69:47–63.

- 454 Haney, S. D., and A. M. Siepielski. 2018. Tipping points in resource abundance drive
455 irreversible changes in community structure. *The American Naturalist* 191:668–675.
- 456 Harestad, A. S., and F. L. Bunnel. 1979. Home range and body weight—a reevaluation.
457 *Ecology* 60:389–402.
- 458 Harvey, P. H., and T. H. Clutton-Brock. 1981. Primate home-range size and metabolic
459 needs. *Behavioral Ecology and Sociobiology* 8:151–155.
- 460 Herfindal, I., J. D. C. Linnell, J. Odden, E. B. Nilsen, and R. Andersen. 2005. Prey density,
461 environmental productivity and home-range size in the eurasian lynx (*lynx lynx*). *Journal*
462 of *Zoology* 265:63–71.
- 463 Hirt, M. R., W. Jetz, B. C. Rall, and U. Brose. 2017. A general scaling law reveals why the
464 largest animals are not the fastest. *Nature Ecology & Evolution* 1:1116–1122.
- 465 Hou, R., C. A. Chapman, O. Jay, S. Guo, B. Li, and D. Raubenheimer. 2020. Cold and
466 hungry: Combined effects of low temperature and resource scarcity on an edge-of-range
467 temperate primate, the golden snub-nose monkey. *Ecography* 43:1672–1682.
- 468 Huang, S., L. Tang, J. P. Hupy, Y. Wang, and G. Shao. 2021. A commentary review on
469 the use of normalized difference vegetation index (NDVI) in the era of popular remote
470 sensing. *Journal of Forestry Research* 32:1–6.
- 471 Jetz, W., C. Carbone, J. Fulford, and J. H. Brown. 2004. The scaling of animal space use.
472 *Science* 306:266–268.
- 473 Kacelnik, A., J. R. Krebs, and C. Bernstein. 1992. The ideal free distribution and predator-
474 prey populations. *Trends in Ecology & Evolution* 7:50–55.
- 475 Le Bot, T., A. Lescroël, J. Fort, C. Péron, O. Gimenez, P. Provost, and D. Grémillet. 2019.
476 Fishery discards do not compensate natural prey shortage in northern gannets from the
477 english channel. *Biological Conservation* 236:375–384.
- 478 Levin, S. A. 1992. The problem of pattern and scale in ecology: The robert h. MacArthur
479 award lecture. *Ecology* 73:1943–1967.

- 480 Levins, R. A. 1974. Evolution in changing environments: Some theoretical explorations.
- 481 Monographs in population biology (3. printing.). Princeton Univ. Press, Princeton, NJ.
- 482 Lindstedt, S. L., and M. S. Boyce. 1985. Seasonality, fasting endurance, and body size in
- 483 mammals. *The American Naturalist* 125:873–878.
- 484 Logares, R., and M. Nuñez. 2012. Black swans in ecology and evolution: The importance
- 485 of improbable but highly influential events. *Ideas in Ecology and Evolution*.
- 486 Lucherini, M., and S. Lovari. 1996. Habitat richness affects home range size in the red fox
- 487 *vulpes vulpes*. *Behavioural Processes* 36:103–105.
- 488 Mariën, B., D. Papadimitriou, T. Kötiläinen, P. Zuccarini, I. Dox, M. Verlinden, T. Heinecke,
- 489 et al. 2022. Timing leaf senescence: A generalized additive models for location, scale and
- 490 shape approach. *Agricultural and Forest Meteorology* 315:108823.
- 491 McClintic, L. F., J. D. Taylor, J. C. Jones, R. D. Singleton, and G. Wang. 2014. Effects of
- 492 spatiotemporal resource heterogeneity on home range size of american beaver. *Journal of*
- 493 *Zoology* 293:134–141.
- 494 McElreath, R. 2016. Statistical rethinking: A bayesian course with examples in r and stan.
- 495 Chapman & hall/CRC texts in statistical science series. CRC Press/Taylor & Francis
- 496 Group, Boca Raton.
- 497 Medici, E. P., S. Mezzini, C. H. Fleming, J. M. Calabrese, and M. J. Noonan. 2022. Move-
- 498 ment ecology of vulnerable lowland tapirs between areas of varying human disturbance.
- 499 *Movement Ecology* 10:14.
- 500 Merkle, J. A., K. L. Monteith, E. O. Aikens, M. M. Hayes, K. R. Hersey, A. D. Middleton, B.
- 501 A. Oates, et al. 2016. Large herbivores surf waves of green-up during spring. *Proceedings*
- 502 *of the Royal Society B: Biological Sciences* 283:20160456.
- 503 Merkle, J. A., H. Sawyer, K. L. Monteith, S. P. H. Dwinnell, G. L. Fralick, and M. J.
- 504 Kauffman. 2019. Spatial memory shapes migration and its benefits: Evidence from a
- 505 large herbivore. (J. Gaillard, ed.)*Ecology Letters* 22:1797–1805.

- 506 Middleton, A. D., J. A. Merkle, D. E. McWhirter, J. G. Cook, R. C. Cook, P. J. White, and
507 M. J. Kauffman. 2018. Green-wave surfing increases fat gain in a migratory ungulate.
508 Oikos 127:1060–1068.
- 509 Morellet, N., C. Bonenfant, L. Börger, F. Ossi, F. Cagnacci, M. Heurich, P. Kjellander,
510 et al. 2013. Seasonality, weather and climate affect home range size in roe deer across
511 a wide latitudinal gradient within europe. (T. Coulson, ed.)Journal of Animal Ecology
512 82:1326–1339.
- 513 Mueller, T., R. B. O'Hara, S. J. Converse, R. P. Urbanek, and W. F. Fagan. 2013. Social
514 learning of migratory performance. Science 341:999–1002.
- 515 Nandintsetseg, D., C. Bracis, P. Leimgruber, P. Kaczensky, B. Buuveibaatar, B. Lkhagva-
516 suren, B. Chimeddorj, et al. 2019. Variability in nomadism: Environmental gradients
517 modulate the movement behaviors of dryland ungulates. Ecosphere 10.
- 518 Nilsen, E. B., I. Herfindal, and J. D. C. Linnell. 2005. Can intra-specific variation in
519 carnivore home-range size be explained using remote-sensing estimates of environmental
520 productivity? Écoscience 12:68–75.
- 521 Noonan, M. J., C. H. Fleming, T. S. Akre, J. Drescher-Lehman, E. Gurarie, A.-L. Harrison,
522 R. Kays, et al. 2019a. Scale-insensitive estimation of speed and distance traveled from
523 animal tracking data. Movement Ecology 7:35.
- 524 Noonan, M. J., R. Martinez-Garcia, C. H. Fleming, B. G. De Figueiredo, A. H. Ali, N.
525 Attias, J. L. Belant, et al. 2023. *The search behavior of terrestrial mammals* (preprint).
526 Ecology.
- 527 Noonan, M. J., M. A. Tucker, C. H. Fleming, T. S. Akre, S. C. Alberts, A. H. Ali, J.
528 Altmann, et al. 2019b. A comprehensive analysis of autocorrelation and bias in home
529 range estimation. Ecological Monographs 89:e01344.
- 530 Pettorelli, N., S. Ryan, T. Mueller, N. Bunnefeld, B. Jedrzejewska, M. Lima, and K. Kausrud.
531 2011. The normalized difference vegetation index (NDVI): Unforeseen successes in animal
532 ecology. Climate Research 46:15–27.

- 533 Phillips, L. B., A. J. Hansen, and C. H. Flather. 2008. Evaluating the species energy
534 relationship with the newest measures of ecosystem energy: NDVI versus MODIS primary
535 production. *Remote Sensing of Environment* 112:4381–4392.
- 536 Poessel, S. A., B. Woodbridge, B. W. Smith, R. K. Murphy, B. E. Bedrosian, D. A. Bell,
537 D. Bittner, et al. 2022. Interpreting long-distance movements of non-migratory golden
538 eagles: Prospecting and nomadism? *Ecosphere* 13.
- 539 Polansky, L., W. Kilian, and G. Wittemyer. 2015. Elucidating the significance of spatial
540 memory on movement decisions by african savannah elephants using state-space models.
541 *Proceedings of the Royal Society B: Biological Sciences* 282:20143042.
- 542 Pretorius, M. D., L. Leeuwner, G. J. Tate, A. Botha, M. D. Michael, K. Durgaprasad, and
543 K. Chetty. 2020. Movement patterns of lesser flamingos *phoeniconaias minor* : Nomadism
544 or partial migration? *Wildlife Biology* 2020:1–11.
- 545 Prox, L., and D. Farine. 2020. A framework for conceptualizing dimensions of social orga-
546 nization in mammals. *Ecology and Evolution* 10:791–807.
- 547 R Core Team. 2023. R: A language and environment for statistical computing. R Foundation
548 for Statistical Computing, Vienna, Austria.
- 549 Reich, P. B. 2001. Body size, geometry, longevity and metabolism: Do plant leaves behave
550 like animal bodies? *Trends in Ecology & Evolution* 16:674–680.
- 551 Relyea, R. A., R. K. Lawrence, and S. Demarais. 2000. Home range of desert mule deer:
552 Testing the body-size and habitat-productivity hypotheses. *The Journal of Wildlife Man-
553 agement* 64:146.
- 554 Rich, L. N., M. S. Mitchell, J. A. Gude, and C. A. Sime. 2012. Anthropogenic mortal-
555 ity, intraspecific competition, and prey availability influence territory sizes of wolves in
556 montana. *Journal of Mammalogy* 93:722–731.
- 557 Rigby, R. A., and D. M. Stasinopoulos. 2005. Generalized additive models for location, scale
558 and shape (with discussion). *Journal of the Royal Statistical Society: Series C (Applied
559 Statistics)* 54:507–554.

- 560 Riotte-Lambert, L., and J. Matthiopoulos. 2020. Environmental predictability as a cause
561 and consequence of animal movement. *Trends in Ecology & Evolution* 35:163–174.
- 562 Rizzuto, M., S. J. Leroux, E. Vander Wal, I. C. Richmond, T. R. Heckford, J. Balluffi-Fry,
563 and Y. F. Wiersma. 2021. Forage stoichiometry predicts the home range size of a small
564 terrestrial herbivore. *Oecologia* 197:327–338.
- 565 Rocha, J. L., R. Godinho, J. C. Brito, and R. Nielsen. 2021. Life in deserts: The genetic
566 basis of mammalian desert adaptation. *Trends in Ecology & Evolution* 36:637–650.
- 567 Samarra, F. I. P., S. B. Tavares, J. Béesau, V. B. Deecke, A. Fennell, P. J. O. Miller, H.
568 Pétursson, et al. 2017. Movements and site fidelity of killer whales (*orcinus orca*) relative
569 to seasonal and long-term shifts in herring (*clupea harengus*) distribution. *Marine Biology*
570 164:159.
- 571 Samuels, J. X., J. A. Meachen, and S. A. Sakai. 2013. Postcranial morphology and the
572 locomotor habits of living and extinct carnivorans. *Journal of Morphology* 274:121–146.
- 573 Schmidt, N. M., C. Grøndahl, A. L. Evans, J.-P. Desforges, J. Blake, L. H. Hansen, L. T.
574 Beumer, et al. 2020. On the interplay between hypothermia and reproduction in a high
575 arctic ungulate. *Scientific Reports* 10:1514.
- 576 Seigle-Ferrand, J., K. Atmeh, J.-M. Gaillard, V. Ronget, N. Morellet, M. Garel, A. Loison,
577 et al. 2021. A systematic review of within-population variation in the size of home range
578 across ungulates: What do we know after 50 years of telemetry studies? *Frontiers in
579 Ecology and Evolution* 8:555429.
- 580 Silva, I., C. H. Fleming, M. J. Noonan, J. Alston, C. Folta, W. F. Fagan, and J. M. Calabrese.
581 2022. Autocorrelation-informed home range estimation: A review and practical guide.
582 *Methods in Ecology and Evolution* 13:534–544.
- 583 Simcharoen, A., T. Savini, G. A. Gale, S. Simcharoen, S. Duangchantrasiri, S. Pakpien, and
584 J. L. D. Smith. 2014. Female tiger *panthera tigris* home range size and prey abundance:
585 Important metrics for management. *Oryx* 48:370–377.

- 586 Singh, N. J., L. Börger, H. Dettki, N. Bunnefeld, and G. Ericsson. 2012. From migration
587 to nomadism: Movement variability in a northern ungulate across its latitudinal range.
588 Ecological Applications 22:2007–2020.
- 589 Southwood, T. R. E. 1977. Habitat, the templet for ecological strategies? The Journal of
590 Animal Ecology 46:336.
- 591 Stasinopoulos, M. D., and R. A. Rigby. 2007. Generalized additive models for location scale
592 and shape (GAMLSS) in r. Journal of Statistical Software 23.
- 593 Stefanescu, C., A. Ubach, and C. Wiklund. 2021. Timing of mating, reproductive status
594 and resource availability in relation to migration in the painted lady butterfly. Animal
595 Behaviour 172:145–153.
- 596 Steixner-Kumar, S., and J. Gläscher. 2020. Strategies for navigating a dynamic world.
597 Science 369:1056–1057.
- 598 Stephens, D. W., and E. L. Charnov. 1982. Optimal foraging: Some simple stochastic
599 models. Behavioral Ecology and Sociobiology 10:251–263.
- 600 Taylor, C. R., N. C. Heglund, and G. M. Maloiy. 1982. Energetics and mechanics of
601 terrestrial locomotion. I. Metabolic energy consumption as a function of speed and body
602 size in birds and mammals. Journal of Experimental Biology 97:1–21.
- 603 Teitelbaum, C. S., W. F. Fagan, C. H. Fleming, G. Dressler, J. M. Calabrese, P. Leimgruber,
604 and T. Mueller. 2015. How far to go? Determinants of migration distance in land
605 mammals. (M. Festa-Bianchet, ed.)Ecology Letters 18:545–552.
- 606 Teitelbaum, C. S., and T. Mueller. 2019. Beyond migration: Causes and consequences of
607 nomadic animal movements. Trends in Ecology & Evolution 34:569–581.
- 608 Tian, F., R. Fensholt, J. Verbesselt, K. Grogan, S. Horion, and Y. Wang. 2015. Evaluat-
609 ing temporal consistency of long-term global NDVI datasets for trend analysis. Remote
610 Sensing of Environment 163:326–340.
- 611 Tórrez-Herrera, L. L., G. H. Davis, and M. C. Crofoot. 2020. Do monkeys avoid areas
612 of home range overlap because they are dangerous? A test of the risk hypothesis in

- 613 white-faced capuchin monkeys (*cebus capucinus*). International Journal of Primatology
614 41:246–264.
- 615 Wang, X., L. Cao, A. D. Fox, R. Fuller, L. Griffin, C. Mitchell, Y. Zhao, et al. 2019.
616 Stochastic simulations reveal few green wave surfing populations among spring migrating
617 herbivorous waterfowl. Nature Communications 10:2187.
- 618 Watson, J. 2020. Ferruginous hawk (*buteo regalis*) home range and resource use on northern
619 grasslands in canada.
- 620 Wessling, E. G., T. Deschner, R. Mundry, J. D. Pruetz, R. M. Wittig, and H. S. Kühl. 2018.
621 Seasonal variation in physiology challenges the notion of chimpanzees (*pan troglodytes*
622 *verus*) as a forest-adapted species. Frontiers in Ecology and Evolution 6:60.
- 623 Wheat, R. E., S. B. Lewis, Y. Wang, T. Levi, and C. C. Wilmers. 2017. To migrate, stay
624 put, or wander? Varied movement strategies in bald eagles (*haliaeetus leucocephalus*).
625 Movement Ecology 5:9.
- 626 Wood, S. N. 2017. Generalized additive models: An introduction with r. Chapman &
627 hall/CRC texts in statistical science (Second edition.). CRC Press/Taylor & Francis
628 Group, Boca Raton.