

A new global measure of environmental unpredictability

Stefano Mezzini^{1,2}

Michael J. Noonan^{1,2,3,*}

¹ Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

² Department of Biology, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

³ Department of Computer Science, Math, Physics, and Statistics, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

* Correspondence: Michael J. Noonan <michael.noonan@ubc.ca>

Article type: Research article

Words in abstract: 0

Words in main text: 155

Figures: XXX

Tables: XXX

References: XXX

Appendices: XXX

Key words: remote sensing, NDVI, environmental stochasticity, XXX

₁ Abstract

2 Introduction

3 <https://www.sciencedirect.com/science/article/abs/pii/S0034425718305625> [https:](https://silvis.forest.wisc.edu/data/dhis/)
4 [//silvis.forest.wisc.edu/data/dhis/](https://silvis.forest.wisc.edu/data/dhis/)

5 Methods

6 maps of input data
7 NDVI palette is colorblind-friendly

8 Results

9 output of `draw()`
10 hex plots of mean and var vs common metrics (e.g., spp diversity, npp, hfi, ...)

11 Discussion

12 Useful references

- 13 • `{khroma}`: doi:10.5281/zenodo.1472077
- 14 • Keith *et al.* (2008)
- 15 • Chevin, Lande & Mace (2010)
- 16 • Rickbeil *et al.* (2019)
- 17 • Mueller *et al.* (2008)
- 18 • Pettoirelli *et al.* (2005)
- 19 • Keys, Barnes & Carter (2021)
- 20 • Nilsen, Herfindal & Linnell (2005)
- 21 • Stephens & Charnov (1982)
- 22 • Merkle *et al.* (2016)

- Tian *et al.* (2015)
- Huang *et al.* (2021)
- Fan & Liu (2016)
- Wang *et al.* (2019)
- Pease (2024)
- Xu *et al.* (2021)
- Mezzini *et al.* (2025)

References

- Chevin L.-M., Lande R. & Mace G.M. (2010). Adaptation, plasticity, and extinction in a changing environment: Towards a predictive theory. *PLoS Biology* **8**, e1000357. <https://doi.org/10.1371/journal.pbio.1000357>
- Fan X. & Liu Y. (2016). A global study of NDVI difference among moderate-resolution satellite sensors. *ISPRS Journal of Photogrammetry and Remote Sensing* **121**, 177–191. <https://doi.org/10.1016/j.isprsjprs.2016.09.008>
- Huang S., Tang L., Hupy J.P., Wang Y. & Shao G. (2021). A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *Journal of Forestry Research* **32**, 1–6. <https://doi.org/10.1007/s11676-020-01155-1>
- Keith D.A., Akçakaya H.R., Thuiller W., Midgley G.F., Pearson R.G., Phillips S.J., *et al.* (2008). Predicting extinction risks under climate change: Coupling stochastic population models with dynamic bioclimatic habitat models. *Biology Letters* **4**, 560–563. <https://doi.org/10.1098/rsbl.2008.0049>
- Keys P.W., Barnes E.A. & Carter N.H. (2021). A machine-learning approach to human footprint index estimation with applications to sustainable development. *Environmental Research Letters* **16**, 044061. <https://doi.org/10.1088/1748-9326/abe00a>
- Merkle J.A., Monteith K.L., Aikens E.O., Hayes M.M., Hersey K.R., Middleton A.D., *et al.* (2016). Large herbivores surf waves of green-up during spring. *Proceedings of the Royal*

Society B: Biological Sciences **283**, 20160456. <https://doi.org/10.1098/rspb.2016.0456>

Mezzini S., Fleming C.H., Medici E.P. & Noonan M.J. (2025). How resource abundance and resource stochasticity affect organisms' range sizes. *Movement Ecology* **13**, 20. <https://doi.org/10.1186/s40462-025-00546-5>

Mueller T., Olson K.A., Fuller T.K., Schaller G.B., Murray M.G. & Leimgruber P. (2008). In search of forage: Predicting dynamic habitats of mongolian gazelles using satellite-based estimates of vegetation productivity. *Journal of Applied Ecology* **45**, 649–658. <https://doi.org/10.1111/j.1365-2664.2007.01371.x>

Nilsen E.B., Herfindal I. & Linnell J.D.C. (2005). Can intra-specific variation in carnivore home-range size be explained using remote-sensing estimates of environmental productivity? *Écoscience* **12**, 68–75. <https://doi.org/10.2980/i1195-6860-12-1-68.1>

Pease B.S. (2024). Ecological scales of effect vary across space and time. *Ecography* **2024**, e07163. <https://doi.org/10.1111/ecog.07163>

Pettorelli N., Vik J.O., Mysterud A., Gaillard J.-M., Tucker C.J. & Stenseth N.Chr. (2005). Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution* **20**, 503–510. <https://doi.org/10.1016/j.tree.2005.05.011>

Rickbeil G.J.M., Merkle J.A., Anderson G., Atwood M.P., Beckmann J.P., Cole E.K., *et al.* (2019). Plasticity in elk migration timing is a response to changing environmental conditions. *Global Change Biology* **25**, 2368–2381. <https://doi.org/10.1111/gcb.14629>

Stephens D.W. & Charnov E.L. (1982). Optimal foraging: Some simple stochastic models. *Behavioral Ecology and Sociobiology* **10**, 251–263. <https://doi.org/10.1007/BF00302814>

Tian F., Fensholt R., Verbesselt J., Grogan K., Horion S. & Wang Y. (2015). Evaluating temporal consistency of long-term global NDVI datasets for trend analysis. *Remote Sensing of Environment* **163**, 326–340. <https://doi.org/10.1016/j.rse.2015.03.031>

Wang X., Cao L., Fox A.D., Fuller R., Griffin L., Mitchell C., *et al.* (2019). Stochastic simulations reveal few green wave surfing populations among spring migrating herbivorous waterfowl. *Nature Communications* **10**, 2187. <https://doi.org/10.1038/s41467-019-09971-8>

76 Xu W., Barker K., Shawler A., Van Scoyoc A., Smith J.A., Mueller T., *et al.* (2021).
77 The plasticity of ungulate migration in a changing world. *Ecology* **102**, e03293. [https:](https://doi.org/10.1002/ecy.3293)
78 [//doi.org/10.1002/ecy.3293](https://doi.org/10.1002/ecy.3293)