

DENVar: A global Dynamic Estimate of NDVI Variance

Stefano Mezzini^{1,2} Gavin L. Simpson³ Michael J. Noonan^{1,2,4,*}

¹ Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

² Department of Biology, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

³ Department of Animal & Veterinary Sciences, Aarhus University, Denmark.

⁴ Department of Computer Science, Math, Physics, and Statistics, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

* Correspondence: [Michael J. Noonan <michael.noonan@ubc.ca>](mailto:michael.noonan@ubc.ca)

Article type: Research article

Words in abstract: 259

Words in main text: 2922

Figures: XXX

Tables: XXX

References: XXX

Appendices: XXX

Key words: remote sensing, NDVI, environmental stochasticity, XXX

Abstract

Remote-sensing estimates of habitat productivity such as the Normalized Difference Vegetation Index (NDVI) have helped develop and support a variety of hypotheses and inform policy and industry-related decisions. The free availability of such data has allowed widespread access and use of well-established metrics of habitat productivity. However, historically, users tended to focus solely on (predictable) trends in productivity. In recent years, people have been recognizing the importance of environmental unpredictability (stochasticity), particularly as climate change and human-induced rapid environmental change transform the ecosystems species have evolved in and adapted to. We address this gap by presenting a new global Dynamic Estimate of NDVI Variance, *DENVar*. We estimate spatiotemporal trends in mean NDVI and the variance around the mean using Hierarchical Generalized Additive Models, which provide a flexible yet transparent model structure, unlike machine learning approaches such as neural networks. We show that *DENVar* can be used as a reliable proxy of environmental stochasticity, strongly correlates with **XXX**, and can be used to test many hypotheses related to environmental stochasticity, such as forage stochasticity, regime shifts, We conclude by offering some considerations around quantifying and interpreting environmental stochasticity, and we provide a publicly and freely available shiny app for estimating mean NDVI and the variance around it, *DENVar*, for given coordinates and optional dates. The app requires no knowledge of statistics or coding. Additionally, we provide a temporally static raster of mean NDVI and *DENVar* for the years 1981-2025 that can be used for GIS applications and does not require any knowledge of R.

22 useful refs:

23 • <https://www.sciencedirect.com/science/article/abs/pii/S0034425711001957?fr=RR->
24 [2&ref=pdf_download&rr=962beb22bcb58443](https://www.sciencedirect.com/science/article/abs/pii/S0034425711001957?fr=RR-2&ref=pdf_download&rr=962beb22bcb58443)

25 • <https://www.nature.com/articles/s41467-025-61585-5>

26 • see to-read folder

27 submitting at journal of remote sensing? (<https://spj.science.org/page/remotesensing/>
28 [for-authors](https://spj.science.org/page/remotesensing/))

29 Introduction

30 The Normalized Difference Vegetation Index (NDVI), an estimate of plant greenness and
31 productivity, was first introduced by Rouse et al. (1974a) and Rouse et al. (1974b) using
32 spectral data from the Landsat 1 satellite. NDVI allowed researchers to evaluate plant pro-
33 ductivity and phenology using satellite data (Tucker, 1979) or hand-held devices (Tucker
34 et al., 1979) via a proxy of primary productivity, forage availability, and vegetative cover.
35 Satellite measurements of NDVI thus allowed analysts to remotely monitor not only veg-
36 etation but also animals' responses to changes in forage availability (Rouse et al., 1974b;
37 Pettorelli et al., 2005; Pettorelli et al., 2011).

38 Since NDVI is not a direct measurement primary productivity but a proxy, it should be
39 interpreted carefully (Pettorelli et al., 2011). NDVI is prone to loss of sensitivity in very
40 green (i.e., highly saturated) environments (Matsushita et al., 2007; Son et al., 2014), and
41 estimates vary across sensors (Beck et al., 2011; Huang et al., 2021). Results can depend on
42 the sensor used to measure NDVI (Fan & Liu, 2016), and the coarse scale of satellite-derived
43 NDVI can be a substantial barrier when when assessing fine-scale changes, heterogeneity,
44 or responses (Reid et al., 2018; Nathan et al., 2022; Goodchild & Quattrochi, 2023). Still,
45 NDVI is often used because it is free, well-established, and available at a wide variety of
46 spatiotemporal scales (Tarnavsky, Garrigues & Brown, 2008) while also requiring minimal
47 work to acquire.

48 NDVI has proven particularly useful in a multitude of fields, including animal movement
49 behavior (Merkle et al., 2016; Geremia et al., 2019), landscape fragmentation (Mallegowda
50 et al., 2015), phenology (Hmimina et al., 2013), habitat classification (Clerici, Weissteiner
51 & Gerard, 2012), and forest recovery after fires (João et al., 2018). Historically, analysts
52 focused on changes in mean NDVI, but recent work has included other metrics of NDVI,
53 including rates of change (Aikens et al., 2017) and changes in variance (i.e., variation around
54 the mean, stochasticity; Zhang et al., 2009; Mutti et al., 2020; Mezzini et al., 2025). However,

estimating trends in NDVI variance presents some additional challenges, particularly when environments are dynamic and (spatio)temporally heterogeneous.

Fig. 1) illustrates some of the difficulties that arise in modeling trends in variance in NDVI. Disentangling signal from noise is a central aspect of estimating trends in NDVI variance, but what constitutes as signal is dependent on the observer’s scale of perception (Levin, 1992; Steixner-Kumar & Gläscher, 2020). A forest fire may be extremely disruptive to a small animal with limited motility and a small range size, while a bird or a large mammal may not struggle to survive after relocating or altering its movement behavior (Nathan et al., 2008; Riotte-Lambert & Matthiopoulos, 2020). Similarly, what may be a severe drought to an annual plant may be no more than an unusually dry year for a plant with a longer generation timescale (Frankham & Brook, 2004). Thus, disentangling trends in mean NDVI from changes in NDVI variance requires a conscious decision about the scale at which to consider the data.

In this article, we present the Dynamic Estimate of NDVI Variance (DENVar). We estimate DENVar using 44 years of NDVI data from the AVHRR and VIIRS collection of sensors and Generalized Additive Models (GAMs). Although satellites with AVHRR and VIIRS sensors provide data at a coarser resolution (0.05°) than other satellites (e.g., 250m for MODIS), they are the only sensors to provide data at a daily frequency and for more than 40 years. Generalized Additive Models provide a flexible yet interpretable approach for estimating trends in mean and variance in NDVI without the rigidity of other more traditional methods (e.g., ARIMA) or the opaqueness of machine learning models such as neural networks (Wood, 2017; Simpson, 2018). We illustrate the relationships between DENVar and relevant ecological metrics, and we provide analysts with three options for incorporating DENVar in their own analyses without requiring any knowledge of R: (1) rasters of long-term averages from 1981 to 2025, rasters of yearly averages from 1982 to 2024, and a simple Shiny app that provides estimates for specific locations and dates.




Figure 1: Estimates of variance in NDVI depend on the estimated trends in mean NDVI. Black lines indicate the estimated mean and variance in simulated NDVI (points). Grey ribbons correspond to 95% credible intervals, while the colored segments highlight significant increases (red) and decreases (blue).

Smoother estimates of the mean (top left) result in wigglier and larger variance estimates (bottom left), since short-term trends in the data are considered to be stochastic events. In contrast, wigglier estimates of the mean (top right) result in smoother and smaller variance estimates (bottom right), since short-term trends in the data are attributed to changes in the mean. The data in the top row is the same in both columns. It was generated as the sum of a sinusoidal function and a mean-reverting, correlated, stochastic process with mean zero. ***ADD METHODS OR REFERENCE APPENDIX***

Materials and Methods

Experimental and Technical Design

- include a diagram to show the entire experimental design and illustrate the most significant elements: materials, treatments, measurements, data collection, methods of data analysis
- This will facilitate the editors and reviewers to understand and follow the whole concept, design, and results.

Choice of color palettes

We represent NDVI using a modified version of Fabio Crameri’s divergent *bukavu* palette (Crameri, 2018a; Crameri, Shephard & Heron, 2020), which has high-contrast for deuteranope and protanope vision. For NDVI values between 0 and 1 (the right half of the palette), the colors also have sufficient contrast for the colors to also be distinguishable by strong tritanope and achromatic vision. Fig. A1 contains an approximate representation of how each vision type would perceive each color palettes used. We obtained all color palettes with three or more colors from the `khroma` package (v. 1.14.0, Frerebeau, 2024) for R (v. 4.4.1, R Core Team, 2024).

Input data

We the NDVI image composites from the AVHRR and VIIRS sensors (Vermote & NOAA CDR Program, 2018, 2022) directly from the NOAA data server (www.ncei.noaa.gov/data/; Fig. A2)). The code for downloading the data is available on GitHub (see the code availability statement). After removing all probably or confidently cloudy pixels, the number of non-NA land raster cells was substantially larger than the maximum data frame size in R, ($3.17 \times 10^{10} > 2^{31} - 1 \approx 0.21 \times 10^{10}$), so we reduced the dataset size by splitting the dataset across the aggregating the averaged rasters by a factor of 2×2 with `terra::aggregate(fact`

= 2). Although this resulted in a reduction in sample size, the temporal averaging also reduced signal-related noise.

The predictor data for the models included: WWF biome (Olson & Dinerstein, 2002, see Fig. A3), ecoregion (the individual polygons from Olson & Dinerstein, 2002, see Fig. A4), integer day of year (1 to 366), integer year (1981 to 2025), and elevation above sea level (Fig. A5). We excluded all polygons that did not have any NDVI data (0.06% of land, 29% of the polygons, maximum area: $452 \text{ km}^2 \approx (21 \text{ km})^2$). We distinguished between biomes in the northern and southern hemisphere to allow for different seasonal trends between hemispheres. The 15-day averages allowed us to have non-repeating values of day of year across years, since 15 is not a factor of 365 or 366. We downloaded the global digital elevation model using the `get_elev_raster()` function from the `elevatr` package for R (v. 0.99.0, Hollister et al., 2023) with a resolution of 0.076 degrees.

Modeling

We estimated spatiotemporal trends in mean NDVI using Hierarchical Generalized Additive Models (HGAMs) via the `mgcv` package for R (v. 1.9-3, Wood, 2017). To reduce modeling fitting times with negligible losses to model accuracy, we used the `bam()` function with fast REstricted Marginal Likelihood (`method = fREML`) and covariate discretization (`discrete = TRUE`). See Wood, Goude & Shaw (2015) and Wood et al. (2017) for more information. Ideally, NDVI should be modeled using beta location-scale models (after the linear transformation $Y^* = \frac{Y+1}{2}$) to account for the fact that: (1) NDVI is bounded between -1 and 1, and (2) the variance in NDVI is dependent on the mean (and vice-versa), since ecosystems with very low NDVI (e.g., rock, ice, or concrete) or very high NDVI (e.g., dense forest) tend to have lower variance in NDVI. However, preliminary tests showed that fitting times for beta models were prohibitive, especially for beta location-scale models. In contrast, Gaussian models fit substantially faster and provided very similar spatialtemporal estimates of mean NDVI (see the `analysis/000-sardinia-test.R` script in the GitHub repository).

131 The global model for the mean NDVI had the structure below:

```
m_mean <- bam(
  ndvi_15_day_mean ~
    biome + # to avoid intercept shrinkage
    s(poly_id, bs = 'mrf', xt = list(nb = nbs)) +
    s(doy, biome, bs = 'fs', xt = list(bs = 'cc'), k = 10) +
    s(year, biome, bs = 'fs', xt = list(bs = 'cr'), k = 10) +
    ti(doy, year, biome, bs = c('cc', 'cr', 're'), k = c(5, 5)) +
    s(elevation_m, bs = 'cr', k = 5),
  family = gaussian(),
  data = d,
  method = 'fREML',
  knots = list(doy = c(0.5, 366.5)),
  drop.unused.levels = TRUE,
  discrete = TRUE,
  samfrac = 0.001, # find initial guesses with a subset of the data
  nthreads = future::availableCores(logical = FALSE) - 2,
  control = gam.control(trace = TRUE))
```

132 Biome-specific fixed-effect intercepts allowed to account for the differences in mean NDVI
133 through biomes without coefficient shrinkage (i.e., reversion towards the global mean). A
134 Markov Random Field of the ecoregions (`bs = mrf`, see page 240? of Wood, 2017) estimated
135 smoothed, ecoregion-specific deviations from the biome-level spatial means. Two factor-
136 smooth interaction terms (`bs = 'fs'`) accounted for biome-specific seasonal and yearly
137 trends. Seasonal terms were made cyclical by using cyclical cubic splines (`xt = list(bs`
138 `= 'cc')`), which are continuous up to and including second derivative at the edge knots
139 (0.5 and 366.5, which correspond to 00:00 of January 1st and 24:00 of December 31st, respec-
140 tively). A tensor product interaction smooth for each biome estimated the change in seasonal
141 trends over the years. Finally, a smooth of elevation above sea level accounted for the effect
142 of altitude. See Pedersen et al. (2019) for more information on hierarchical modeling and
143 the use and interpretation of factor smooth interaction terms.

144 A second HGAM estimated the variance in NDVI around the mean estimated by the
145 model above. The model had an identical structure to the model for the mean, with the
146 exception that (1) the response variable was the squared residual from the first model, such

that the model estimated the mean squared residual (i.e., the variance) for a given point in time and space, and (2) a smooth term of the estimated mean NDVI. The model is available below:

```
## *CHANGE TO SOS BASIS*
m_var <- bam(
  e_2 ~
    biome +
    s(poly_id, bs = 'mrf', xt = list(nb = nbs)) +
    s(doy, biome, bs = 'fs', xt = list(bs = 'cc'), k = 10) +
    s(year, biome, bs = 'fs', xt = list(bs = 'cr'), k = 10) +
    ti(doy, year, biome, bs = c('cc', 'cr', 're'), k = c(5, 5)) +
    s(elevation_m, bs = 'cr', k = 5) +
    s(mu_hat, bs = 'cr', k = 5),
  family = gaussian(),
  data = d,
  method = 'fREML',
  knots = list(doy = c(0.5, 366.5)),
  drop.unused.levels = TRUE,
  discrete = TRUE,
  samfrac = 0.001,
  nthreads = future::availableCores(logical = FALSE) - 2,
  control = gam.control(trace = TRUE))
```

Results

figures:

- plot grid of mean over space, `s(year)`, `s(doy)`, `s(elev)`
- plot grid of DENVar over space, `s(year)`, `s(doy)`, `s(elev)`, `s(mean)`
- histograms of DENVar around flat map of biomes (change colors)
- hex plots of DENVar with other variables (add temperature, annual precip, extreme weather events, fires, invasive species, max body size, disease occurrence, ...)

General results

- two Orthographic projections of DENVar: Northern and southern hemispheres on spring solstice

Correlations with other metrics

- temperature




Figure 2: Hex plots of the relationship between the global Dynamic Estimate of NDVI Variance (DENVar) and common ecological metrics. The fill of the hexagonal cells represents the log-10-transformed number of points within the cell. Rowwise, the variables are: mean NDVI (as estimated in this manuscript); the machine-learning Human Footprint Index (ml-HFI) of Keys, Barnes & Carter (2021); the 2015 cumulative, minimum, and seasonal range in the Dynamic Habitat Index of XXX, (F).

- seasonal temperature range
- precip
- seasonal temperature range
- spp diversity
- gross primary productivity
- max animal weight
- <https://silvis.forest.wisc.edu/globalwui/>
- <https://silvis.forest.wisc.edu/data/dhis/>: NDVI16 (Normalized Difference Vegetation Index), EVI16 (Enhanced Vegetation Index), FPAR8 (Fraction absorbed Photosynthetically Active Radiation), LAI8 (Leaf Area Index), GPP8 (Gross Primary Productivity)

How to obtain the data

Static rasters of mean and variance (1981-2025)

- for GIS users (tif)

Shiny app for obtaining values

- get values for specific coordinates: 1981-2025 average
- get values for specific coordinates and dates (upload as a CSV, download as a CSV)
- screenshot of the shiny app

Discussion

We have presented a new spatiotemporally dynamic metric of environmental stochasticity, which we have called the Dynamic Estimate of NDVI Variance, DENVar. In this section, we discuss the interpretation of DENVar in greater depth, and we argue the advantages and limitations of the statistical methods used. We conclude by suggesting some use cases of DENVar within animal movement ecology, landscape ecology, and phenology, but we invite readers to consider how DENVar may be of use in their field and adjacent disciplines.

Why use complicated statistics instead of machine learning?

Admittedly, the title of this section is misleading for two reasons. Firstly, some may argue that Generalized Additive Models are a form of machine-learning (here abbreviated as ML), but we intend to contrast GAMs with more “black-box” types of machine learning, such as neural networks and random forests. The second reason for which the title is misleading is that it implies that machine learning is not complicated, or at least that it is less complicated than the models we used. However, we argue that while traditional statistical models may seem more intimidating to some, their complexity is what allows analysts to produce interpretable results and measures of uncertainty (Wood, 2017). And while the theoretical basis for GAMs can be discouraging for those who do not have a strong background in mathematics or statistics, outputs from GAMs can be presented in an accessible manner even to those without a strong quantitative background through the use of well-designed figures, especially in a Bayesian context (when the aim is to estimate a parameter). In contrast, while ML

methods will often give understandable results, it may not be possible to discern *why* such results were “chosen” by the model. Instead, GAMs produce flexible yet interpretable estimated effects that can be represented with relatively simple figures, even if the mathematical formula(e) for the model may be complicated.

Ideally, a model should estimate or explore the relationships among variables such that results can be interpreted in a transparent and unambiguous manner, such that readers can scrutinize the underlying assumptions, methods, and decisions. We recognize that the assessing GAMs requires a certain level of expertise, but we hope the readers can appreciate the option to assess the GAMs more easily than a series of consecutive, weighted regressions (as in the case of neural networks). In contrast, GAMs learn from data with no initial starting assumptions other than user-imposed limitations of wiggleness, which are clearly outlined in each basis size of the models’ terms. The structure and wuggleness of GAMs can thus be grossly designed based on the system’s complexity and structure while allowing the data to inform the model trends, wiggleness, and coefficient size.

Interpreting DENVar: the importance of scale

- smoothness of mean impacts smoothness and size of variance: depends on scale of interest (e.g., animals’ ability to respond to, learn, and predict changes in the mean). Changes not attributed to trends in the mean are stochastic
- DENVar is a good metric as long as the spatiotemporal scale of interest is similar or larger to that of DENVar. May not work well for small-scale processes such as the movement dynamics of a small rodent, the phenology of a small orchard when spatiotemporal heterogeneity is finer than DENVar’s resolution

Environmental stochasticity and animal movement ecology

The field of animal movement ecology has long shown that animals alter their movement in response to resource abundance or availability (Burt, 1943; Charnov, 1976; Southwood,

1977; Broekman et al., 2024). More recently, movement ecologists have started focusing more on resource stochasticity (e.g., Stephens & Charnov, 1982; Rizzuto et al., 2021), but it is worth distinguishing between resource stochasticity and seasonality (Nilsen, Herfindal & Linnell, 2005). Seasonality is generally best interpreted as a cyclical change in the mean that many animals can learn to predict, as in the case of green-wave surfing (Middleton et al., 2018; Geremia et al., 2019). In contrast, environmental stochasticity refers to unpredictable changes, such as forest fires (ref?), sudden snow storms (Aikens et al., 2025), and unusual phenology (ref?). Animals also respond to stochastic changes (Mezzini et al., 2025), but their ability to adapt depends on behavioral plasticity (Rickbeil et al., 2019; Steixner-Kumar & Gläser, 2020), and memory (Polansky, Kilian & Wittemyer, 2015; Abrahms et al., 2019; Riotte-Lambert & Matthiopoulos, 2020; Falcón-Cortés et al., 2021; Ranc, Cagnacci & Moorcroft, 2022).

Botero et al. (2015) tipping points

Polazzo et al. (2024)

extreme events, black swans (Logares & Nuñez, 2012; Anderson et al., 2017)

- migration routes are key habitat (Ortega et al., in prep), especially in stochastic environments
- can fail to adapt to changes in conditions Sawyer et al. (2019)
- <https://www.sciencedirect.com/science/article/pii/S0960982220308484>
- <https://onlinelibrary.wiley.com/doi/full/10.1111/gcb.15169>
- industrial development affects animals' ability to respond to both predictable and unpredictable change (<https://www.nature.com/articles/s41559-022-01887-9>)
- effects of sudden deep snow in red desert on proghorn (<https://www.sciencedirect.com/science/article/abs/pii/S0960982225002957>)
- <https://www.nature.com/articles/s41467-023-37750-z>
- <https://esajournals.onlinelibrary.wiley.com/doi/full/10.1002/ecy.4238>
- detecting regime shifts (in lakes: Bjorndahl et al., 2022)

Environmental stochasticity and population dynamics

read Lande (1993)

Environmental stochasticity reduces habitats' energetic balance and carrying capacity (Chevin, Lande & Mace, 2010) as individuals struggle to rely on unpredictable resources and increase the instability of populations and communities (Lande, 1993). Additionally, stochastic environments reduce the ability of specialist species to adapt and evolve while selecting for more generalist species (Levins, 1974). Consequently, stochastic environments tend to have lower species diversity (**marcus?**).

Environmental stochasticity and fire ecology

Recent work by (**devries?**) has suggested that NDVI stochasticity is an important metric for evaluating forest recovery after large fires, while (**collinson?**) has suggested that variance in NDVI is an important predictor of the occurrence of forest fires due to ignition by lightning.

Environmental stochasticity and phenology

- Jonzén et al. (2006)
- effects can be compounded by cumulative effects (e.g., loss of species diversity, see Wolf, Zavaleta & Selman (2017))
- climate change affects plant phenology, but changes in situ may not match experimental estimates (Wolkovich et al., 2012)
- reproduction of painted lady butterfly (Stefanescu, Ubach & Wiklund, 2021)
- Wessling et al. (2018)

Useful references

- Bro-Jørgensen, Brown & Pettoirelli (2008): in the dry season preceding the rut topi
antelope (*Damaliscus lunatus*) respond to forage availability at the large scale but not
the fine scale
- Keith et al. (2008)
- Chevin, Lande & Mace (2010)
- Rickbeil et al. (2019)
- Mueller et al. (2008)
- Pettoirelli et al. (2005)
- Keys, Barnes & Carter (2021)
- Nilsen, Herfindal & Linnell (2005)
- Merkle et al. (2016)
- Tian et al. (2015)
- Huang et al. (2021)
- Fan & Liu (2016)
- Wang et al. (2019)
- Pease (2024)
- Xu et al. (2021)
- calibrating across sensors: <https://ieeexplore.ieee.org/abstract/document/6809983>

TO READ: * Site fidelity as a maladaptive behavior in the Anthropocene: [https://](https://esajournals.onlinelibrary.wiley.com/doi/full/10.1002/fee.2456)
* Pettoirelli N., Vik J.O., Mysterud A., Gaillard J.-M., Tucker C.J. & Stenseth N.Ch. (2005). Using the satellite-derived
NDVI to assess ecological responses to environmental change. Trends in Ecology & Evolution
20, 503–510. <https://doi.org/10.1016/j.tree.2005.05.011>

296 supplementary material:

297 • hex plots of pred vs obs for mean and var

298 • histograms of mean around flat map of biomes (change colors)

References

- Abrahms B, Hazen EL, Aikens EO, Savoca MS, Goldbogen JA, Bograd SJ, Jacox MG, Irvine LM, Palacios DM, and Mate BR. 2019. Memory and resource tracking drive blue whale migrations. *Proceedings of the National Academy of Sciences* 116: 5582–5587. [DOI [10.1073/pnas.1819031116](https://doi.org/10.1073/pnas.1819031116)]
- Aikens EO, Kauffman MJ, Merkle JA, Dwinnell SPH, Fralick GL, and Monteith KL. 2017. The greenscape shapes surfing of resource waves in a large migratory herbivore. *Ecology Letters* 20: 741–750. [DOI [10.1111/ele.12772](https://doi.org/10.1111/ele.12772)]
- Aikens EO, Merkle JA, Xu W, and Sawyer H. 2025. Pronghorn movements and mortality during extreme weather highlight the critical importance of connectivity. *Current Biology* 35: 1927–1934.e2. [DOI [10.1016/j.cub.2025.03.010](https://doi.org/10.1016/j.cub.2025.03.010)]
- Anderson SC, Branch TA, Cooper AB, and Dulvy NK. 2017. Black-swan events in animal populations. *Proceedings of the National Academy of Sciences* 114: 3252–3257. [DOI [10.1073/pnas.1611525114](https://doi.org/10.1073/pnas.1611525114)]
- Beck HE, McVicar TR, Van Dijk AIJM, Schellekens J, De Jeu RAM, and Bruijnzeel LA. 2011. Global evaluation of four AVHRR–NDVI data sets: Intercomparison and assessment against Landsat imagery. *Remote Sensing of Environment* 115: 2547–2563. [DOI [10.1016/j.rse.2011.05.012](https://doi.org/10.1016/j.rse.2011.05.012)]
- Bjorndahl JA, Gushulak CAC, Mezzini S, Simpson GL, Haig HA, Leavitt PR, and Finlay K. 2022. Abrupt changes in the physical and biological structure of endorheic upland lakes due to 8-m lake-level variation during the 20th century. *Limnology and Oceanography* 67: 1022–1039. [DOI [10.1002/lno.12054](https://doi.org/10.1002/lno.12054)]
- Botero CA, Weissing FJ, Wright J, and Rubenstein DR. 2015. Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences* 112: 184–189. [DOI [10.1073/pnas.1408589111](https://doi.org/10.1073/pnas.1408589111)]
- Broekman MJE, Hilbers JP, Hoeks S, Huijbregts MAJ, Schipper AM, and Tucker MA. 2024.

Environmental drivers of global variation in home range size of terrestrial and marine mammals. *Journal of Animal Ecology* 93: 488–500. [DOI [10.1111/1365-2656.14073](https://doi.org/10.1111/1365-2656.14073)]

Bro-Jørgensen J, Brown ME, and Pettorelli N. 2008. Using the satellite-derived normalized difference vegetation index (NDVI) to explain ranging patterns in a lek-breeding antelope: The importance of scale. *Oecologia* 158: 177–182. [DOI [10.1007/s00442-008-1121-z](https://doi.org/10.1007/s00442-008-1121-z)]

Burt WH. 1943. Territoriality and Home Range Concepts as Applied to Mammals. *Journal of Mammalogy* 24: 346. [DOI [10.2307/1374834](https://doi.org/10.2307/1374834)]

Charnov EL. 1976. Optimal foraging, the marginal value theorem. *Theoretical Population Biology* 9: 129–136. [DOI [10.1016/0040-5809\(76\)90040-X](https://doi.org/10.1016/0040-5809(76)90040-X)]

Chevin L-M, Lande R, and Mace GM. 2010. Adaptation, Plasticity, and Extinction in a Changing Environment: Towards a Predictive Theory. *PLoS Biology* 8: e1000357. [DOI [10.1371/journal.pbio.1000357](https://doi.org/10.1371/journal.pbio.1000357)]

Clerici N, Weissteiner CJ, and Gerard F. 2012. Exploring the Use of MODIS NDVI-Based Phenology Indicators for Classifying Forest General Habitat Categories. *Remote Sensing* 4: 1781–1803. [DOI [10.3390/rs4061781](https://doi.org/10.3390/rs4061781)]

Crameri F. 2018b. Scientific colour-maps. [DOI [10.5281/ZENODO.1287763](https://doi.org/10.5281/ZENODO.1287763)]

Crameri F. 2018a. Geodynamic diagnostics, scientific visualisation and StagLab 3.0. *Geoscientific Model Development* 11: 2541–2562. [DOI [10.5194/gmd-11-2541-2018](https://doi.org/10.5194/gmd-11-2541-2018)]

Crameri F, Shephard GE, and Heron PJ. 2020. The misuse of colour in science communication. *Nature Communications* 11: 5444. [DOI [10.1038/s41467-020-19160-7](https://doi.org/10.1038/s41467-020-19160-7)]

Falcón-Cortés A, Boyer D, Merrill E, Frair JL, and Morales JM. 2021. Hierarchical, Memory-Based Movement Models for Translocated Elk (*Cervus canadensis*). *Frontiers in Ecology and Evolution* 9: 702925. [DOI [10.3389/fevo.2021.702925](https://doi.org/10.3389/fevo.2021.702925)]

Fan X and Liu Y. 2016. A global study of NDVI difference among moderate-resolution satellite sensors. *ISPRS Journal of Photogrammetry and Remote Sensing* 121: 177–191. [DOI [10.1016/j.isprsjprs.2016.09.008](https://doi.org/10.1016/j.isprsjprs.2016.09.008)]

Frankham R and Brook BW. 2004. The importance of time scale in conservation biology

and ecology. *Annales Zoologici Fennici* 41: 459–463.

Frerebeau N. 2024. *Khroma: Colour Schemes for Scientific Data Visualization*. Pessac, France: Université Bordeaux Montaigne. [DOI [10.5281/zenodo.1472077](https://doi.org/10.5281/zenodo.1472077)]

Geremia C, Merkle JA, Eacker DR, Wallen RL, White PJ, Hebblewhite M, and Kauffman MJ. 2019. Migrating bison engineer the green wave. *Proceedings of the National Academy of Sciences* 116: 25707–25713. [DOI [10.1073/pnas.1913783116](https://doi.org/10.1073/pnas.1913783116)]

Goodchild MF and Quattrochi DA. 2023. *Scale in Remote Sensing and GIS*. New York: Routledge. [DOI [10.1201/9780203740170](https://doi.org/10.1201/9780203740170)]

Hmimina G, Dufrêne E, Pontailier J-Y, Delpierre N, Aubinet M, Caquet B, De Grandcourt A, Burban B, Flechard C, Granier A, Gross P, Heinesch B, Longdoz B, Moureaux C, Ourcival J-M, Rambal S, Saint André L, and Soudani K. 2013. Evaluation of the potential of MODIS satellite data to predict vegetation phenology in different biomes: An investigation using ground-based NDVI measurements. *Remote Sensing of Environment* 132: 145–158. [DOI [10.1016/j.rse.2013.01.010](https://doi.org/10.1016/j.rse.2013.01.010)]

Hollister J, Shah T, Nowosad J, Robitaille AL, Beck MW, and Johnson M. 2023. *Elevatr: Access Elevation Data from Various APIs*. [DOI [10.5281/zenodo.8335450](https://doi.org/10.5281/zenodo.8335450)]

Huang S, Tang L, Hupy JP, Wang Y, and Shao G. 2021. A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *Journal of Forestry Research* 32: 1–6. [DOI [10.1007/s11676-020-01155-1](https://doi.org/10.1007/s11676-020-01155-1)]

João T, João G, Bruno M, and João H. 2018. Indicator-based assessment of post-fire recovery dynamics using satellite NDVI time-series. *Ecological Indicators* 89: 199–212. [DOI [10.1016/j.ecolind.2018.02.008](https://doi.org/10.1016/j.ecolind.2018.02.008)]

Jonzén N, Lindén A, Ergon T, Knudsen E, Vik JO, Rubolini D, Piacentini D, Brinch C, Spina F, Karlsson L, Stervander M, Andersson A, Waldenström J, Lehikoinen A, Edvardsen E, Solvang R, and Stenseth NChr. 2006. Rapid Advance of Spring Arrival Dates in Long-Distance Migratory Birds. *Science* 312: 1959–1961. [DOI [10.1126/science.1126119](https://doi.org/10.1126/science.1126119)]

Keith DA, Akçakaya HR, Thuiller W, Midgley GF, Pearson RG, Phillips SJ, Regan HM,

- Araújo MB, and Rebelo TG. 2008. Predicting extinction risks under climate change: Coupling stochastic population models with dynamic bioclimatic habitat models. *Biology Letters* 4: 560–563. [DOI [10.1098/rsbl.2008.0049](https://doi.org/10.1098/rsbl.2008.0049)]
- Keys PW, Barnes EA, and Carter NH. 2021. A machine-learning approach to human footprint index estimation with applications to sustainable development. *Environmental Research Letters* 16: 044061. [DOI [10.1088/1748-9326/abe00a](https://doi.org/10.1088/1748-9326/abe00a)]
- Lande R. 1993. [Risks of Population Extinction from Demographic and Environmental Stochasticity and Random Catastrophes](#). *The American Naturalist* 142: 911–927.
- Levin SA. 1992. The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture. *Ecology* 73: 1943–1967. [DOI [10.2307/1941447](https://doi.org/10.2307/1941447)]
- Levins RA. 1974. *Evolution in changing environments: Some theoretical explorations*. Princeton, NJ: Princeton Univ. Press.
- Logares R and Nuñez M. 2012. Black Swans in ecology and evolution: The importance of improbable but highly influential events. *Ideas in Ecology and Evolution*. [DOI [10.4033/iee.2012.5.5.n](https://doi.org/10.4033/iee.2012.5.5.n)]
- Mallegowda P, Rengaiyan G, Krishnan J, and Niphadkar M. 2015. Assessing Habitat Quality of Forest-Corridors through NDVI Analysis in Dry Tropical Forests of South India: Implications for Conservation. *Remote Sensing* 7: 1619–1639. [DOI [10.3390/rs70201619](https://doi.org/10.3390/rs70201619)]
- Matsushita B, Yang W, Chen J, Onda Y, and Qiu G. 2007. Sensitivity of the Enhanced Vegetation Index (EVI) and Normalized Difference Vegetation Index (NDVI) to Topographic Effects: A Case Study in High-density Cypress Forest. *Sensors* 7: 2636–2651. [DOI [10.3390/s7112636](https://doi.org/10.3390/s7112636)]
- Merkle JA, Monteith KL, Aikens EO, Hayes MM, Hersey KR, Middleton AD, Oates BA, Sawyer H, Scurlock BM, and Kauffman MJ. 2016. Large herbivores surf waves of green-up during spring. *Proceedings of the Royal Society B: Biological Sciences* 283: 20160456. [DOI [10.1098/rspb.2016.0456](https://doi.org/10.1098/rspb.2016.0456)]
- Mezzini S, Fleming CH, Medici EP, and Noonan MJ. 2025. How resource abundance and

resource stochasticity affect organisms' range sizes. *Movement Ecology* 13: 20. [DOI [10.1186/s40462-025-00546-5](https://doi.org/10.1186/s40462-025-00546-5)]

Middleton AD, Merkle JA, McWhirter DE, Cook JG, Cook RC, White PJ, and Kauffman MJ. 2018. Green-wave surfing increases fat gain in a migratory ungulate. *Oikos* 127: 1060–1068. [DOI [10.1111/oik.05227](https://doi.org/10.1111/oik.05227)]

Mueller T, Olson KA, Fuller TK, Schaller GB, Murray MG, and Leimgruber P. 2008. In search of forage: Predicting dynamic habitats of Mongolian gazelles using satellite-based estimates of vegetation productivity. *Journal of Applied Ecology* 45: 649–658. [DOI [10.1111/j.1365-2664.2007.01371.x](https://doi.org/10.1111/j.1365-2664.2007.01371.x)]

Mutti PR, Lúcio PS, Dubreuil V, and Bezerra BG. 2020. NDVI time series stochastic models for the forecast of vegetation dynamics over desertification hotspots. *International Journal of Remote Sensing* 41: 2759–2788. [DOI [10.1080/01431161.2019.1697008](https://doi.org/10.1080/01431161.2019.1697008)]

Nathan R, Getz WM, Revilla E, Holyoak M, Kadmon R, Saltz D, and Smouse PE. 2008. A movement ecology paradigm for unifying organismal movement research. *Proceedings of the National Academy of Sciences* 105: 19052–19059. [DOI [10.1073/pnas.0800375105](https://doi.org/10.1073/pnas.0800375105)]

Nathan R, Monk CT, Arlinghaus R, Adam T, Alós J, Assaf M, Baktoft H, Beardsworth CE, Bertram MG, Bijleveld AI, Brodin T, Brooks JL, Campos-Candela A, Cooke SJ, Gjelland KØ, Gupte PR, Harel R, Hellström G, Jeltsch F, Killen SS, Klefoth T, Langrock R, Lennox RJ, Lourie E, Madden JR, Orchan Y, Pauwels IS, Říha M, Roeleke M, Schlägel UE, Shohami D, Signer J, Toledo S, Vilk O, Westrelin S, Whiteside MA, and Jarić I. 2022. Big-data approaches lead to an increased understanding of the ecology of animal movement. *Science* 375: eabg1780. [DOI [10.1126/science.abg1780](https://doi.org/10.1126/science.abg1780)]

Nilsen EB, Herfindal I, and Linnell JDC. 2005. Can intra-specific variation in carnivore home-range size be explained using remote-sensing estimates of environmental productivity? *Écoscience* 12: 68–75. [DOI [10.2980/i1195-6860-12-1-68.1](https://doi.org/10.2980/i1195-6860-12-1-68.1)]

Okabe M and Ito K. 2008. [Color universal design \(CUD\): How to make figures and presentations that are friendly to colorblind people.](#)

- Olson DM and Dinerstein E. 2002. The Global 200: Priority Ecoregions for Global Conservation. *Annals of the Missouri Botanical Garden* 89: 199. [DOI [10.2307/3298564](https://doi.org/10.2307/3298564)]
- Pease BS. 2024. Ecological scales of effect vary across space and time. *Ecography* 2024: e07163. [DOI [10.1111/ecog.07163](https://doi.org/10.1111/ecog.07163)]
- Pedersen EJ, Miller DL, Simpson GL, and Ross N. 2019. Hierarchical generalized additive models in ecology: An introduction with mgcv. *PeerJ* 7: e6876. [DOI [10.7717/peerj.6876](https://doi.org/10.7717/peerj.6876)]
- Pettorelli N, Ryan S, Mueller T, Bunnefeld N, Jedrzejewska B, Lima M, and Kausrud K. 2011. The Normalized Difference Vegetation Index (NDVI): Unforeseen successes in animal ecology. *Climate Research* 46: 15–27. [DOI [10.3354/cr00936](https://doi.org/10.3354/cr00936)]
- Pettorelli N, Vik JO, Mysterud A, Gaillard J-M, Tucker CJ, and Stenseth NChr. 2005. Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution* 20: 503–510. [DOI [10.1016/j.tree.2005.05.011](https://doi.org/10.1016/j.tree.2005.05.011)]
- Polansky L, Kilian W, and Wittemyer G. 2015. Elucidating the significance of spatial memory on movement decisions by African savannah elephants using state–space models. *Proceedings of the Royal Society B: Biological Sciences* 282: 20143042. [DOI [10.1098/rspb.2014.3042](https://doi.org/10.1098/rspb.2014.3042)]
- Polazzo F, Limberger R, Pennekamp F, Ross SRP-J, Simpson GL, and Petchey OL. 2024. Measuring the Response Diversity of Ecological Communities Experiencing Multifarious Environmental Change. *Global Change Biology* 30: e17594. [DOI [10.1111/gcb.17594](https://doi.org/10.1111/gcb.17594)]
- R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing.
- Ranc N, Cagnacci F, and Moorcroft PR. 2022. Memory drives the formation of animal home ranges: Evidence from a reintroduction. *Ecology Letters* 25: 716–728. [DOI [10.1111/ele.13869](https://doi.org/10.1111/ele.13869)]
- Reid CE, Kubzansky LD, Li J, Shmool JL, and Clougherty JE. 2018. It’s not easy assessing greenness: A comparison of NDVI datasets and neighborhood types and their associations with self-rated health in New York City. *Health & Place* 54: 92–101. [DOI

460 [10.1016/j.healthplace.2018.09.005](https://doi.org/10.1016/j.healthplace.2018.09.005)]

- 461 Rickbeil GJM, Merkle JA, Anderson G, Atwood MP, Beckmann JP, Cole EK, Courtemanch
462 AB, Dewey S, Gustine DD, Kauffman MJ, McWhirter DE, Mong T, Proffitt K, White PJ,
463 and Middleton AD. 2019. Plasticity in elk migration timing is a response to changing en-
464 vironmental conditions. *Global Change Biology* 25: 2368–2381. [DOI [10.1111/gcb.14629](https://doi.org/10.1111/gcb.14629)]
- 465 Riotte-Lambert L and Matthiopoulos J. 2020. Environmental Predictability as a Cause and
466 Consequence of Animal Movement. *Trends in Ecology & Evolution* 35: 163–174. [DOI
467 [10.1016/j.tree.2019.09.009](https://doi.org/10.1016/j.tree.2019.09.009)]
- 468 Rizzuto M, Leroux SJ, Vander Wal E, Richmond IC, Heckford TR, Balluffi-Fry J, and
469 Wiersma YF. 2021. Forage stoichiometry predicts the home range size of a small terres-
470 trial herbivore. *Oecologia* 197: 327–338. [DOI [10.1007/s00442-021-04965-0](https://doi.org/10.1007/s00442-021-04965-0)]
- 471 Rouse JW Jr., Haas RH, Schell JA, and Deering DW. 1974a. Monitoring Vegetation Systems
472 in the Great Plains with ERTS. NASA Special Publication. 309.
- 473 Rouse JW Jr., Haas RH, Schell JA, and Deering DW. 1974b. *Monitoring the vernal ad-
474 vancement and retrogradation (green wave effect) of natural vegetation*. Remote Sensing
475 Center, Texas A&M University, College Station.
- 476 Sawyer H, Merkle JA, Middleton AD, Dwinnell SPH, and Monteith KL. 2019. Migratory
477 plasticity is not ubiquitous among large herbivores. *Journal of Animal Ecology* 88: 450–
478 460. [DOI [10.1111/1365-2656.12926](https://doi.org/10.1111/1365-2656.12926)]
- 479 Simpson GL. 2018. Modelling Palaeoecological Time Series Using Generalised Additive
480 Models. *Frontiers in Ecology and Evolution* 6: 149. [DOI [10.3389/fevo.2018.00149](https://doi.org/10.3389/fevo.2018.00149)]
- 481 Son NT, Chen CF, Chen CR, Minh VQ, and Trung NH. 2014. A comparative analysis of
482 multitemporal MODIS EVI and NDVI data for large-scale rice yield estimation. *Agri-
483 cultural and Forest Meteorology* 197: 52–64. [DOI [10.1016/j.agrformet.2014.06.007](https://doi.org/10.1016/j.agrformet.2014.06.007)]
- 484 Southwood TRE. 1977. Habitat, the Templet for Ecological Strategies? *The Journal of
485 Animal Ecology* 46: 336. [DOI [10.2307/3817](https://doi.org/10.2307/3817)]
- 486 Stefanescu C, Ubach A, and Wiklund C. 2021. Timing of mating, reproductive status and

resource availability in relation to migration in the painted lady butterfly. *Animal Behaviour* 172: 145–153. [DOI [10.1016/j.anbehav.2020.12.013](https://doi.org/10.1016/j.anbehav.2020.12.013)]

Steixner-Kumar S and Gläscher J. 2020. Strategies for navigating a dynamic world. *Science* 369: 1056–1057. [DOI [10.1126/science.abd7258](https://doi.org/10.1126/science.abd7258)]

Stephens DW and Charnov EL. 1982. Optimal foraging: Some simple stochastic models. *Behavioral Ecology and Sociobiology* 10: 251–263. [DOI [10.1007/BF00302814](https://doi.org/10.1007/BF00302814)]

Tarnavsky E, Garrigues S, and Brown ME. 2008. Multiscale geostatistical analysis of AVHRR, SPOT-VGT, and MODIS global NDVI products. *Remote Sensing of Environment* 112: 535–549. [DOI [10.1016/j.rse.2007.05.008](https://doi.org/10.1016/j.rse.2007.05.008)]

Tian F, Fensholt R, Verbesselt J, Grogan K, Horion S, and Wang Y. 2015. Evaluating temporal consistency of long-term global NDVI datasets for trend analysis. *Remote Sensing of Environment* 163: 326–340. [DOI [10.1016/j.rse.2015.03.031](https://doi.org/10.1016/j.rse.2015.03.031)]

Tucker CJ. 1979. Red and photographic infrared linear combinations for monitoring vegetation. *Remote Sensing of Environment* 8: 127–150. [DOI [10.1016/0034-4257\(79\)90013-0](https://doi.org/10.1016/0034-4257(79)90013-0)]

Tucker CJ, Elgin JH, McMurtrey JE, and Fan CJ. 1979. Monitoring corn and soybean crop development with hand-held radiometer spectral data. *Remote Sensing of Environment* 8: 237–248. [DOI [10.1016/0034-4257\(79\)90004-x](https://doi.org/10.1016/0034-4257(79)90004-x)]

Vermote E and NOAA CDR Program. 2018. NOAA Climate Data Record (CDR) of AVHRR Normalized Difference Vegetation Index (NDVI), Version 5. [DOI [10.7289/V5ZG6QH9](https://doi.org/10.7289/V5ZG6QH9)]

Vermote E and NOAA CDR Program. 2022. NOAA Climate Data Record (CDR) of VIIRS Normalized Difference Vegetation Index (NDVI), Version 1. [DOI [10.25921/GAKH-ST76](https://doi.org/10.25921/GAKH-ST76)]

Wang X, Cao L, Fox AD, Fuller R, Griffin L, Mitchell C, Zhao Y, Moon O-K, Cabot D, Xu Z, Batbayar N, Kölzsch A, Van Der Jeugd HP, Madsen J, Chen L, and Nathan R. 2019. Stochastic simulations reveal few green wave surfing populations among spring migrating herbivorous waterfowl. *Nature Communications* 10: 2187. [DOI [10.1038/s41467-019-09971-8](https://doi.org/10.1038/s41467-019-09971-8)]

Wessling EG, Deschner T, Mundry R, Pruetz JD, Wittig RM, and Köhl HS. 2018. Sea-

- sonal Variation in Physiology Challenges the Notion of Chimpanzees (*Pan troglodytes* versus) as a Forest-Adapted Species. *Frontiers in Ecology and Evolution* 6: 60. [DOI [10.3389/fevo.2018.00060](https://doi.org/10.3389/fevo.2018.00060)]
- Wolf AA, Zavaleta ES, and Selmants PC. 2017. Flowering phenology shifts in response to biodiversity loss. *Proceedings of the National Academy of Sciences* 114: 3463–3468. [DOI [10.1073/pnas.1608357114](https://doi.org/10.1073/pnas.1608357114)]
- Wolkovich EM, Cook BI, Allen JM, Crimmins TM, Betancourt JL, Travers SE, Pau S, Regetz J, Davies TJ, Kraft NJB, Ault TR, Bolmgren K, Mazer SJ, McCabe GJ, McGill BJ, Parmesan C, Salamin N, Schwartz MD, and Cleland EE. 2012. Warming experiments underpredict plant phenological responses to climate change. *Nature* 485: 494–497. [DOI [10.1038/nature11014](https://doi.org/10.1038/nature11014)]
- Wood SN. 2017. *Generalized additive models: An introduction with R*. Boca Raton: CRC Press/Taylor & Francis Group.
- Wood SN, Goude Y, and Shaw S. 2015. Generalized additive models for large data sets. *Journal of the Royal Statistical Society: Series C (Applied Statistics)* 64: 139–155. [DOI [10.1111/rssc.12068](https://doi.org/10.1111/rssc.12068)]
- Wood SN, Li Z, Shaddick G, and Augustin NH. 2017. Generalized Additive Models for Gigadata: Modeling the U.K. Black Smoke Network Daily Data. *Journal of the American Statistical Association* 112: 1199–1210. [DOI [10.1080/01621459.2016.1195744](https://doi.org/10.1080/01621459.2016.1195744)]
- Xu W, Barker K, Shawler A, Van Scoyoc A, Smith JA, Mueller T, Sawyer H, Andreozzi C, Bidder OR, Karandikar H, Mumme S, Templin E, and Middleton AD. 2021. The plasticity of ungulate migration in a changing world. *Ecology* 102: e03293. [DOI [10.1002/ecy.3293](https://doi.org/10.1002/ecy.3293)]
- Zhang X, Hu Y, Zhuang D, Qi Y, and Ma X. 2009. NDVI spatial pattern and its differentiation on the Mongolian Plateau. *Journal of Geographical Sciences* 19: 403–415. [DOI [10.1007/s11442-009-0403-7](https://doi.org/10.1007/s11442-009-0403-7)]