

DENVar: A Global Dynamic Estimate of NDVI Variance

Stefano Mezzini^{1,2}, Gavin L Simpson³, and Michael J Noonan^{1,2,4,*}

¹Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

²Department of Biology, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

³Department of Animal and Veterinary Sciences, Aarhus University, Denmark.

⁴Department of Computer Science, Math, Physics, and Statistics, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

*Address correspondence to: michael.noonan@ubc.ca

Abstract

Remote-sensing proxies of habitat productivity such as the Normalized Difference Vegetation Index (NDVI) have helped develop and support many hypotheses and projects. However, historically, users tended to focus solely on trends in mean productivity. In recent years, the variance around the mean (i.e., stochasticity) has gained more attention, particularly as climate change and human-induced rapid environmental change transform the ecosystems species have evolved in and adapted to. Still, calculating trends in NDVI stochasticity requires careful modeling, since stochasticity estimates are conditional on the estimated mean and the scale at which the data are collected and modeled. We thus present a new global Dynamic Estimate of NDVI Variance, *DENVar*. We estimate spatiotemporal trends in NDVI mean and variance using Generalized Additive Models, which provide a flexible yet transparent and interpretable model structure. We show that DENVar correlates nonlinearly with **XXX** and can be used to test many hypotheses related to environmental stochasticity, such as optimal foraging theory and regime shifts as well as for conservation. We provide a publicly and freely available repository for downloading rasters of daily and long-term averages of DENVar and the related estimates of mean NDVI.

1 Introduction

The Normalized Difference Vegetation Index (NDVI) is a proxy of primary productivity used in a wide range of disciplines, including ecology [6, 7], landscape phenology [8, 9], agriculture [10], forestry [11], and conservation [12, 13]. High-resolution, remotely-sensed NDVI data have allowed analysts to estimate plant biomass without the need for fieldwork [14, 15] and leverage such estimates for studying animal movement behavior [8, 16]. Still, most work is limited to quantifying changes in and responses to mean NDVI while ignoring changes in the variance around the mean.

Environmental stochasticity has been studied for decades [17–20], but recent work has drawn more attention to the effects environmental stochasticity has at a wide range of scales, from individual organisms [21, 22] to across landscapes [13, 23, 24]. However, estimating trends in NDVI variance requires more nuanced modeling than estimating changes in mean NDVI, particularly when environments are temporally dynamic and spatially heterogeneous. The scale at which processes are observed determine how individuals’ perception of and ability to respond to change [25]. A forest fire is severely disruptive and deadly for many insects [26], while birds and large mammals may not struggle to escape the fire and or establish a new range [27, 28]. Similarly, the effects of the fire recovery will depend on its duration relative to other organisms’ phenology and lifespan [29]. Therefore, distinguishing between a change in mean NDVI and a stochastic fluctuation requires a conscious decision about the scale at which to consider the data.

Add a paragraph on what effects stochasticity may have/why it’s important?

We present a new measure of environmental stochasticity, which we name the Dynamic Estimate of NDVI Variance (DENVar). We estimate DENVar using 44 years of NDVI data from the AVHRR and VIIRS collection of sensors and Generalized Additive Models (GAMs). Although satellites with AVHRR and VIIRS sensors provide data at a coarser spatial resolution (0.05°) than other satellites (e.g., 250m for MODIS), they are the only sensors to provide data at a daily frequency and for more than 40 years. Our use of GAMs provides a flexible yet interpretable approach for estimating trends in mean and variance in NDVI without the rigidity of other more traditional methods (e.g., ARIMA) or the opaqueness of machine learning models such as neural networks [30, 31]. We illustrate the relationships between DENVar and multiple, relevant ecological metrics, and we provide access to a free repository of daily, global estimates of DENVar and the corresponding estimates of mean NDVI.

2 Materials and Methods

2.1 Choice of color palettes

We represent NDVI using a modified version of Fabio Crameri’s divergent *bukavu* palette [1, 32], which has high-contrast for deuteranope and protanope vision. For NDVI values between -0.1 and 1, the colors also have sufficient contrast for the colors to also be distinguishable by strong tritanope and achromatic vision. Fig. A1 contains an approximate representation of how each vision type would perceive each color palette used. We obtained all color palettes with three or more colors

from the `khroma` package (v. 1.14.0, [4]) for R (v. 4.4.1, [33]).

2.2 Experimental and Technical Design

- include a diagram to show the entire experimental design and illustrate the most significant elements: materials, treatments, measurements, data collection, methods of data analysis
- This will facilitate the editors and reviewers to understand and follow the whole concept, design, and results.

We downloaded NDVI image composites directly from the servers for data from AVHRR sensors (1981-06-24 to 2013-12-31; <https://ladsweb.modaps.eosdis.nasa.gov/archive/allData>; [34]) and VIIRS sensors (2014-01-01 to 2025-06-29; <https://ncei.noaa.gov/data/>; [35]). Fig. A2 shows the availability of rasters over time for both sets of sensors. We removed all pixels labeled as "cloudy" for AVHRR rasters and "probably cloudy" or "confidently cloudy" for VIIRS rasters. Additionally, we removed pixels with erroneously high NDVI values at high latitudes between September and March using latitudinal cutoffs based on the monthly empirical cumulative distribution function of NDVI across latitude (Fig. B2). We did not remove values at latitudes $< 60^\circ\text{S}$ since we fully excluded Antarctica from the analyses.

Elevation above sea level was obtained using the `get_elev_raster()` function from the `elevatr` package for R (v. 0.99.0, [36]) with an original resolution of 0.076° (Fig. A6). To remove any biases towards negative NDVI values due to water, we removed all raster cells with $> 40\%$ water cover. We estimated the proportion of water in each cell by rasterizing a shapefile of main water bodies at a resolution of 0.05° (Fig. A7).

To ensure the number of rows for each dataset was below the maximum data frame size in R ($3.17 \times 10^{10} > 2^{31} - 1 \approx 0.21 \times 10^{10}$), we reduced the dataset size by splitting the dataset across four spatial groups (Fig. A4) and spatiotemporally averaged the (cleaned) rasters by a factor of four. The final dataset thus had one raster every four days with a spatial resolution of $0.05^\circ \times 4 = 0.2^\circ$. Despite the substantial reduction in dataset size, the dataset creation and modeling were still challenging for the machine used, which had 2.2 TB of RAM and an Intel Xeon Platinum 8462Y+ processor (32 cores).

2.3 Statistical Analysis

We estimated spatiotemporal trends in mean NDVI using Generalized Additive Models (GAMs) via the `mgcv` package for R (v. 1.9-3, [31]). To reduce modeling fitting times with negligible losses to model accuracy, we used the `bam()` function with fast REstricted Marginal Likelihood (`method = fREML`) and covariate discretization (`discrete = TRUE`; [37, 38]). Ideally, NDVI should be modeled using beta location-scale models (after the linear transformation $Y^* = \frac{Y+1}{2}$) to account for the fact that: (1) NDVI is bounded between -1 and 1, and (2) the variance in NDVI is dependent on the mean (and vice-versa), since ecosystems with very low NDVI (e.g., rock, ice, or concrete) or very high NDVI (e.g., dense forest) tend to have lower variance in NDVI. However, preliminary tests showed that fitting times for beta models were prohibitive, especially for beta location-scale models. In contrast,

101 Gaussian models fit substantially faster and provided very similar spatialtemporal estimates of mean
 102 NDVI (see Appendix B). For each region shown in Fig. A4, the model for the mean NDVI had the
 103 structure below:

$$\begin{cases} \text{NDVI} \sim \text{Normal}(\mu, \sigma^2) \\ \mu = \beta_0 + f_1(\text{prop_water}) + f_2(\text{elev_m}) + f_3(\text{doy}) + f_4(\text{year}) + f_5(\text{lat}, \text{long}) + \\ \quad f_{6T}(\text{doy}, \text{year}) + f_{7T}(\text{doy}, \text{lat}, \text{long}) + f_{8T}(\text{year}, \text{lat}, \text{long}) \end{cases}, \quad (1)$$

104 where, for a given (aggregated) pixel, **prop_water** is the proportion of water within the pixel, **elev_m**
 105 is elevation above sea level in meters, **doy** is integer day of year (1-366), **year** is integer year (1981-
 106 2025), and **lat** and **long** are latitude and longitude, respectively. $f_j(\cdot)$ indicate the j^{th} marginal
 107 smooth functions, while $f_{jT}(\cdot)$ indicate the j^{th} tensor product interaction terms. Latitude and
 108 longitude were modeled using a spline-on-the-sphere basis (**bs** = 'sos'; [31]), while seasonal terms
 109 used cyclical cubic splines (**bs** = 'cc') with edge knots 0.5 and 366.5.

110 **HERE**

111 A second GAM estimated the variance in NDVI around the mean estimated by the model above.
 112 The model had an identical structure to the model for the mean, with the exception that (1) the
 113 response variable was the squared residual from the first model, such that the model estimated the
 114 mean squared residual (i.e., the variance) for a given point in time and space, and (2) a smooth term
 115 of the estimated mean NDVI. *Estimating the variance as the spatially or spatiotemporally varying*
 116 *mean squared residuals included the pixel-level squared bias.* The model is available below:

117 **formula**

118 3 Results

119 variance is likely overestimated in areas near sharp changes in NDVI, but this depends on scale
 120 Figures:

- 121 • two orthographic projections of DENVar: northern and southern hemispheres on spring solstice
- 122 OR Robinson projection of the world with biomes to see if fill matches biome boundaries
- 123 • plot grid of mean over space, **s(year)**, **s(doy)**, **s(elev)**
- 124 • plot grid of DENVar over space, **s(year)**, **s(doy)**, **s(elev)**, **s(mean NDVI)**


- 125 • hex plots of pred vs obs for mean and var
- 126 • hex plots of DENVar with other variables (add temperature, annual precip, extreme weather
127 event, fires, invasive species, max body size, disease occurrence, ...)
- 128 • supplementary material: histograms of mean around flat map of biomes (already created, but
129 change colors?)

130 **3.1 Correlations with other metrics**

- 131 • temperature
- 132 • seasonal temperature range
- 133 • precip
- 134 • seasonal temperature range
- 135 • spp diversity
- 136 • gross primary productivity
- 137 • max animal weight
- 138 • <https://silvis.forest.wisc.edu/globalwui/>
- 139 • <https://silvis.forest.wisc.edu/data/dhis/>: NDVI16 (Normalized Difference Vegetation Index),
140 EVI16 (Enhanced Vegetation Index), FPAR8 (Fraction absorbed Photosynthetically Active
141 Radiation), LAI8 (Leaf Area Index), GPP8 (Gross Primary Productivity)

142 **3.2 Obtaining the data**

- 143 • Data should be available as daily rasters on a github repo?
- 144 • make an app for data annotation?



4 Discussion

We have presented a new spatiotemporally dynamic metric of environmental stochasticity, which we have called the Dynamic Estimate of NDVI Variance, DENVar. In this section, we discuss the interpretation of DENVar in greater depth, and we argue the advantages and limitations of the statistical methods used. We conclude by suggesting some use cases of DENVar within animal movement ecology, landscape ecology, and phenology, but we invite readers to consider how DENVar may be of use in their field and adjacent disciplines.

4.1 Interpreting DENVar: the importance of scale

difficulties in estimating variance

what counts as a change? predictable vs stochastic changes

smoothness of mean impacts smoothness and size of variance: depends on scale of interest (e.g., animals' ability to respond to, learn, and predict changes in the mean). Changes not attributed to trends in the mean are stochastic

DENVar is a good metric as long as the spatiotemporal scale of interest is similar or larger to that of DENVar. May not work well for small-scale processes such as the movement dynamics of a small rodent, the phenology of a small orchard when spatiotemporal heterogeneity is finer than DENVar's resolution

4.2 Environmental stochasticity and animal movement ecology

The field of animal movement ecology has long shown that animals alter their movement in response to resource abundance or availability [39–42]. More recently, movement ecologists have started focusing more on resource stochasticity [18, 22, 43], but it is worth distinguishing between resource stochasticity and seasonality [44]. Seasonality is generally best interpreted as a cyclical change in the mean that many animals can learn to predict, as in the case of green-wave surfing [45, 46]. In contrast, environmental stochasticity refers to unpredictable changes, such as forest fires [ref](#), sudden snow storms [47], and unusual phenology [ref](#). Animals also respond to stochastic changes [21], but their ability to adapt depends on behavioral plasticity [48, 49], and memory [50–54].

evolutionary tipping points [55]

multifarious change [56]

extreme events, black swans [57, 58]

- migration routes are key habitat (Ortega et al., in prep), especially in stochastic environments

- can fail to adapt to changes in conditions [59]

- <https://www.sciencedirect.com/science/article/pii/S0960982220308484>

- <https://onlinelibrary.wiley.com/doi/full/10.1111/gcb.15169>

- industrial development affects animals’ ability to respond to both predictable and unpredictable change (<https://www.nature.com/articles/s41559-022-01887-9>)

- effects of sudden deep snow in red desert on proghorn (<https://www.sciencedirect.com/science/article/abs/pii/S0960982220308484>)

- <https://www.nature.com/articles/s41467-023-37750-z>

- <https://esajournals.onlinelibrary.wiley.com/doi/full/10.1002/ecy.4238>

- detecting regime shifts (in lakes: [60])

4.3 Environmental stochasticity and population dynamics

read [lande_risks_1993](#)

Environmental stochasticity reduces habitats’ energetic balance and carrying capacity [61] as individuals struggle to rely on unpredictable resources and increase the instability of populations and

188 communities [19]. Additionally, stochastic environments reduce the ability of specialist species to
189 adapt and evolve while selecting for more generalist species [17]. Consequently, stochastic environ-
190 ments tend to have lower species diversity [13].

191 Recent work by Devries has suggested that NDVI stochasticity is an important metric for evaluating
192 forest recovery after large fires, while Collinson has suggested that variance in NDVI is an important
193 predictor of the occurrence of lightning-induced forest fires.

- 194 • jonzen_rapid_2006
- 195 • effects can be compounded by cumulative effects (e.g., loss of species diversity; [62])
- 196 • climate change affects plant phenology, but changes in situ may not match experimental esti-
197 mates [63]
- 198 • reproduction of painted lady butterfly [64]
- 199 • wessling_seasonal_2018

200 4.4 Environmental stochasticity and fire ecology

201 4.5 Environmental stochasticity and phenology

202 5 Other useful references

- 203 • **READ**: Site fidelity as a maladaptive behavior in the Anthropocene: <https://esajournals.onlinelibrary.wiley.com/doi>
- 204 • **READ**: Pettorelli N., Vik J.O., Mysterud A., Gaillard J.-M., Tucker C.J. & Stenseth N.Chr.
205 (2005). Using the satellite-derived NDVI to assess ecological responses to environmental
206 change. *Trends in Ecology & Evolution* 20, 503–510. <https://doi.org/10.1016/j.tree.2005.05.011>
- 207 • bro-jorgensen_using_2008: in the dry season preceding the rut topi antelope (*Damaliscus*
208 *lunatus*) respond to forage availability at the large scale but not the fine scale
- 209 • keith_predicting_2008
- 210 • chevin_adaptation_2010
- 211 • rickbeil_plasticity_2019
- 212 • mueller_search_2008
- 213 • pettorelli_using_2005
- 214 • nilsen_can_2005
- 215 • merkle_large_2016
- 216 • tian_evaluating_2015
- 217 • huang_commentary_2021
- 218 • fan_global_2016
- 219 • wang_stochastic_2019
- 220 • pease_ecological_2024
- 221 • xu_plasticity_2021
- 222 • calibrating across sensors: <https://ieeexplore.ieee.org/abstract/document/6809983>

223 Acknowledgments

224 Anyone who made a contribution to the research or manuscript, but who is not a listed author,
225 should be acknowledged (with their permission). Types of acknowledgements include:

226 General

227 Thank others for any contributions, whether it be direct technical help or indirect assistance

228 Author Contributions

229 Describe contributions of each author to the paper, using the first initial and full last name.

230 Examples:

231 “S. Zhang conceived the idea and designed the experiments.”

232 “E. F. Mustermann and J. F. Smith conducted the experiments.”

233 “All authors contributed equally to the writing of the manuscript.”

234 Funding

235 Name financially supporting bodies (written out in full), followed by the funding awardee and asso-
236 ciated grant numbers (if applicable) in square brackets.

237 Example:

238 “This work was supported by the Engineering and Physical Sciences Research Council grant
239 numbers xxxx, yyyy; the National Science Foundation grant number zzzz; and a Leverhulme Trust
240 Research Project Grant.”

241 If the research did not receive specific funding, but was performed as part of the employment
242 of the authors, please name this employer. If the funder was involved in the manuscript writing,
243 editing, approval, or decision to publish, please declare this.

244 Conflicts of Interest

245 Conflicts of interest (COIs, also known as “competing interests”) occur when issues outside research
246 could be reasonably perceived to affect the neutrality or objectivity of the work or its assessment.

247 Authors must declare all potential interests – whether or not they actually had an influence – in a
248 ‘Conflicts of Interest’ section, which should explain why the interest may be a conflict. Authors must
249 declare current or recent funding (including for Article Processing Charges) and other payments,
250 goods or services that might influence the work. All funding, whether a conflict or not, must be
251 declared in a “Funding Statement.” The involvement of anyone other than the authors who 1) has
252 an interest in the outcome of the work; 2) is affiliated to an organization with such an interest; or 3)
253 was employed or paid by a funder, in the commissioning, conception, planning, design, conduct, or
254 analysis of the work, the preparation or editing of the manuscript, or the decision to publish must
255 be declared.

256 If there are none, the authors should state “The author(s) declare(s) that there is no conflict of
257 interest regarding the publication of this article.” Submitting authors are responsible for coauthors
258 declaring their interests. Declared conflicts of interest will be considered by the editor and reviewers
259 and included in the published article.

260 **Data Availability**

261 A data availability statement is compulsory for all research articles. This statement describes
262 whether and how others can access the data supporting the findings of the paper, including 1)
263 what the nature of the data is, 2) where the data can be accessed, and 3) any restrictions on data
264 access and why.

265 If data are in an archive, include the accession number or a placeholder for it. Also include any
266 materials that must be obtained through a Material Transfer Agreements (MTA).

267 **Supplementary Materials**

268 Describe any supplementary materials submitted with the manuscript (e.g., audio files, video clips
269 or datasets).

270 Please group supplementary materials in the following order: materials and methods, figures,
271 tables, and other files (such as movies, data, interactive images, or database files).

272 Example: Fig. S1. Title of the first supplementary figure.

273 Fig. S2. Title of the second supplementary figure.

274 Table S1. Title of the first supplementary table.

275 Data file S1. Title of the first supplementary data file.

276 Movie S1. Title of the first supplementary movie.

277 Be sure to submit all supplementary materials with the manuscript and remember to reference
278 the supplementary materials at appropriate points within the manuscript. We recommend citing
279 specific items, rather than referring to the supplementary materials in general, for example: “See
280 Figures S1-S10 in the Supplementary Material for comprehensive image analysis.”

281 A link to access the supplementary materials will be provided in the published article.

282 Supplementary Materials may include additional author notes—for example, a list of group
283 authors.

References

1. Crameri F. Geodynamic diagnostics, scientific visualisation and StagLab 3.0. *Geoscientific Model Development* 2018;11:2541–62.
2. Crameri F. Scientific colour-maps. In collab. with Shephard GE. Version 3.0.4. Language: en. 2018. DOI: 10.5281/ZENODO.1287763. URL: <https://zenodo.org/record/1287763> (visited on 04/04/2025).
3. Okabe M and Ito K. Color universal design (CUD): How to make figures and presentations that are friendly to colorblind people. 2008. URL: <https://jfly.uni-koeln.de/color/#pallet>.
4. Frerebeau N. khroma: Colour Schemes for Scientific Data Visualization. Pessac, France: Université Bordeaux Montaigne, 2024. DOI: 10.5281/zenodo.1472077. URL: <https://packages.tesselle.org/khroma/>.
5. Olson DM, Dinerstein E, Wikramanayake ED, et al. Terrestrial Ecoregions of the World: A New Map of Life on Earth. *BioScience* 2001;51. Publisher: Oxford University Press (OUP):933.
6. Pettorelli N, Vik JO, Mysterud A, Gaillard JM, Tucker CJ, and Stenseth NC. Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution* 2005;20:503–10.
7. Pettorelli N, Ryan S, Mueller T, et al. The Normalized Difference Vegetation Index (NDVI): unforeseen successes in animal ecology. *Climate Research* 2011;46:15–27.
8. Merkle JA, Monteith KL, Aikens EO, et al. Large herbivores surf waves of green-up during spring. *Proceedings of the Royal Society B: Biological Sciences* 2016;283:20160456.
9. Hmimina G, Dufrêne E, Pontailier JY, et al. Evaluation of the potential of MODIS satellite data to predict vegetation phenology in different biomes: An investigation using ground-based NDVI measurements. *Remote Sensing of Environment* 2013;132. Publisher: Elsevier BV:145–58.
10. Franch B, Vermote E, Roger JC, et al. A 30+ Year AVHRR Land Surface Reflectance Climate Data Record and Its Application to Wheat Yield Monitoring. *Remote Sensing* 2017;9:296.
11. João T, João G, Bruno M, and João H. Indicator-based assessment of post-fire recovery dynamics using satellite NDVI time-series. *Ecological Indicators* 2018;89. Publisher: Elsevier BV:199–212.

- 313 12. Mallegowda P, Rengaiian G, Krishnan J, and Niphadkar M. Assessing Habitat Quality of Forest-
314 Corridors through NDVI Analysis in Dry Tropical Forests of South India: Implications for
315 Conservation. *Remote Sensing* 2015;7. Publisher: MDPI AG:1619–39.
- 316 13. Marcus R, Mezzini S, Desai D, and Noonan M. Planning for a variable future: Environmental
317 stochasticity, species richness, and protected areas in Canada. 2025. DOI: 10.21203/rs.3.rs-
318 7665738/v1. URL: <https://www.researchsquare.com/article/rs-7665738/v1> (visited on
319 10/21/2025).
- 320 14. Bhatti MT, Gilani H, Ashraf M, Iqbal MS, and Munir S. Field validation of NDVI to identify
321 crop phenological signatures. *Precision Agriculture* 2024;25:2245–70.
- 322 15. Borowik T, Pettorelli N, Sönnichsen L, and Jędrzejewska B. Normalized difference vegetation
323 index (NDVI) as a predictor of forage availability for ungulates in forest and field habitats.
324 *European Journal of Wildlife Research* 2013;59:675–82.
- 325 16. Aikens EO, Kauffman MJ, Merkle JA, Dwinell SPH, Fralick GL, and Monteith KL. The
326 greenscape shapes surfing of resource waves in a large migratory herbivore. *Ecology Letters*
327 2017;20. Ed. by Nathan R. Publisher: Wiley:741–50.
- 328 17. Levins RA. Evolution in changing environments: some theoretical explorations. 3. printing.
329 *Monographs in population biology* 2. Princeton, NJ: Princeton Univ. Press, 1974. 120 pp.
- 330 18. Stephens DW and Charnov EL. Optimal foraging: Some simple stochastic models. *Behavioral*
331 *Ecology and Sociobiology* 1982;10:251–63.
- 332 19. Lande R. Risks of Population Extinction from Demographic and Environmental Stochasticity
333 and Random Catastrophes. *The American Naturalist* 1993;142. Publisher: [The University of
334 Chicago Press, The American Society of Naturalists]:911–27.
- 335 20. Sæther BE. Environmental stochasticity and population dynamics of large herbivores: a search
336 for mechanisms. *Trends in Ecology & Evolution* 1997;12. Publisher: Elsevier:143–9.
- 337 21. Mezzini S, Fleming CH, Medici EP, and Noonan MJ. How resource abundance and resource
338 stochasticity affect organisms’ range sizes. *Movement Ecology* 2025;13:20.
- 339 22. Rizzuto M, Leroux SJ, Vander Wal E, et al. Forage stoichiometry predicts the home range size
340 of a small terrestrial herbivore. *Oecologia* 2021;197:327–38.

23. Zhang X, Hu Y, Zhuang D, Qi Y, and Ma X. NDVI spatial pattern and its differentiation on the Mongolian Plateau. *Journal of Geographical Sciences* 2009;19. Publisher: Springer Science and Business Media LLC:403–15.
24. Mutti PR, Lúcio PS, Dubreuil V, and Bezerra BG. NDVI time series stochastic models for the forecast of vegetation dynamics over desertification hotspots. *International Journal of Remote Sensing* 2020;41. Publisher: Informa UK Limited:2759–88.
25. Levin SA. The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture. *Ecology* 1992;73:1943–67.
26. Fettig CJ, Runyon JB, Homicz CS, James PMA, and Ulyshen MD. Fire and Insect Interactions in North American Forests. *Current Forestry Reports* 2022;8:301–16.
27. Franklin MJ, Major RE, Bedward M, and Bradstock RA. Relative avian mobility linked to use of fire-affected resources in forested landscapes. *Forest Ecology and Management* 2021;497:119484.
28. Calhoun KL, Connor T, Gaynor KM, et al. Movement behavior in a dominant ungulate underlies successful adjustment to a rapidly changing landscape following megafire. *Movement Ecology* 2024;12:53.
29. Frankham R and Brook BW. The importance of time scale in conservation biology and ecology. *Annales Zoologici Fennici* 2004;41:459–63.
30. Simpson GL. Modelling Palaeoecological Time Series Using Generalised Additive Models. *Frontiers in Ecology and Evolution* 2018;6:149.
31. Wood SN. Generalized additive models: an introduction with R. Second edition. Chapman & Hall/CRC texts in statistical science. Boca Raton: CRC Press/Taylor & Francis Group, 2017.
32. Crameri F, Shephard GE, and Heron PJ. The misuse of colour in science communication. *Nature Communications* 2020;11:5444.
33. R Core Team. R: A Language and Environment for Statistical Computing. Vienna, Austria: R Foundation for Statistical Computing, 2024. URL: <https://www.R-project.org/>.
34. Vermote E and NOAA CDR Program. NOAA Climate Data Record (CDR) of AVHRR Normalized Difference Vegetation Index (NDVI), Version 5. 2018. DOI: 10.7289/V5ZG6QH9. URL: <https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01558/html> (visited on 04/14/2025).

- 370 35. Vermote E and NOAA CDR Program. NOAA Climate Data Record (CDR) of VIIRS Normal-
371 ized Difference Vegetation Index (NDVI), Version 1. 2022. DOI: 10.25921/GAKH-ST76. URL:
372 [https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:](https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01677/html)
373 [C01677/html](https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01677/html) (visited on 04/14/2025).
- 374 36. Hollister J, Shah T, Nowosad J, Robitaille AL, Beck MW, and Johnson M. elevatr: Access
375 Elevation Data from Various APIs. 2023. DOI: 10.5281/zenodo.8335450. URL: [https://](https://github.com/jhollist/elevatr/)
376 github.com/jhollist/elevatr/.
- 377 37. Wood SN, Goude Y, and Shaw S. Generalized additive models for large data sets. *Journal of*
378 *the Royal Statistical Society: Series C (Applied Statistics)* 2015;64:139–55.
- 379 38. Wood SN, Li Z, Shaddick G, and Augustin NH. Generalized Additive Models for Gigadata:
380 Modeling the U.K. Black Smoke Network Daily Data. *Journal of the American Statistical*
381 *Association* 2017;112:1199–210.
- 382 39. Burt WH. Territoriality and Home Range Concepts as Applied to Mammals. *Journal of Mam-*
383 *malogy* 1943;24:346.
- 384 40. Southwood TRE. Habitat, the Templet for Ecological Strategies? *The Journal of Animal Ecol-*
385 *ogy* 1977;46:336.
- 386 41. Charnov EL. Optimal foraging, the marginal value theorem. *Theoretical Population Biology*
387 1976;9:129–36.
- 388 42. Broekman MJE, Hilbers JP, Hoeks S, Huijbregts MAJ, Schipper AM, and Tucker MA. Envi-
389 ronmental drivers of global variation in home range size of terrestrial and marine mammals.
390 *Journal of Animal Ecology* 2024;93:488–500.
- 391 43. Marcus R and Noonan MJ. Changing variability is an overlooked aspect of protected area
392 planning. 2023. DOI: 10.1101/2023.10.26.564062. URL: [http://biorxiv.org/lookup/doi/](http://biorxiv.org/lookup/doi/10.1101/2023.10.26.564062)
393 [10.1101/2023.10.26.564062](http://biorxiv.org/lookup/doi/10.1101/2023.10.26.564062) (visited on 10/23/2025).
- 394 44. Nilsen EB, Herfindal I, and Linnell JDC. Can intra-specific variation in carnivore home-range
395 size be explained using remote-sensing estimates of environmental productivity? *Écoscience*
396 2005;12:68–75.
- 397 45. Middleton AD, Merkle JA, McWhirter DE, et al. Green-wave surfing increases fat gain in a
398 migratory ungulate. *Oikos* 2018;127:1060–8.

- 399 46. Geremia C, Merkle JA, Eacker DR, et al. Migrating bison engineer the green wave. *Proceedings*
400 *of the National Academy of Sciences* 2019;116:25707–13.
- 401 47. Aikens EO, Merkle JA, Xu W, and Sawyer H. Pronghorn movements and mortality during ex-
402 tremite weather highlight the critical importance of connectivity. *Current Biology* 2025;35:1927–
403 1934.e2.
- 404 48. Steixner-Kumar S and Gläscher J. Strategies for navigating a dynamic world. *Science* 2020;369:1056–
405 7.
- 406 49. Rickbeil GJM, Merkle JA, Anderson G, et al. Plasticity in elk migration timing is a response
407 to changing environmental conditions. *Global Change Biology* 2019;25:2368–81.
- 408 50. Polansky L, Kilian W, and Wittemyer G. Elucidating the significance of spatial memory on
409 movement decisions by African savannah elephants using state–space models. *Proceedings of*
410 *the Royal Society B: Biological Sciences* 2015;282:20143042.
- 411 51. Abrahms B, Hazen EL, Aikens EO, et al. Memory and resource tracking drive blue whale
412 migrations. *Proceedings of the National Academy of Sciences* 2019;116:5582–7.
- 413 52. Riotte-Lambert L and Matthiopoulos J. Environmental Predictability as a Cause and Conse-
414 quence of Animal Movement. *Trends in Ecology & Evolution* 2020;35:163–74.
- 415 53. Falcón-Cortés A, Boyer D, Merrill E, Frair JL, and Morales JM. Hierarchical, Memory-Based
416 Movement Models for Translocated Elk (*Cervus canadensis*). *Frontiers in Ecology and Evolu-*
417 *tion* 2021;9:702925.
- 418 54. Ranc N, Cagnacci F, and Moorcroft PR. Memory drives the formation of animal home ranges:
419 Evidence from a reintroduction. *Ecology Letters* 2022;25. Ed. by Coulson T:716–28.
- 420 55. Botero CA, Weissing FJ, Wright J, and Rubenstein DR. Evolutionary tipping points in the
421 capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences*
422 2015;112:184–9.
- 423 56. Polazzo F, Limberger R, Pennekamp F, Ross SRPJ, Simpson GL, and Petchey OL. Measuring
424 the Response Diversity of Ecological Communities Experiencing Multifarious Environmental
425 Change. *Global Change Biology* 2024;30:e17594.
- 426 57. Anderson SC, Branch TA, Cooper AB, and Dulvy NK. Black-swan events in animal popula-
427 tions. *Proceedings of the National Academy of Sciences* 2017;114:3252–7.

- 428 58. Logares R and Nuñez M. Black Swans in ecology and evolution: The importance of improbable
429 but highly influential events. *Ideas in Ecology and Evolution* 2012.
- 430 59. Sawyer H, Merkle JA, Middleton AD, Dwinnell SPH, and Monteith KL. Migratory plasticity is
431 not ubiquitous among large herbivores. *Journal of Animal Ecology* 2019;88. Ed. by Chapman
432 J:450–60.
- 433 60. Bjorndahl JA, Gushulak CA, Mezzini S, et al. Abrupt changes in the physical and biological
434 structure of endorheic upland lakes due to 8-m lake-level variation during the 20th century.
435 *Limnology and Oceanography* 2022;67:1022–39.
- 436 61. Chevin LM, Lande R, and Mace GM. Adaptation, Plasticity, and Extinction in a Changing En-
437 vironment: Towards a Predictive Theory. *PLoS Biology* 2010;8. Ed. by Kingsolver JG:e1000357.
- 438 62. Wolf AA, Zavaleta ES, and Selmants PC. Flowering phenology shifts in response to biodiversity
439 loss. *Proceedings of the National Academy of Sciences* 2017;114:3463–8.
- 440 63. Wolkovich EM, Cook BI, Allen JM, et al. Warming experiments underpredict plant phenological
441 responses to climate change. *Nature* 2012;485:494–7.
- 442 64. Stefanescu C, Ubach A, and Wiklund C. Timing of mating, reproductive status and resource
443 availability in relation to migration in the painted lady butterfly. *Animal Behaviour* 2021;172:145–
444 53.