

DENVar: A Global Dynamic Estimate of NDVI Variance

Stefano Mezzini^{1,2}, Gavin L Simpson³, and Michael J Noonan^{1,2,4,*}

¹Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

²Department of Biology, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

³Department of Animal and Veterinary Sciences, Aarhus University, Denmark.

⁴Department of Computer Science, Math, Physics, and Statistics, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

*Address correspondence to: michael.noonan@ubc.ca

Abstract

Remote-sensing proxies of habitat productivity such as the Normalized Difference Vegetation Index (NDVI) have helped develop and support many hypotheses and projects. However, historically, users tended to focus solely on trends in mean productivity. In recent years, the variance around the mean (i.e., stochasticity) has gained more attention, particularly as climate change and human-induced rapid environmental change transform the ecosystems species have evolved in and adapted to. Still, calculating trends in NDVI stochasticity requires careful modeling, since stochasticity estimates are conditional on the estimated mean and the scale at which the data are collected and modeled. We thus present a new global Dynamic Estimate of NDVI Variance, *DENVar*. We estimate spatiotemporal trends in NDVI mean and variance using Generalized Additive Models, which provide a flexible yet transparent and interpretable model structure. We show that *DENVar* varies nonlinearly with **XXX** and can be used to test many hypotheses related to environmental stochasticity, such as optimal foraging theory and regime shifts as well as for conservation. We provide a publicly and freely available repository for downloading rasters of daily and long-term averages of *DENVar* and the related estimates of mean NDVI.

1 Introduction

The Normalized Difference Vegetation Index (NDVI) is a proxy of primary productivity used in a wide range of disciplines, including ecology (Pettorelli et al., 2005, 2011), landscape phenology (Merkle et al., 2016; Hmimina et al., 2013), agriculture (Franch et al., 2017), forestry (João et al., 2018), and conservation (Mallegowda et al., 2015; Marcus et al., 2025). High-resolution, remotely-sensed NDVI data have allowed analysts to estimate plant biomass without fieldwork (Bhatti et al., 2024; Borowik et al., 2013) and leverage such estimates for studying large-scale plant dynamics (Pettorelli et al., 2005) and their effects on animal movement behavior (Merkle et al., 2016; Aikens et al., 2017). Still, most NDVI work is limited to quantifying changes in and responses to mean NDVI while ignoring changes in the variance around the mean.

Environmental stochasticity has been studied for decades (Levins, 1974; Stephens and Charnov, 1982; Lande, 1993; Sæther, 1997), but recent work has drawn more attention to the effects environmental stochasticity has at a wide range of scales, from individual organisms (Mezzini et al., 2025; Rizzuto et al., 2021) to landscapes (Zhang et al., 2009; Mutti et al., 2020; Marcus et al., 2025). However, estimating trends in NDVI variance requires more nuanced modeling than estimating changes in mean NDVI, particularly when environments are temporally dynamic and spatially heterogeneous. The scale at which processes are observed determine how individuals’ perception of and ability to respond to change (Levin, 1992). A forest fire is severely disruptive and deadly for many insects (Fettig et al., 2022), while birds and large mammal may not struggle to escape the fire and or establish a new range (Franklin et al., 2021; Calhoun et al., 2024). Similarly, the effects of the fire recovery will depend on its duration relative to other organisms’ phenology and lifespan (Frankham and Brook, 2004). Therefore, distinguishing between a change in mean NDVI and a stochastic fluctuation requires a conscious decision about the scale at which to consider the data.

Previous work suggests that environmental stochasticity can reduce limit species’ ability to adapt or specialize (Chevin et al., 2010), alter species’ movement behavior (Mezzini et al., 2025), and even affect extinction risk at both the population and species levels (Lande, 1993; Bull et al., 2007; Fung et al., 2018). Still, the effects of environmental stochasticity on species dynamics and ecosystem stability remain understudied (Marcus and Noonan, 2023). Global, temporally dynamic estimates of NDVI variance would facilitate research by building upon the widespread use and interpretability of NDVI while also removing the barriers of computational demand and statistical knowledge that are required for quantifying environmental stochasticity. However, since NDVI is not a direct measurement primary productivity but a proxy, it should be interpreted carefully (Pettorelli et al., 2011). NDVI is prone to sensitivity loss in very green environments (Matsushita et al., 2007; Son et al., 2014), values vary across sensors (Beck et al., 2011; Fan and Liu, 2016; Huang et al., 2021), and the coarse spatiotemporal scale of satellite-derived NDVI can be a substantial barrier when assessing fine-scale trends (Goodchild and Quattrochi, 2023; Reid et al., 2018). However, careful statistical modeling can help overcome some of these limitations. In particular, Generalized Additive Models (GAMs) are particularly convenient due to their ability to learn flexibly from data (Wood, 2017; Simpson, 2018) and handle very large spatiotemporal datasets (Wood et al., 2015, 2017).

We present a new measure of environmental stochasticity, which we name the Dynamic Estimate

of NDVI Variance (DENVar). We calculate DENVar using 44 years of NDVI data from the AVHRR and VIIRS collection of sensors and GAMs. Although satellites with AVHRR and VIIRS sensors provide data at a coarser spatial resolution (0.05°) than other satellites (e.g., 250m for MODIS), they are the only sensors to provide data at a daily frequency and for more than 40 years. Our use of GAMs provides a flexible yet interpretable approach for estimating trends in mean and variance in NDVI without the rigidity of other more traditional methods (e.g., ARIMA) or the opaqueness of machine learning models such as neural networks (Simpson, 2018; Wood, 2017). We illustrate the relationships between DENVar and multiple, relevant ecological metrics, and we provide access to a free repository of daily, global estimates of DENVar and the corresponding estimates of mean NDVI.

2 Methods

2.1 Choice of color palettes

We represent NDVI using a modified version of Fabio Crameri’s divergent *bukavu* palette (Crameri, 2018a; Crameri et al., 2020), which has high-contrast for deuteranope and protanope vision. For NDVI values between -0.1 and 1, the colors also have sufficient contrast for the colors to also be distinguishable by strong tritanope and achromatic vision. Fig. A.1 contains an approximate representation of how each vision type would perceive each color palettes used. We obtained all color palettes with three or more colors from the *khroma* package (v. 1.14.0 Frerebeau, 2024) for R (v. 4.4.0; R Core Team, 2024).

2.2 Datasets used

Fig. 1 illustrates the processes of data collection, cleaning, and modeling used in this project. We downloaded NDVI image composites directly from the servers for data from AVHRR sensors (1981-06-24 to 2013-12-31; United States National Centers for Environmental Information (NCEI) and United States National Aeronautics and Space Administration, 2025; Vermote and NOAA CDR Program, 2018) and VIIRS sensors (2014-01-01 to 2025-06-29; United States National Aeronautics and Space Administration, 2025; Vermote and NOAA CDR Program, 2022).

We removed all pixels labeled as "cloudy" for AVHRR rasters and "probably cloudy" or "confidently cloudy" for VIIRS rasters. To remove pixels that included large portions of water, we used a shapefile of major water bodies (Garmin International, Inc. and Esri, 2025) to produce a static raster of the percentage of water within each pixel, at the same spatial resolution as the NDVI rasters (0.05° ; Fig. A.2). We removed all pixels with $> 40\%$ water cover.

Many rasters had horizontal bands of $\text{NDVI} > 0.9$ at unrealistically northern latitudes between September and March, likely due to limited visibility during fall and winter in the Northern hemisphere. We identified and removed problematic pixels using the monthly empirical cumulative distribution functions of NDVI across latitude (Fig. A.3). We did not filter problematic values at latitudes $< 60^\circ\text{S}$ since we fully excluded Antarctica from the analyses (Fig. A.4).




Figure 1: **Flowchart for the analyses used in the manuscript.** The curly braces on the right indicate the subsections in the Methods section that detail the dataset collation and cleaning (2.2), and statistical analyses (2.3).

To ensure the number of rows for each dataset was below the maximum data frame size in R ($3.17 \times 10^{10} > 2^{31} - 1 \approx 0.21 \times 10^{10}$), we reduced the dataset size by splitting it across four spatial groups (Figs. A.4 and A.5) and spatially and temporally aggregated the rasters by a factor of four. The final dataset thus had one raster every four days with a spatial resolution of $0.05^\circ \times 4 = 0.2^\circ$ (Fig. A.6). Finally, we annotated the dataset with columns of day of year (1-366), integer year (1981-2025), percent water cover, and elevation above sea level (m). The digital elevation model was obtained using the `get_elev_raster()` function from the `elevatr` package for R (v. 0.99.0; Hollister et al., 2023) with an original resolution of 0.076° (Fig. A.6). Despite the substantial reduction in dataset size, the dataset creation was still challenging for the machine used, which had 2.2 TB of RAM and two Intel Xeon Platinum 8462Y+ processors (64 cores total).

2.3 Statistical Analyses

We estimated spatiotemporal trends in mean and variance in NDVI using Generalized Additive Models (GAMs) via the `mgcv` package for R (v. 1.9-3 Wood, 2017). To reduce modeling fitting times

with negligible losses to model accuracy, we used the `bam()` function with fast REstricted Marginal Likelihood (`method = fREML`) and covariate discretization (`discrete = TRUE`; Wood et al., 2015, 2017). Ideally, NDVI should be modeled using beta location-scale models (after the linear transformation $Y^* = (Y + 1)/2$) to account for the fact that: (1) NDVI is bounded between -1 and 1, and (2) the variance in NDVI is dependent on the mean (and vice-versa), since ecosystems with very low NDVI (e.g., rock, ice, or concrete) or very high NDVI (e.g., dense forest) tend to have lower variance in NDVI. However, preliminary tests and work by (Marcus et al., 2025) showed that beta GAMs and particularly beta location-scale GAMs had prohibitive requirements for both fitting times and memory requirements, since both requirements scale $O(Mnp^2)$, for M smoothing parameters and a model matrix with n rows and p columns (Wood et al., 2015). A beta location-scale model for only the island of Sardinia (24,000 km²; Appendix B) was interrupted after 144 hours, during which the model had only completed three steps using the Newton method described by Wood and Fasiolo (2017). In contrast, Gaussian models with fixed scaled parameters (i.e., constant variance) fit substantially faster and provided very similar spatiotemporal estimates of mean and variance in NDVI (Appendix B). Thus, we estimated trends in mean and variance in NDVI using a pair of GAMs for each region shown in Fig. A.4; one for estimating mean NDVI over space and time (assuming a constant variance), and one for estimating the mean squared residual (i.e., variance) in NDVI (again, assuming a constant scale parameter).

Together, the sets of models for mean NDVI (i.e., μ) and variance in NDVI (i.e., σ^2) had the following structure, with ν indicating NDVI:

$$\left\{ \begin{array}{l} \nu \sim \text{Normal}(\mu, \sigma^2) \\ \mu = \mathbb{E}(\nu) \approx \hat{\mathbb{E}}(\nu) = \hat{\mu} \\ \hat{\mu} = \beta_{\mu} + s(\text{prop_water}) + s(\text{elev_m}) + s(\text{doy}) + s(\text{year}) + s(\text{lat}, \text{long}) + \\ \quad t(\text{doy}, \text{year}) + t(\text{doy}, \text{lat}, \text{long}) + t(\text{year}, \text{lat}, \text{long}) \\ \sigma^2 = \mathbb{E}((\nu - \mu)^2) \approx \hat{\mathbb{E}}((\nu - \hat{\mu})^2) = \hat{\sigma}^2 \\ \hat{\sigma}^2 = \beta_{\sigma^2} + s(\text{prop_water}) + s(\text{elev_m}) + s(\text{doy}) + s(\text{year}) + s(\text{lat}, \text{long}) + \\ \quad t(\text{doy}, \text{year}) + t(\text{doy}, \text{lat}, \text{long}) + t(\text{year}, \text{lat}, \text{long}) + s(\hat{\mu}) \end{array} \right. , \quad (1)$$

where, for a given (spatiotemporally aggregated) pixel, `prop_water` is the proportion of water within the pixel, `elev_m` is elevation above sea level in meters, `doy` is integer day of year (1-366), `year` is integer year (1981-2025), `lat` and `long` are latitude and longitude, respectively. For ease of readability, we omit the pixel-level indices. $s(\cdot)$ indicates marginal smooth functions, while $t(\cdot)$ indicates tensor product interaction terms. Seasonal terms used cyclical cubic splines (`bs = 'cc'`) with edge knots 0.5 and 366.5, while spatial smooths used a basis with second-order spline on the sphere (`bs = 'sos'`; Wood, 2017). The linear predictors for μ and σ^2 have the same structure, other than the $s(\mu)$ smooth term for σ^2 , but the functions for σ^2 are independent of those for μ . Despite the data aggregation procedures and faster fitting methods, including parallelizing computation across

145 60 cores, each of the four models required multiple days to fit. Predicting from the models for $\hat{\mu}$ was
 146 also had substantial time and memory demands, since predictions were calculated without discrete
 147 prediction methods (`discrete = FALSE`). While discrete predictions were orders of magnitude faster,
 148 they were excessively spatially coarse. See Appendix C and the code availability statement for
 149 additional information.

150 Since there are no corrections to any bias in residuals from models 1, the estimates of variance
 151 from models 1 may be inflated near areas or periods of rapid and substantial change. Still, the
 152 appreciability of this inflation is dependent on the spatiotemporal scale over which changes are
 153 perceived (Levin, 1992; Frankham and Brook, 2004; Goodchild and Quattrochi, 2023; Mezzini et al.,
 154 2025).

155 From the two sets of models, we produced daily estimates of (1) the Dynamic Estimate of NDVI
 156 Variance (DENVar) and (2) the estimated mean NDVI they are conditional on, at a **XXX**°spatial
 157 scale. The rasters are freely available on GitHub (see code availability statement), along with
 158 long-term (1981-2025) averages of both DENVar and the estimated mean NDVI. Since estimates of
 159 variance are conditional on the corresponding estimates of the mean, we suggest users always use
 160 the corresponding rasters of mean NDVI in their analyses rather than using independent estimates
 161 of mean NDVI.

162 3 Results

163 Other variables to potentially add to hexplots:

- 164 • temperature
- 165 • seasonal temperature range
- 166 • precip
- 167 • seasonal temperature range
- 168 • spp diversity
- 169 • gross primary productivity
- 170 • max body size
- 171 • <https://silvis.forest.wisc.edu/globalwui/>
- 172 • <https://silvis.forest.wisc.edu/data/dhis/>: NDVI16 (Normalized Difference Vegetation Index),
 173 EVI16 (Enhanced Vegetation Index), FPAR8 (Fraction absorbed Photosynthetically Active
 174 Radiation), LAI8 (Leaf Area Index), GPP8 (Gross Primary Productivity)
- 175 • extreme weather event
- 176 • invasive species
- 177 • disease occurrence?

3.1 Obtaining the data

- Data should be available as daily rasters on a github repo?
- make an app for data annotation?

4 Discussion

We have presented a new spatiotemporally dynamic metric of environmental stochasticity, which we have called the Dynamic Estimate of NDVI Variance, DENVar. In this section, we discuss the interpretation of DENVar in greater depth, and we argue the advantages and limitations of the statistical methods used. We conclude by suggesting some use cases of DENVar within animal movement ecology, landscape ecology, and phenology, but we invite readers to consider how DENVar may be of use in their field and adjacent disciplines.

4.1 Interpreting DENVar: the importance of scale

difficulties in estimating variance

what counts as a change? predictable vs stochastic changes

smoothness of mean impacts smoothness and size of variance: depends on scale of interest (e.g., animals' ability to respond to, learn, and predict changes in the mean). Changes not attributed to trends in the mean are stochastic

DENVar is a good metric as long as the spatiotemporal scale of interest is similar or larger to that of DENVar. May not work well for small-scale processes such as the movement dynamics of a small rodent, the phenology of a small orchard when spatiotemporal heterogeneity is finer than DENVar's resolution

4.2 Practical applications

Convert the subsections to a single subsection:

- spatial ecology
- movement behavior: (Mezzini et al., 2025)
- conservation (Marcus et al., 2025), etc.
- anthropology (navigating paper)
- industry (agriculture)

Environmental stochasticity and animal movement ecology

The field of animal movement ecology has long shown that animals alter their movement in response to resource abundance or availability (Burt, 1943; Southwood, 1977; Charnov, 1976; Broekman et al., 2024). More recently, movement ecologists have started focusing more on resource stochasticity

(Stephens and Charnov, 1982; Rizzuto et al., 2021; Marcus and Noonan, 2023), but it is worth distinguishing between resource stochasticity and seasonality (Nilsen et al., 2005). Seasonality is generally best interpreted as a cyclical change in the mean that many animals can learn to predict, as in the case of green-wave surfing (Middleton et al., 2018; Geremia et al., 2019). In contrast, environmental stochasticity refers to unpredictable changes, such as forest fires [ref](#), sudden snow storms (Aikens et al., 2025), and unusual phenology [ref](#). Animals also respond to stochastic changes (Mezzini et al., 2025), but their ability to adapt depends on behavioral plasticity (Steixner-Kumar and Gläser, 2020; Rickbeil et al., 2019), and memory (Polansky et al., 2015; Abrahms et al., 2019; Riotte-Lambert and Matthiopoulos, 2020; Falcón-Cortés et al., 2021; Ranc et al., 2022).

evolutionary tipping points (Botero et al., 2015)

multifarious change (Polazzo et al., 2024)

extreme events, black swans (Anderson et al., 2017; Logares and Nuñez, 2012)

- migration routes are key habitat (Ortega et al., in prep), especially in stochastic environments

- can fail to adapt to changes in conditions (Sawyer et al., 2019)

- <https://www.sciencedirect.com/science/article/pii/S0960982220308484>

- <https://onlinelibrary.wiley.com/doi/full/10.1111/gcb.15169>

- industrial development affects animals' ability to respond to both predictable and unpredictable change (<https://www.nature.com/articles/s41559-022-01887-9>)

- effects of sudden deep snow in red desert on proghorn (<https://www.sciencedirect.com/science/article/abs/pii/S0960982220308484>)

- <https://www.nature.com/articles/s41467-023-37750-z>

- <https://esajournals.onlinelibrary.wiley.com/doi/full/10.1002/ecy.4238>

- detecting regime shifts (in lakes: (Bjorndahl et al., 2022))

Environmental stochasticity and population dynamics

read lande_risks_1993

Environmental stochasticity reduces habitats' energetic balance and carrying capacity (Chevin et al., 2010) as individuals struggle to rely on unpredictable resources and increase the instability of populations and communities (Lande, 1993). Additionally, stochastic environments reduce the ability of specialist species to adapt and evolve while selecting for more generalist species (Levins, 1974). Consequently, stochastic environments tend to have lower species diversity (Marcus et al., 2025).

Recent work by Devries has suggested that NDVI stochasticity is an important metric for evaluating forest recovery after large fires, while Collinson has suggested that variance in NDVI is an important predictor of the occurrence of lightning-induced forest fires.

- jonzen_rapid_2006

- 242 • effects can be compounded by cumulative effects (e.g., loss of species diversity; (Wolf et al.,
243 2017))
- 244 • climate change affects plant phenology, but changes in situ may not match experimental esti-
245 mates (Wolkovich et al., 2012)
- 246 • reproduction of painted lady butterfly (Stefanescu et al., 2021)
- 247 • wessling_seasonal_2018




Figure 2: Estimated mean NDVI (A) and DENVar (B) for 2003-07-02. **CHANGE TO LONG-TERM MEAN?**




Figure 3: Surface plots of mean NDVI (A) and DENVar (B) over time and latitudes.




Figure 4: Hex plots of the relationship between the long-term global Dynamic Estimate of NDVI Variance (DENVar) and common ecological metrics. The fill of the hexagonal cells represents the log-10 number of points within the cell. Rowwise, the variables are: long-term mean NDVI (as estimated in this manuscript), the machine-learning Human Footprint Index (ml-HFI) of , and the 2015 cumulative, minimum, and seasonal range in the Dynamic Habitat Index of (Radeloff et al., 2019).




Figure 5: Map of terrestrial biomes along with hex plots of mean NDVI and DENVar for each biome. **UPDATE FIGURE TO HEXPLOTS.**

5 Other useful references

- **READ**: Methods for Detecting Early Warnings of Critical Transitions in Time Series Illustrated Using Simulated Ecological Data (<https://doi.org/10.1371/journal.pone.0041010>)
- **READ**: <https://doi.org/10.1029/2024GL109298>
- **READ**: Site fidelity as a maladaptive behavior in the Anthropocene: <https://esajournals.onlinelibrary.wiley.com/doi/10.1002/ecs.25888>
- **READ**: Pettorelli N., Vik J.O., Mysterud A., Gaillard J.-M., Tucker C.J. & Stenseth N. Chr. (2005). Using the satellite-derived NDVI to assess ecological responses to environmental change. *Trends in Ecology & Evolution* 20, 503–510. <https://doi.org/10.1016/j.tree.2005.05.011>
- Ecological and evolutionary consequences of changing seasonality: <https://www.science.org/doi/10.1126/science.adc1234>
- increase in variance before eutrophication in lakes: <https://doi.org/10.1002/lno.10355>
- uses ndvi variance: <https://esajournals.onlinelibrary.wiley.com/doi/full/10.1002/ecm.1348>
- bro-jorgensen_using_2008: in the dry season preceding the rut topi antelope (*Damaliscus lunatus*) respond to forage availability at the large scale but not the fine scale
- keith_predicting_2008
- chevin_adaptation_2010
- rickbeil_plasticity_2019
- mueller_search_2008
- pettorelli_using_2005
- nilsen_can_2005
- merkle_large_2016
- tian_evaluating_2015
- huang_commentary_2021
- fan_global_2016
- wang_stochastic_2019
- pease_ecological_2024
- xu_plasticity_2021
- calibrating across sensors: <https://ieeexplore.ieee.org/abstract/document/6809983>

275 **Acknowledgments**

276 The authors would like to thank Dr. Simon Wood for writing the custom Beta location-scale family
277 even though it was not used in the final models.

278 **Author Contributions**

279 **Funding**

280 **Conflicts of Interest**

281 The authors declare that there is no conflict of interest regarding the publication of this article.

282 **Code and Data Availability**

283 All code is available on GitHub at <https://github.com/QuantitativeEcologyLab/ndvi-stochasticity>,
284 including code for downloading the original NDVI rasters directly from the servers they are housed
285 in. Most data are not included due to GitHub's size limitations, but all objects can be created using
286 the code provided.

287 **Supplementary Materials**

- 288 • Appendix A: Color palettes used and input data.
- 289 • Appendix B: Sensitivity analyses.
- 290 • Appendix C: Code for fitting the models, along with supporting supplementary figures.

References

- Briana Abrahms, Elliott L. Hazen, Ellen O. Aikens, Matthew S. Savoca, Jeremy A. Goldbogen, Steven J. Bograd, Michael G. Jacox, Ladd M. Irvine, Daniel M. Palacios, and Bruce R. Mate. Memory and resource tracking drive blue whale migrations. *Proceedings of the National Academy of Sciences*, 116(12):5582–5587, March 2019. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.1819031116. URL <https://pnas.org/doi/full/10.1073/pnas.1819031116>.
- Ellen O. Aikens, Matthew J. Kauffman, Jerod A. Merkle, Samantha P. H. Dwinnell, Gary L. Fralick, and Kevin L. Monteith. The greenscape shapes surfing of resource waves in a large migratory herbivore. *Ecology Letters*, 20(6):741–750, June 2017. ISSN 1461-023X, 1461-0248. doi: 10.1111/ele.12772. URL <https://onlinelibrary.wiley.com/doi/10.1111/ele.12772>. Publisher: Wiley.
- Ellen O. Aikens, Jerod A. Merkle, Wenjing Xu, and Hall Sawyer. Pronghorn movements and mortality during extreme weather highlight the critical importance of connectivity. *Current Biology*, 35(8):1927–1934.e2, April 2025. ISSN 09609822. doi: 10.1016/j.cub.2025.03.010. URL <https://linkinghub.elsevier.com/retrieve/pii/S0960982225002957>.
- Sean C. Anderson, Trevor A. Branch, Andrew B. Cooper, and Nicholas K. Dulvy. Black-swan events in animal populations. *Proceedings of the National Academy of Sciences*, 114(12):3252–3257, March 2017. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.1611525114. URL <http://www.pnas.org/lookup/doi/10.1073/pnas.1611525114>.
- Hylke E. Beck, Tim R. McVicar, Albert I.J.M. Van Dijk, Jaap Schellekens, Richard A.M. De Jeu, and L. Adrian Bruijnzeel. Global evaluation of four AVHRR–NDVI data sets: Intercomparison and assessment against Landsat imagery. *Remote Sensing of Environment*, 115(10):2547–2563, October 2011. ISSN 0034-4257. doi: 10.1016/j.rse.2011.05.012. URL <https://linkinghub.elsevier.com/retrieve/pii/S0034425711001957>. Publisher: Elsevier BV.
- Muhammad Tousif Bhatti, Hammad Gilani, Muhammad Ashraf, Muhammad Shahid Iqbal, and Sarfraz Munir. Field validation of NDVI to identify crop phenological signatures. *Precision Agriculture*, 25(5):2245–2270, October 2024. ISSN 1385-2256, 1573-1618. doi: 10.1007/s11119-024-10165-6. URL <https://link.springer.com/10.1007/s11119-024-10165-6>.
- Judith A. Bjorndahl, Cale A.C. Gushulak, Stefano Mezzini, Gavin L. Simpson, Heather A. Haig, Peter R. Leavitt, and Kerri Finlay. Abrupt changes in the physical and biological structure of endorheic upland lakes due to 8-m lake-level variation during the 20th century. *Limnology and Oceanography*, 67(5):1022–1039, May 2022. ISSN 0024-3590, 1939-5590. doi: 10.1002/lno.12054. URL <https://aslopubs.onlinelibrary.wiley.com/doi/10.1002/lno.12054>.
- Tomasz Borowik, Nathalie Pettorelli, Leif Sönnichsen, and Bogumiła Jędrzejewska. Normalized difference vegetation index (NDVI) as a predictor of forage availability for ungulates in forest and field habitats. *European Journal of Wildlife Research*, 59(5):675–682, October 2013. ISSN

1612-4642, 1439-0574. doi: 10.1007/s10344-013-0720-0. URL <http://link.springer.com/10.1007/s10344-013-0720-0>.

Carlos A. Botero, Franz J. Weissing, Jonathan Wright, and Dustin R. Rubenstein. Evolutionary tipping points in the capacity to adapt to environmental change. *Proceedings of the National Academy of Sciences*, 112(1):184–189, January 2015. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.1408589111. URL <https://pnas.org/doi/full/10.1073/pnas.1408589111>.

Maarten J. E. Broekman, Jelle P. Hilbers, Selwyn Hoeks, Mark A. J. Huijbregts, Aafke M. Schipper, and Marlee A. Tucker. Environmental drivers of global variation in home range size of terrestrial and marine mammals. *Journal of Animal Ecology*, 93(4):488–500, April 2024. ISSN 0021-8790, 1365-2656. doi: 10.1111/1365-2656.14073. URL <https://besjournals.onlinelibrary.wiley.com/doi/10.1111/1365-2656.14073>.

James C Bull, Nicola J Pickup, Brian Pickett, Michael P Hassell, and Michael B Bonsall. Metapopulation extinction risk is increased by environmental stochasticity and assemblage complexity. *Proceedings of the Royal Society B: Biological Sciences*, 274(1606):87–96, January 2007. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2006.3691. URL <https://royalsocietypublishing.org/doi/10.1098/rspb.2006.3691>.

William Henry Burt. Territoriality and Home Range Concepts as Applied to Mammals. *Journal of Mammalogy*, 24(3):346, August 1943. ISSN 00222372. doi: 10.2307/1374834. URL <https://academic.oup.com/jmammal/article-lookup/doi/10.2307/1374834>.

Kendall L. Calhoun, Thomas Connor, Kaitlyn M. Gaynor, Amy Van Scoyoc, Alex McInturff, Samantha E.S. Kreling, and Justin S. Brashares. Movement behavior in a dominant ungulate underlies successful adjustment to a rapidly changing landscape following megafire. *Movement Ecology*, 12(1):53, July 2024. ISSN 2051-3933. doi: 10.1186/s40462-024-00488-4. URL <https://movementecologyjournal.biomedcentral.com/articles/10.1186/s40462-024-00488-4>.

Eric L. Charnov. Optimal foraging, the marginal value theorem. *Theoretical Population Biology*, 9(2):129–136, April 1976. ISSN 00405809. doi: 10.1016/0040-5809(76)90040-X. URL <https://linkinghub.elsevier.com/retrieve/pii/004058097690040X>.

Luis-Miguel Chevin, Russell Lande, and Georgina M. Mace. Adaptation, Plasticity, and Extinction in a Changing Environment: Towards a Predictive Theory. *PLoS Biology*, 8(4):e1000357, April 2010. ISSN 1545-7885. doi: 10.1371/journal.pbio.1000357. URL <https://dx.plos.org/10.1371/journal.pbio.1000357>.

Fabio Cramer. Geodynamic diagnostics, scientific visualisation and StagLab 3.0. *Geoscientific Model Development*, 11(6):2541–2562, June 2018a. ISSN 1991-9603. doi: 10.5194/gmd-11-2541-2018. URL <https://gmd.copernicus.org/articles/11/2541/2018/>.

Fabio Cramer. Scientific colour-maps, June 2018b. URL <https://zenodo.org/record/1287763>.
Language: en.

363 Fabio Crameri, Grace E. Shephard, and Philip J. Heron. The misuse of colour in science
 364 communication. *Nature Communications*, 11(1):5444, October 2020. ISSN 2041-1723. doi:
 365 10.1038/s41467-020-19160-7. URL <https://www.nature.com/articles/s41467-020-19160-7>.

366 Andrea Falcón-Cortés, Denis Boyer, Evelyn Merrill, Jacqueline L. Frair, and Juan Manuel Morales.
 367 Hierarchical, Memory-Based Movement Models for Translocated Elk (*Cervus canadensis*). *Frontiers in Ecology and Evolution*, 9:702925, August 2021. ISSN 2296-701X. doi: 10.3389/fevo.2021.
 368 702925. URL <https://www.frontiersin.org/articles/10.3389/fevo.2021.702925/full>.

370 Xingwang Fan and Yuanbo Liu. A global study of NDVI difference among moderate-resolution
 371 satellite sensors. *ISPRS Journal of Photogrammetry and Remote Sensing*, 121:177–191, November
 372 2016. ISSN 09242716. doi: 10.1016/j.isprsjprs.2016.09.008. URL <https://linkinghub.elsevier.com/retrieve/pii/S0924271616303975>.

374 Christopher J. Fettig, Justin B. Runyon, Crystal S. Homicz, Patrick M. A. James, and Michael D.
 375 Ulyshen. Fire and Insect Interactions in North American Forests. *Current Forestry Reports*,
 376 8(4):301–316, August 2022. ISSN 2198-6436. doi: 10.1007/s40725-022-00170-1. URL <https://link.springer.com/10.1007/s40725-022-00170-1>.

378 Belen Franch, Eric Vermote, Jean-Claude Roger, Emilie Murphy, Inbal Becker-Reshef, Chris Justice,
 379 Martin Claverie, Jyoteshwar Nagol, Ivan Csizar, Dave Meyer, Frederic Baret, Edward Masuoka,
 380 Robert Wolfe, and Sadashiva Devadiga. A 30+ Year AVHRR Land Surface Reflectance Climate
 381 Data Record and Its Application to Wheat Yield Monitoring. *Remote Sensing*, 9(3):296, March
 382 2017. ISSN 2072-4292. doi: 10.3390/rs9030296. URL <https://www.mdpi.com/2072-4292/9/3/296>.

384 Richard Frankham and Barry W. Brook. The importance of time scale in conservation biology and
 385 ecology. *Annales Zoologici Fennici*, 41:459–463, 2004. ISSN 0003-455X.

386 Michael J.M. Franklin, Richard E. Major, Michael Bedward, and Ross A. Bradstock. Relative
 387 avian mobility linked to use of fire-affected resources in forested landscapes. *Forest Ecology and*
 388 *Management*, 497:119484, October 2021. ISSN 03781127. doi: 10.1016/j.foreco.2021.119484. URL
 389 <https://linkinghub.elsevier.com/retrieve/pii/S0378112721005739>.

390 Nicolas Frerebeau. *khroma: Colour Schemes for Scientific Data Visualization*. Université Bor-
 391 deaux Montaigne, Pessac, France, 2024. doi: 10.5281/zenodo.1472077. URL <https://packages.tesselle.org/khroma/>.

393 Tak Fung, James P. O'Dwyer, and Ryan A. Chisholm. Quantifying species extinction risk under
 394 temporal environmental variance. *Ecological Complexity*, 34:139–146, May 2018. ISSN 1476945X.
 395 doi: 10.1016/j.ecocom.2017.09.006. URL <https://linkinghub.elsevier.com/retrieve/pii/S1476945X1730082X>.

397 Garmin International, Inc. and Esri. World Water Bodies, April 2025. URL <https://hub.arcgis.com/content/e750071279bf450cbd510454a80f2e63/about>.

- Chris Geremia, Jerod A. Merkle, Daniel R. Eacker, Rick L. Wallen, P. J. White, Mark Hebblewhite, and Matthew J. Kauffman. Migrating bison engineer the green wave. *Proceedings of the National Academy of Sciences*, 116(51):25707–25713, December 2019. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.1913783116. URL <http://www.pnas.org/lookup/doi/10.1073/pnas.1913783116>.
- Michael F. Goodchild and Dale A. Quattrochi. *Scale in Remote Sensing and GIS*. Routledge, New York, 1 edition, January 2023. ISBN 978-0-203-74017-0. doi: 10.1201/9780203740170. URL <https://www.taylorfrancis.com/books/9780203740170>.
- G. Hmimina, E. Dufrêne, J.-Y. Pontailier, N. Delpierre, M. Aubinet, B. Caquet, A. De Grandcourt, B. Burban, C. Flechard, A. Granier, P. Gross, B. Heinesch, B. Longdoz, C. Moureaux, J.-M. Ourcival, S. Rambal, L. Saint André, and K. Soudani. Evaluation of the potential of MODIS satellite data to predict vegetation phenology in different biomes: An investigation using ground-based NDVI measurements. *Remote Sensing of Environment*, 132:145–158, May 2013. ISSN 0034-4257. doi: 10.1016/j.rse.2013.01.010. URL <https://linkinghub.elsevier.com/retrieve/pii/S0034425713000229>. Publisher: Elsevier BV.
- Jeffrey Hollister, Tarak Shah, Jakub Nowosad, Alec L. Robitaille, Marcus W. Beck, and Mike Johnson. *elevatr: Access Elevation Data from Various APIs*. 2023. doi: 10.5281/zenodo.8335450. URL <https://github.com/jhollist/elevatr/>.
- Sha Huang, Lina Tang, Joseph P. Hupy, Yang Wang, and Guofan Shao. A commentary review on the use of normalized difference vegetation index (NDVI) in the era of popular remote sensing. *Journal of Forestry Research*, 32(1):1–6, February 2021. ISSN 1007-662X, 1993-0607. doi: 10.1007/s11676-020-01155-1. URL <https://link.springer.com/10.1007/s11676-020-01155-1>.
- Torres João, Gonçalves João, Marcos Bruno, and Honrado João. Indicator-based assessment of post-fire recovery dynamics using satellite NDVI time-series. *Ecological Indicators*, 89:199–212, June 2018. ISSN 1470-160X. doi: 10.1016/j.ecolind.2018.02.008. URL <https://linkinghub.elsevier.com/retrieve/pii/S1470160X18300852>. Publisher: Elsevier BV.
- Russell Lande. Risks of Population Extinction from Demographic and Environmental Stochasticity and Random Catastrophes. *The American Naturalist*, 142(6):911–927, 1993. ISSN 00030147, 15375323. URL <http://www.jstor.org/stable/2462690>. Publisher: [The University of Chicago Press, The American Society of Naturalists].
- Simon A. Levin. The Problem of Pattern and Scale in Ecology: The Robert H. MacArthur Award Lecture. *Ecology*, 73(6):1943–1967, December 1992. ISSN 0012-9658, 1939-9170. doi: 10.2307/1941447. URL <https://esajournals.onlinelibrary.wiley.com/doi/10.2307/1941447>.
- Richard A. Levins. *Evolution in changing environments: some theoretical explorations*. Number 2 in Monographs in population biology. Princeton Univ. Press, Princeton, NJ, 3. printing edition, 1974. ISBN 978-0-691-08062-8.
- Ramiro Logares and Martin Nuñez. Black Swans in ecology and evolution: The importance of improbable but highly influential events. *Ideas in Ecology and Evolution*, 2012. ISSN 19183178.

doi: 10.4033/iee.2012.5.5.n. URL <https://ojs.library.queensu.ca/index.php/IEE/article/view/4311>.

Paramesha Mallegowda, Ganesan Rengaiyan, Jayalakshmi Krishnan, and Madhura Niphadkar. Assessing Habitat Quality of Forest-Corridors through NDVI Analysis in Dry Tropical Forests of South India: Implications for Conservation. *Remote Sensing*, 7(2):1619–1639, February 2015. ISSN 2072-4292. doi: 10.3390/rs70201619. URL <https://www.mdpi.com/2072-4292/7/2/1619>. Publisher: MDPI AG.

Rekha Marcus and Michael J. Noonan. Changing variability is an overlooked aspect of protected area planning, October 2023. URL <http://biorxiv.org/lookup/doi/10.1101/2023.10.26.564062>.

Rekha Marcus, Stefano Mezzini, Dwija Desai, and Michael Noonan. Planning for a variable future: Environmental stochasticity, species richness, and protected areas in Canada, October 2025. URL <https://www.researchsquare.com/article/rs-7665738/v1>.

Bunkei Matsushita, Wei Yang, Jin Chen, Yuyichi Onda, and Guoyu Qiu. Sensitivity of the Enhanced Vegetation Index (EVI) and Normalized Difference Vegetation Index (NDVI) to Topographic Effects: A Case Study in High-density Cypress Forest. *Sensors*, 7(11):2636–2651, November 2007. ISSN 1424-8220. doi: 10.3390/s7112636. URL <https://www.mdpi.com/1424-8220/7/11/2636>.

Jerod A. Merkle, Kevin L. Monteith, Ellen O. Aikens, Matthew M. Hayes, Kent R. Hersey, Arthur D. Middleton, Brendan A. Oates, Hall Sawyer, Brandon M. Scurlock, and Matthew J. Kauffman. Large herbivores surf waves of green-up during spring. *Proceedings of the Royal Society B: Biological Sciences*, 283(1833):20160456, June 2016. ISSN 0962-8452, 1471-2954. doi: 10.1098/rspb.2016.0456. URL <https://royalsocietypublishing.org/doi/10.1098/rspb.2016.0456>.

Stefano Mezzini, Christen H. Fleming, E. Patrícia Medici, and Michael J. Noonan. How resource abundance and resource stochasticity affect organisms’ range sizes. *Movement Ecology*, 13(1):20, March 2025. ISSN 2051-3933. doi: 10.1186/s40462-025-00546-5. URL <https://movementecologyjournal.biomedcentral.com/articles/10.1186/s40462-025-00546-5>.

Arthur D. Middleton, Jerod A. Merkle, Douglas E. McWhirter, John G. Cook, Rachel C. Cook, P. J. White, and Matthew J. Kauffman. Green-wave surfing increases fat gain in a migratory ungulate. *Oikos*, 127(7):1060–1068, July 2018. ISSN 00301299. doi: 10.1111/oik.05227. URL <https://onlinelibrary.wiley.com/doi/10.1111/oik.05227>.

Pedro R. Mutti, Paulo S. Lúcio, Vincent Dubreuil, and Bergson G. Bezerra. NDVI time series stochastic models for the forecast of vegetation dynamics over desertification hotspots. *International Journal of Remote Sensing*, 41(7):2759–2788, April 2020. ISSN 0143-1161, 1366-5901. doi: 10.1080/01431161.2019.1697008. URL <https://www.tandfonline.com/doi/full/10.1080/01431161.2019.1697008>. Publisher: Informa UK Limited.

Erlend B. Nilsen, Ivar Herfindal, and John D. C. Linnell. Can intra-specific variation in carnivore home-range size be explained using remote-sensing estimates of environmental productivity? *Éco-*

472 *science*, 12(1):68–75, January 2005. ISSN 1195-6860, 2376-7626. doi: 10.2980/i1195-6860-12-1-68.
 473 1. URL <https://www.tandfonline.com/doi/full/10.2980/i1195-6860-12-1-68.1>.

474 Masataka Okabe and Kei Ito. Color universal design (CUD): How to make figures and presentations
 475 that are friendly to colorblind people, 2008. URL <https://jfly.uni-koeln.de/color/#pallett>.

476 David M. Olson, Eric Dinerstein, Eric D. Wikramanayake, Neil D. Burgess, George V. N. Powell,
 477 Emma C. Underwood, Jennifer A. D’amico, Illanga Itoua, Holly E. Strand, John C. Morrison,
 478 Colby J. Loucks, Thomas F. Allnutt, Taylor H. Ricketts, Yumiko Kura, John F. Lamoreux,
 479 Wesley W. Wettengel, Prashant Hedao, and Kenneth R. Kassem. Terrestrial Ecoregions of the
 480 World: A New Map of Life on Earth. *BioScience*, 51(11):933, 2001. ISSN 0006-3568. doi: 10.
 481 1641/0006-3568(2001)051[0933:teotwa]2.0.co;2. URL [https://academic.oup.com/bioscience/](https://academic.oup.com/bioscience/article/51/11/933-938/227116)
 482 [article/51/11/933-938/227116](https://academic.oup.com/bioscience/article/51/11/933-938/227116). Publisher: Oxford University Press (OUP).

483 N Pettorelli, S Ryan, T Mueller, N Bunnefeld, B Jedrzejewska, M Lima, and K Kausrud. The
 484 Normalized Difference Vegetation Index (NDVI): unforeseen successes in animal ecology. *Climate*
 485 *Research*, 46(1):15–27, January 2011. ISSN 0936-577X, 1616-1572. doi: 10.3354/cr00936. URL
 486 <http://www.int-res.com/abstracts/cr/v46/n1/p15-27/>.

487 Nathalie Pettorelli, Jon Olav Vik, Atle Mysterud, Jean-Michel Gaillard, Compton J. Tucker,
 488 and Nils Chr. Stenseth. Using the satellite-derived NDVI to assess ecological responses to en-
 489 vironmental change. *Trends in Ecology & Evolution*, 20(9):503–510, September 2005. ISSN
 490 01695347. doi: 10.1016/j.tree.2005.05.011. URL [https://linkinghub.elsevier.com/retrieve/](https://linkinghub.elsevier.com/retrieve/pii/S016953470500162X)
 491 [pii/S016953470500162X](https://linkinghub.elsevier.com/retrieve/pii/S016953470500162X).

492 Leo Polansky, Werner Kilian, and George Wittemyer. Elucidating the significance of spatial memory
 493 on movement decisions by African savannah elephants using state–space models. *Proceedings of*
 494 *the Royal Society B: Biological Sciences*, 282(1805):20143042, April 2015. ISSN 0962-8452, 1471-
 495 2954. doi: 10.1098/rspb.2014.3042. URL [https://royalsocietypublishing.org/doi/10.1098/](https://royalsocietypublishing.org/doi/10.1098/rspb.2014.3042)
 496 [rspb.2014.3042](https://royalsocietypublishing.org/doi/10.1098/rspb.2014.3042).

497 Francesco Polazzo, Romana Limberger, Frank Pennekamp, Samuel R. P.-J. Ross, Gavin L. Simpson,
 498 and Owen L. Petchey. Measuring the Response Diversity of Ecological Communities Experiencing
 499 Multifarious Environmental Change. *Global Change Biology*, 30(11):e17594, November 2024. ISSN
 500 1354-1013, 1365-2486. doi: 10.1111/gcb.17594. URL [https://onlinelibrary.wiley.com/doi/](https://onlinelibrary.wiley.com/doi/10.1111/gcb.17594)
 501 [10.1111/gcb.17594](https://onlinelibrary.wiley.com/doi/10.1111/gcb.17594).

502 R Core Team. *R: A Language and Environment for Statistical Computing*. R Foundation for
 503 Statistical Computing, Vienna, Austria, 2024. URL <https://www.R-project.org/>.

504 V.C. Radeloff, M. Dubinin, N.C. Coops, A.M. Allen, T.M. Brooks, M.K. Clayton, G.C. Costa, C.H.
 505 Graham, D.P. Helmers, A.R. Ives, D. Kolesov, A.M. Pidgeon, G. Rapacciuolo, E. Razenkova,
 506 N. Suttidate, B.E. Young, L. Zhu, and M.L. Hobi. The Dynamic Habitat Indices (DHIs) from
 507 MODIS and global biodiversity. *Remote Sensing of Environment*, 222:204–214, March 2019. ISSN

00344257. doi: 10.1016/j.rse.2018.12.009. URL <https://linkinghub.elsevier.com/retrieve/pii/S0034425718305625>.

Nathan Ranc, Francesca Cagnacci, and Paul R. Moorcroft. Memory drives the formation of animal home ranges: Evidence from a reintroduction. *Ecology Letters*, 25(4):716–728, April 2022. ISSN 1461-023X, 1461-0248. doi: 10.1111/ele.13869. URL <https://onlinelibrary.wiley.com/doi/10.1111/ele.13869>.

Colleen E. Reid, Laura D. Kubzansky, Jiayue Li, Jessie L. Shmool, and Jane E. Clougherty. It’s not easy assessing greenness: A comparison of NDVI datasets and neighborhood types and their associations with self-rated health in New York City. *Health & Place*, 54:92–101, November 2018. ISSN 1353-8292. doi: 10.1016/j.healthplace.2018.09.005. URL <https://linkinghub.elsevier.com/retrieve/pii/S1353829218301151>. Publisher: Elsevier BV.

Gregory J. M. Rickbeil, Jerod A. Merkle, Greg Anderson, M. Paul Atwood, Jon P. Beckmann, Eric K. Cole, Alyson B. Courtemanch, Sarah Dewey, David D. Gustine, Matthew J. Kauffman, Douglas E. McWhirter, Tony Mong, Kelly Proffitt, Patrick J. White, and Arthur D. Middleton. Plasticity in elk migration timing is a response to changing environmental conditions. *Global Change Biology*, 25(7):2368–2381, July 2019. ISSN 1354-1013, 1365-2486. doi: 10.1111/gcb.14629. URL <https://onlinelibrary.wiley.com/doi/10.1111/gcb.14629>.

Louise Riotte-Lambert and Jason Matthiopoulos. Environmental Predictability as a Cause and Consequence of Animal Movement. *Trends in Ecology & Evolution*, 35(2):163–174, February 2020. ISSN 01695347. doi: 10.1016/j.tree.2019.09.009. URL <https://linkinghub.elsevier.com/retrieve/pii/S0169534719302885>.

Matteo Rizzuto, Shawn J. Leroux, Eric Vander Wal, Isabella C. Richmond, Travis R. Heckford, Juliana Balluffi-Fry, and Yolanda F. Wiersma. Forage stoichiometry predicts the home range size of a small terrestrial herbivore. *Oecologia*, 197(2):327–338, October 2021. ISSN 0029-8549, 1432-1939. doi: 10.1007/s00442-021-04965-0. URL <https://link.springer.com/10.1007/s00442-021-04965-0>.

Hall Sawyer, Jerod A. Merkle, Arthur D. Middleton, Samantha P. H. Dwinnell, and Kevin L. Monteith. Migratory plasticity is not ubiquitous among large herbivores. *Journal of Animal Ecology*, 88(3):450–460, March 2019. ISSN 0021-8790, 1365-2656. doi: 10.1111/1365-2656.12926. URL <https://besjournals.onlinelibrary.wiley.com/doi/10.1111/1365-2656.12926>.

Gavin L. Simpson. Modelling Palaeoecological Time Series Using Generalised Additive Models. *Frontiers in Ecology and Evolution*, 6:149, October 2018. ISSN 2296-701X. doi: 10.3389/fevo.2018.00149. URL <https://www.frontiersin.org/article/10.3389/fevo.2018.00149/full>.

N.T. Son, C.F. Chen, C.R. Chen, V.Q. Minh, and N.H. Trung. A comparative analysis of multitemporal MODIS EVI and NDVI data for large-scale rice yield estimation. *Agricultural and Forest Meteorology*, 197:52–64, October 2014. ISSN 0168-1923. doi: 10.1016/j.agrformet.2014.06.007.

544 URL <https://linkinghub.elsevier.com/retrieve/pii/S016819231400149X>. Publisher: El-
545 sevier BV.

546 T. R. E. Southwood. Habitat, the Templet for Ecological Strategies? *The Journal of Animal*
547 *Ecology*, 46(2):336, June 1977. ISSN 00218790. doi: 10.2307/3817. URL [https://www.jstor.](https://www.jstor.org/stable/3817?origin=crossref)
548 [org/stable/3817?origin=crossref](https://www.jstor.org/stable/3817?origin=crossref).

549 Constantî Stefanescu, Andreu Ubach, and Christer Wiklund. Timing of mating, reproductive status
550 and resource availability in relation to migration in the painted lady butterfly. *Animal Behaviour*,
551 172:145–153, February 2021. ISSN 00033472. doi: 10.1016/j.anbehav.2020.12.013. URL [https:](https://linkinghub.elsevier.com/retrieve/pii/S0003347220303742)
552 [//linkinghub.elsevier.com/retrieve/pii/S0003347220303742](https://linkinghub.elsevier.com/retrieve/pii/S0003347220303742).

553 Saurabh Steixner-Kumar and Jan Gläsker. Strategies for navigating a dynamic world. *Science*, 369
554 (6507):1056–1057, August 2020. ISSN 0036-8075, 1095-9203. doi: 10.1126/science.abd7258. URL
555 <https://www.science.org/doi/10.1126/science.abd7258>.

556 D. W. Stephens and Eric L. Charnov. Optimal foraging: Some simple stochastic models. *Behavioral*
557 *Ecology and Sociobiology*, 10(4):251–263, July 1982. ISSN 0340-5443, 1432-0762. doi: 10.1007/
558 [BF00302814](http://link.springer.com/10.1007/BF00302814). URL <http://link.springer.com/10.1007/BF00302814>.

559 Bernt-Erik Sæther. Environmental stochasticity and population dynamics of large herbivores: a
560 search for mechanisms. *Trends in Ecology & Evolution*, 12(4):143–149, April 1997. ISSN 0169-
561 5347. doi: 10.1016/S0169-5347(96)10068-9. URL [https://doi.org/10.1016/S0169-5347\(96\)](https://doi.org/10.1016/S0169-5347(96)10068-9)
562 [10068-9](https://doi.org/10.1016/S0169-5347(96)10068-9). Publisher: Elsevier.

563 United States National Aeronautics and Space Administration. Level-1 and Atmosphere Archive &
564 Distribution System Distributed Active Archive Center, 2025. URL [https://ladsweb.modaps.](https://ladsweb.modaps.eosdis.nasa.gov/archive/allData)
565 [eosdis.nasa.gov/archive/allData](https://ladsweb.modaps.eosdis.nasa.gov/archive/allData).

566 United States National Centers for Environmental Information (NCEI) and United States National
567 Aeronautics and Space Administration. NCEI Database, 2025. URL [https://ncei.noaa.gov/](https://ncei.noaa.gov/dataver)
568 [dataver](https://ncei.noaa.gov/dataver).

569 Eric Vermote and NOAA CDR Program. NOAA Climate Data Record (CDR) of AVHRR Normal-
570 ized Difference Vegetation Index (NDVI), Version 5, 2018. URL [https://www.ncei.noaa.gov/](https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01558/html)
571 [metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01558/html](https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01558/html).

572 Eric Vermote and NOAA CDR Program. NOAA Climate Data Record (CDR) of VIIRS Normal-
573 ized Difference Vegetation Index (NDVI), Version 1, 2022. URL [https://www.ncei.noaa.gov/](https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01677/html)
574 [metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01677/html](https://www.ncei.noaa.gov/metadata/geoportal/rest/metadata/item/gov.noaa.ncdc:C01677/html).

575 Amelia A. Wolf, Erika S. Zavaleta, and Paul C. Selman. Flowering phenology shifts in response
576 to biodiversity loss. *Proceedings of the National Academy of Sciences*, 114(13):3463–3468, March
577 2017. ISSN 0027-8424, 1091-6490. doi: 10.1073/pnas.1608357114. URL [http://www.pnas.org/](http://www.pnas.org/lookup/doi/10.1073/pnas.1608357114)
578 [lookup/doi/10.1073/pnas.1608357114](http://www.pnas.org/lookup/doi/10.1073/pnas.1608357114).

579 E. M. Wolkovich, B. I. Cook, J. M. Allen, T. M. Crimmins, J. L. Betancourt, S. E. Travers, S. Pau,
580 J. Regetz, T. J. Davies, N. J. B. Kraft, T. R. Ault, K. Bolmgren, S. J. Mazer, G. J. McCabe, B. J.
581 McGill, C. Parmesan, N. Salamin, M. D. Schwartz, and E. E. Cleland. Warming experiments
582 underpredict plant phenological responses to climate change. *Nature*, 485(7399):494–497, May
583 2012. ISSN 0028-0836, 1476-4687. doi: 10.1038/nature11014. URL [http://www.nature.com/
584 articles/nature11014](http://www.nature.com/articles/nature11014).

585 Simon N. Wood. *Generalized additive models: an introduction with R*. Chapman & Hall/CRC texts
586 in statistical science. CRC Press/Taylor & Francis Group, Boca Raton, second edition edition,
587 2017. ISBN 978-1-4987-2833-1.

588 Simon N. Wood and Matteo Fasiolo. A Generalized Fellner-Schall Method for Smoothing Parameter
589 Optimization with Application to Tweedie Location, Scale and Shape Models. *Biometrics*, 73
590 (4):1071–1081, December 2017. ISSN 0006-341X, 1541-0420. doi: 10.1111/biom.12666. URL
591 <https://academic.oup.com/biometrics/article/73/4/1071-1081/7537785>.

592 Simon N. Wood, Yannig Goude, and Simon Shaw. Generalized additive models for large data sets.
593 *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 64(1):139–155, January
594 2015. ISSN 00359254. doi: 10.1111/rssc.12068. URL [https://onlinelibrary.wiley.com/doi/
595 10.1111/rssc.12068](https://onlinelibrary.wiley.com/doi/10.1111/rssc.12068).

596 Simon N. Wood, Zheyuan Li, Gavin Shaddick, and Nicole H. Augustin. Generalized Additive
597 Models for Gigadata: Modeling the U.K. Black Smoke Network Daily Data. *Journal of the
598 American Statistical Association*, 112(519):1199–1210, July 2017. ISSN 0162-1459, 1537-274X.
599 doi: 10.1080/01621459.2016.1195744. URL [https://www.tandfonline.com/doi/full/10.1080/
600 01621459.2016.1195744](https://www.tandfonline.com/doi/full/10.1080/01621459.2016.1195744).

601 Xueyan Zhang, Yunfeng Hu, Dafang Zhuang, Yongqing Qi, and Xin Ma. NDVI spatial pattern
602 and its differentiation on the Mongolian Plateau. *Journal of Geographical Sciences*, 19(4):403–
603 415, August 2009. ISSN 1009-637X, 1861-9568. doi: 10.1007/s11442-009-0403-7. URL [http:
604 //link.springer.com/10.1007/s11442-009-0403-7](http://link.springer.com/10.1007/s11442-009-0403-7). Publisher: Springer Science and Business
605 Media LLC.