

# DENVar: A global Dynamic Estimate of NDVI Variance

## Appendix A

Stefano Mezzini<sup>1,2</sup>      Gavin L. Simpson<sup>3</sup>      Michael J. Noonan<sup>1,2,4,\*</sup>

<sup>1</sup> Okanagan Institute for Biodiversity, Resilience, and Ecosystem Services, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.


<sup>2</sup> Department of Biology, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

<sup>3</sup> Department of Animal & Veterinary Sciences, Aarhus University, Denmark.


<sup>4</sup> Department of Computer Science, Math, Physics, and Statistics, The University of British Columbia Okanagan, Kelowna, British Columbia, Canada.

\* Correspondence: [Michael J. Noonan <michael.noonan@ubc.ca>](mailto:michael.noonan@ubc.ca)


Figure A1 below illustrates the color palettes used in the main manuscript and this appendix, for each type of vision (trichromatic, deuteranope, protanope, tritanope, and achromatic). The NDVI palette is a modified version of the Bukavu palette by Fabio Crameri (Crameri, 2018a b). Palette A3 is a modified version of a palette by Masataka Okabe and Kei Ito (Okabe & Ito, 2008). All other palettes were created by Fabio Crameri (Crameri, 2018a b). All original palettes are available via the `khroma` package (Frerebeau, 2024) for R (R Core Team, 2024). Figures A2-A5 show some of the data used for fitting the models for estimating mean NDVI and the variance around the mean. The code for generating the figures is available on GitHub at <https://github.com/QuantitativeEcologyLab/ndvi-stochasticity/blob/main/analysis/figures/input-data.R>.




**Figure A1:** Color palettes used in the main manuscript and the figures below.




**Figure A2:** Number of days with a raster within each 2-day period. used for temporal aggregation of the data.



**Figure A3:** To avoid data size limitations in R, data were split into groups of WWF realms. Also see Fig. A4 below.



**Figure A4:** Map of WWF biomes. Also see Fig. A3 above.



**Figure A5:** Map of elevation above sea level. All elevations below the lowest altitude in the Dead Sea (-432 m) were set to -432 m to remove excessively negative altitudes near coast lines. The `elevatr` package for R did not provide elevations for the southernmost region of Antarctica.



## References

- Crameri F. (2018a). Geodynamic diagnostics, scientific visualisation and StagLab 3.0. *Geoscientific Model Development* **11**, 2541–2562. <https://doi.org/10.5194/gmd-11-2541-2018>
- Crameri F. (2018b). [Scientific colour-maps](#)
- Frerebeau N. (2024). *Khroma: Colour Schemes for Scientific Data Visualization*. Université Bordeaux Montaigne, Pessac, France.
- Okabe M. & Ito K. (2008). [Color universal design \(CUD\): How to make figures and presentations that are friendly to colorblind people](#)
- R Core Team (2024). *R: A Language and Environment for Statistical Computing*. R Foundation for Statistical Computing, Vienna, Austria.