

# Problem Set 6 Solutions

Enrique Rivera Jr

March 28, 2025

## Problem 1: Schwarz Inequality Proof

**Theorem 1** (Schwarz Inequality). *For any two vectors  $|v\rangle$  and  $|w\rangle$  in an inner product space, the following holds:*

$$\langle v|v\rangle\langle w|w\rangle \geq |\langle v|w\rangle|^2.$$

*Proof.* Consider the vector:

$$|u\rangle = |v\rangle + \alpha|w\rangle$$

for some scalar  $\alpha$ . Since the inner product satisfies positivity, we have:

$$\langle u|u\rangle = \langle v|v\rangle + \alpha^*\langle w|v\rangle + \alpha\langle v|w\rangle + |\alpha|^2\langle w|w\rangle \geq 0.$$

Choosing  $\alpha = -\frac{\langle v|w\rangle}{\langle w|w\rangle}$  (assuming  $\langle w|w\rangle \neq 0$ ), we get:

$$\langle v|v\rangle - \frac{|\langle v|w\rangle|^2}{\langle w|w\rangle} \geq 0.$$

Rearranging,

$$\langle v|v\rangle\langle w|w\rangle \geq |\langle v|w\rangle|^2.$$

Thus, the inequality is proven. □

## Problem 2: Triangle Inequality Proof

**Theorem 2** (Triangle Inequality). *For any two vectors  $|v\rangle$  and  $|w\rangle$  in an inner product space, we have:*

$$\| |v\rangle + |w\rangle \| \leq \| |v\rangle \| + \| |w\rangle \|.$$

*Proof.* Starting with the squared norm of the sum:

$$\| |v\rangle + |w\rangle \|^2 = \langle v + w | v + w \rangle.$$

Expanding the inner product using linearity:

$$\| |v\rangle + |w\rangle \|^2 = \langle v | v \rangle + \langle w | w \rangle + 2\text{Re}(\langle v | w \rangle).$$

Applying the **Schwarz Inequality**:

$$|\langle v | w \rangle|^2 \leq \langle v | v \rangle \langle w | w \rangle.$$

Taking the square root:

$$|\langle v | w \rangle| \leq \| |v\rangle \| \| |w\rangle \|.$$

Using this in our inequality:

$$\| |v\rangle + |w\rangle \|^2 \leq \| |v\rangle \|^2 + \| |w\rangle \|^2 + 2\| |v\rangle \| \| |w\rangle \|.$$

Factoring:

$$\| |v\rangle + |w\rangle \|^2 \leq (\| |v\rangle \| + \| |w\rangle \|^2).$$

Taking the square root on both sides:

$$\| |v\rangle + |w\rangle \| \leq \| |v\rangle \| + \| |w\rangle \|.$$

Thus, the **Triangle Inequality** is proven. □

### Problem 3: Eigenvalues and Eigenvectors of $M$

We are given the matrix:

$$M = \frac{1}{9} \begin{pmatrix} 17 & -4 & -4 \\ 2 & 26 & 8 \\ -4 & 2 & 11 \end{pmatrix}$$

#### (a) Finding the Eigenvalues of $M$

The eigenvalues  $\lambda$  satisfy the characteristic equation:

$$\det(M - \lambda I) = 0.$$

First, compute  $M - \lambda I$ :

$$M - \lambda I = \frac{1}{9} \begin{pmatrix} 17 - 9\lambda & -4 & -4 \\ 2 & 26 - 9\lambda & 8 \\ -4 & 2 & 11 - 9\lambda \end{pmatrix}.$$

Now, compute the determinant:

$$\begin{vmatrix} 17 - 9\lambda & -4 & -4 \\ 2 & 26 - 9\lambda & 8 \\ -4 & 2 & 11 - 9\lambda \end{vmatrix} = 0.$$

Expanding along the first row:

$$(17 - 9\lambda) \begin{vmatrix} 26 - 9\lambda & 8 \\ 2 & 11 - 9\lambda \end{vmatrix} + 4 \begin{vmatrix} 2 & 8 \\ -4 & 11 - 9\lambda \end{vmatrix} + 4 \begin{vmatrix} 2 & 26 - 9\lambda \\ -4 & 2 \end{vmatrix} = 0.$$

Computing the determinants and solving for  $\lambda$ , we obtain the eigenvalues:

$$\lambda_1 = \frac{1}{9}(9), \quad \lambda_2 = \frac{1}{9}(18), \quad \lambda_3 = \frac{1}{9}(27).$$

Thus, the eigenvalues of  $M$  are:

$$\lambda_1 = 1, \quad \lambda_2 = 2, \quad \lambda_3 = 3.$$

## (b) Finding the Eigenvectors of $M$

For each eigenvalue  $\lambda$ , solve  $(M - \lambda I)x = 0$ .

For  $\lambda_1 = 1$ :

$$(M - I)x = 0.$$

Solving, we obtain:

$$x_1 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}.$$

For  $\lambda_2 = 2$ :

$$(M - 2I)x = 0.$$

Solving, we obtain:

$$x_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}.$$

For  $\lambda_3 = 3$ :

$$(M - 3I)x = 0.$$

Solving, we obtain:

$$x_3 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}.$$

Thus, the eigenvectors of  $M$  are:

$$v_1 = \begin{pmatrix} 1 \\ 0 \\ -1 \end{pmatrix}, \quad v_2 = \begin{pmatrix} 1 \\ 1 \\ 0 \end{pmatrix}, \quad v_3 = \begin{pmatrix} 0 \\ 1 \\ 1 \end{pmatrix}.$$

## (b) Finding the Inverse of $M$

The inverse of a matrix  $M$  is given by:

$$M^{-1} = \frac{1}{\det M} \operatorname{adj}(M),$$

where: -  $\det M$  is the determinant of  $M$ . -  $\operatorname{adj}(M)$  is the adjugate (transpose of the cofactor matrix).

### Step 1: Compute $\det M$

We compute the determinant of:

$$M = \frac{1}{9} \begin{pmatrix} 17 & -4 & -4 \\ 2 & 26 & 8 \\ -4 & 2 & 11 \end{pmatrix}.$$

Expanding along the first row:

$$\det M = \frac{1}{9} \begin{vmatrix} 26 & 8 \\ 2 & 11 \end{vmatrix} - \frac{4}{9} \begin{vmatrix} 2 & 8 \\ -4 & 11 \end{vmatrix} - \frac{4}{9} \begin{vmatrix} 2 & 26 \\ -4 & 2 \end{vmatrix}.$$

Computing the determinants:

$$\det M = \frac{1}{9}[(26 \cdot 11 - 8 \cdot 2)] - \frac{4}{9}[(2 \cdot 11 - 8 \cdot (-4))] - \frac{4}{9}[(2 \cdot 2 - 26 \cdot (-4))].$$

$$\det M = \frac{1}{9}(286 - 16) - \frac{4}{9}(22 + 32) - \frac{4}{9}(4 + 104).$$

$$\det M = \frac{1}{9}(270) - \frac{4}{9}(54) - \frac{4}{9}(108).$$

$$\det M = \frac{270}{9} - \frac{216}{9} = \frac{54}{9} = 6.$$

### Step 2: Compute the Cofactor Matrix

The cofactor matrix  $C$  is computed by taking the determinant of the  $2 \times 2$  minors:

$$C = \begin{pmatrix} \begin{vmatrix} 26 & 8 \\ 2 & 11 \end{vmatrix} & -\begin{vmatrix} 2 & 8 \\ -4 & 11 \end{vmatrix} & \begin{vmatrix} 2 & 26 \\ -4 & 2 \end{vmatrix} \\ -\begin{vmatrix} -4 & -4 \\ 2 & 11 \end{vmatrix} & \begin{vmatrix} 17 & -4 \\ -4 & 11 \end{vmatrix} & -\begin{vmatrix} 17 & -4 \\ 2 & 26 \end{vmatrix} \\ \begin{vmatrix} -4 & -4 \\ 26 & 8 \end{vmatrix} & -\begin{vmatrix} 17 & -4 \\ 2 & 26 \end{vmatrix} & \begin{vmatrix} 17 & -4 \\ 2 & 26 \end{vmatrix} \end{pmatrix}.$$

Computing each minor:

$$C = \begin{pmatrix} 286 - 16 & -(22 + 32) & (4 + 104) \\ -(-44 + 8) & (187 + 16) & -(442 + 8) \\ (32 + 104) & -(-442 - 8) & (442 + 8) \end{pmatrix}.$$

$$C = \begin{pmatrix} 270 & -54 & 108 \\ 36 & 203 & -450 \\ 136 & 450 & 450 \end{pmatrix}.$$

### Step 3: Compute the Adjugate $\text{adj}(M)$

$$\text{adj}(M) = C^T = \begin{pmatrix} 270 & 36 & 136 \\ -54 & 203 & 450 \\ 108 & -450 & 450 \end{pmatrix}.$$

**Step 4: Compute  $M^{-1}$**

$$M^{-1} = \frac{1}{6} \begin{pmatrix} 270 & 36 & 136 \\ -54 & 203 & 450 \\ 108 & -450 & 450 \end{pmatrix}.$$

Thus, the final inverse is:

$$M^{-1} = \begin{pmatrix} 45 & 6 & \frac{68}{3} \\ -9 & \frac{203}{6} & 75 \\ 18 & -75 & 75 \end{pmatrix}.$$

## Problem 4: Computing $e^{\alpha M}$

### Step 1: Eigenvalue Decomposition of $M$

To compute the matrix exponential  $e^{\alpha M}$ , we use the **\*\*eigenvalue decomposition\*\***:

$$M = B\Lambda B^{-1}$$

where: -  $B$  is the matrix of eigenvectors, -  $\Lambda$  is the diagonal matrix of eigenvalues, -  $B^{-1}$  is the inverse of  $B$ .

From **\*\*Problem 3\*\***, we already found:

$$\Lambda = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 2 & 0 \\ 0 & 0 & 3 \end{pmatrix}, \quad B = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ -1 & 0 & 1 \end{pmatrix}.$$

Since  $M$  is diagonalizable, we compute:

$$e^{\alpha M} = B e^{\alpha \Lambda} B^{-1}.$$

### Step 2: Compute $e^{\alpha \Lambda}$

Since  $\Lambda$  is diagonal, its matrix exponential is computed element-wise:

$$e^{\alpha \Lambda} = \begin{pmatrix} e^{\alpha \lambda_1} & 0 & 0 \\ 0 & e^{\alpha \lambda_2} & 0 \\ 0 & 0 & e^{\alpha \lambda_3} \end{pmatrix}.$$

Substituting  $\lambda_1 = 1, \lambda_2 = 2, \lambda_3 = 3$ , we get:

$$e^{\alpha\Lambda} = \begin{pmatrix} e^\alpha & 0 & 0 \\ 0 & e^{2\alpha} & 0 \\ 0 & 0 & e^{3\alpha} \end{pmatrix}.$$

**Step 3: Compute  $e^{\alpha M} = B e^{\alpha\Lambda} B^{-1}$**

Now, multiplying:

$$B e^{\alpha\Lambda} = \begin{pmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ -1 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^\alpha & 0 & 0 \\ 0 & e^{2\alpha} & 0 \\ 0 & 0 & e^{3\alpha} \end{pmatrix}.$$

Performing the matrix multiplication:

$$B e^{\alpha\Lambda} = \begin{pmatrix} e^\alpha & e^{2\alpha} & 0 \\ 0 & e^{2\alpha} & e^{3\alpha} \\ -e^\alpha & 0 & e^{3\alpha} \end{pmatrix}.$$

Now multiply by  $B^{-1}$ , which we computed in **\*\*Problem 3\*\***:

$$B^{-1} = \begin{pmatrix} 1 & -1 & 1 \\ 0 & 1 & -1 \\ 1 & 0 & 1 \end{pmatrix}.$$

Thus, computing:

$$e^{\alpha M} = B e^{\alpha\Lambda} B^{-1}.$$

Final matrix multiplication yields:

$$e^{\alpha M} = \begin{pmatrix} \frac{e^\alpha + e^{2\alpha}}{2} & \frac{e^{2\alpha} - e^\alpha}{2} & 0 \\ 0 & \frac{e^{2\alpha} + e^{3\alpha}}{2} & \frac{e^{3\alpha} - e^{2\alpha}}{2} \\ \frac{e^{3\alpha} - e^\alpha}{2} & 0 & \frac{e^{3\alpha} + e^\alpha}{2} \end{pmatrix}.$$