

Problem Set 8 Solutions

Enrique Rivera Jr

March 26, 2025

Question 1(a): Ground State and First Excited State of the 3D Isotropic Harmonic Oscillator

Hamiltonian and Separation of Variables

The Hamiltonian for the 3D isotropic harmonic oscillator is:

$$\hat{H} = \frac{\hat{p}_x^2 + \hat{p}_y^2 + \hat{p}_z^2}{2m} + \frac{1}{2}m\omega^2(x^2 + y^2 + z^2). \quad (1)$$

Because it *separates* into a sum of three identical 1D harmonic oscillators, the wavefunction can be written as a product:

$$\Psi_{n_x, n_y, n_z}(x, y, z) = \psi_{n_x}(x) \psi_{n_y}(y) \psi_{n_z}(z), \quad (2)$$

where each $\psi_n(x)$ is the standard 1D harmonic oscillator solution.

1D Review

1D ground state:

$$\psi_0(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} \exp\left[-\frac{m\omega}{2\hbar} x^2\right]. \quad (3)$$

1D first excited state:

$$\psi_1(x) = \left(\frac{m\omega}{\pi\hbar}\right)^{1/4} \sqrt{2} \alpha x \exp\left[-\frac{m\omega}{2\hbar} x^2\right] \quad \text{where} \quad \alpha = \left(\frac{m\omega}{\hbar}\right)^{1/2}. \quad (4)$$

Ground State in 3D

For $n_x = n_y = n_z = 0$, the 3D ground state is:

$$\Psi_{0,0,0}(x, y, z) = \psi_0(x) \psi_0(y) \psi_0(z) = \left(\frac{m\omega}{\pi\hbar}\right)^{3/4} \exp\left[-\frac{m\omega}{2\hbar}(x^2 + y^2 + z^2)\right]. \quad (5)$$

First Excited States in 3D

The energy in 3D is $\hbar\omega (n_x + n_y + n_z + \frac{3}{2})$. The first excited level occurs when $n_x + n_y + n_z = 1$. That yields three possible combinations:

$$\begin{aligned}(n_x, n_y, n_z) &= (1, 0, 0), \\ (n_x, n_y, n_z) &= (0, 1, 0), \\ (n_x, n_y, n_z) &= (0, 0, 1).\end{aligned}$$

Thus we have three degenerate first-excited states:

$$\begin{aligned}\Psi_{1,0,0}(x, y, z) &= \psi_1(x) \psi_0(y) \psi_0(z), \\ \Psi_{0,1,0}(x, y, z) &= \psi_0(x) \psi_1(y) \psi_0(z), \\ \Psi_{0,0,1}(x, y, z) &= \psi_0(x) \psi_0(y) \psi_1(z),\end{aligned}$$

all with energy $E = \frac{5}{2}\hbar\omega$.

Final Results

- **Ground State:**

$$\Psi_{0,0,0}(x, y, z) = \left(\frac{m\omega}{\pi\hbar}\right)^{3/4} \exp\left[-\frac{m\omega}{2\hbar}(x^2 + y^2 + z^2)\right]. \quad (6)$$

- **First-Excited States (3-fold degenerate):**

$$\begin{aligned}\Psi_{1,0,0}(x, y, z) &= \psi_1(x) \psi_0(y) \psi_0(z), \\ \Psi_{0,1,0}(x, y, z) &= \psi_0(x) \psi_1(y) \psi_0(z), \\ \Psi_{0,0,1}(x, y, z) &= \psi_0(x) \psi_0(y) \psi_1(z),\end{aligned}$$

- **Energy at first-excited level:** $E = \frac{5}{2}\hbar\omega$.

Question 1(b): Energy and Degeneracy of the 3D Isotropic Harmonic Oscillator

Energy of the n-th Excited State

For a 1D harmonic oscillator, the energy levels are

$$E_n^{(1D)} = \hbar\omega\left(n + \frac{1}{2}\right), \quad n = 0, 1, 2, \dots \quad (7)$$

In 3D, we have three independent 1D oscillators, so the total energy is

$$E_{n_x, n_y, n_z} = \hbar\omega \left(n_x + n_y + n_z + \frac{3}{2} \right). \quad (8)$$

Defining $n = n_x + n_y + n_z$, the n -th excited level has energy

$$\boxed{E_n = \hbar\omega \left(n + \frac{3}{2} \right)}. \quad (9)$$

Degeneracy of the n -th Excited State

The states with $n_x + n_y + n_z = n$ share the same energy E_n . The number of nonnegative integer solutions to $n_x + n_y + n_z = n$ is

$$\binom{n+3-1}{3-1} = \binom{n+2}{2} = \frac{(n+1)(n+2)}{2}. \quad (10)$$

Hence, the degeneracy is

$$\boxed{g_n = \frac{(n+1)(n+2)}{2}}. \quad (11)$$

Construction Using Raising Operators

Let a_i, a_i^\dagger be the annihilation and creation operators for each Cartesian direction $i = 1, 2, 3$. The 3D ground state $|\psi_0\rangle$ satisfies

$$a_i |\psi_0\rangle = 0, \quad i = 1, 2, 3. \quad (12)$$

Then any excited state with quantum numbers (n_x, n_y, n_z) can be built by applying the appropriate creation operators:

$$|n_x, n_y, n_z\rangle \propto (a_1^\dagger)^{n_x} (a_2^\dagger)^{n_y} (a_3^\dagger)^{n_z} |\psi_0\rangle. \quad (13)$$

Hence, the n -th excited state ($n = n_x + n_y + n_z$) can be written in the form

$$\boxed{(a_1^\dagger)^{n_1} (a_2^\dagger)^{n_2} (a_3^\dagger)^{n_3} |\psi_0\rangle \quad \text{with} \quad n_1 + n_2 + n_3 = n.} \quad (14)$$

Summary for Part (b)

- **Energy:** $E_n = \hbar\omega \left(n + \frac{3}{2} \right)$.
- **Degeneracy:** $g_n = \frac{(n+1)(n+2)}{2}$.
- **Raising-Operator Representation:** states in the n -th manifold are generated by distributing n creation-operator actions among the three coordinates.