

Learning-Induced Metal-Insulator Transition

Exploratory DIII monitored-Majorana study with an XY-line validation gate

We reproduce the known XY-line learning transition, then test a generic symmetry-class-DIII measurement cut with Born-sampled Gaussian Majorana trajectories. The report explains the mapping, algorithms, parameters, finite-size analysis, anisotropy calibration, uncertainty, and strict exploratory claim gates.

Team 卧龙凤雏

Executive result and scope

Exploratory status

The frozen status is `xy_reproduced_diii_inconclusive`. The directly fitted universal candidate is `c_eff`
 $\alpha=0.349033$; α is not stable, so no standalone `c_eff` is published.




Figure 1. XY validation scan and declared reference window.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Basic concepts: monitoring and phases

Repeated quantum measurements can either preserve extended Majorana correlations or localize them. The metal-insulator distinction is diagnosed from size trends, entanglement-arc model weights, correlations, and record free energy rather than from one smooth curve.

$$\sigma(\theta, \varphi) = \sin\theta \cos\varphi X + \sin\theta \sin\varphi Y + \cos\theta Z$$

The two angles define the physical single-qubit measurement axis.

Surface-code mapping and effective couplings

The surface-code Born tensor network maps to alternating Gaussian measurements and rotations. The real coupling controls information gain; the complex phase controls norm-preserving Majorana rotation.

$$J(\theta) = \text{atanh}(\cos\theta)$$

This converts the polar measurement angle into a real monitoring strength.




Figure 2. Representative entanglement arcs across angles and widths.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Why generic DIII differs from the XY class-D line

The XY line has a special class-D block decomposition. Moving to $\theta=0.45\pi$ with nonzero azimuth removes that decomposition and realizes the generic DIII cut targeted by the open challenge.




Figure 3. Generic DIII coarse evidence and selected refinement bracket.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Gaussian covariance implementation

Rust evolves a real antisymmetric $2L \times 2L$ covariance matrix. Rational measurement updates and orthogonal rotations preserve purity analytically; after each period, an orthogonal polar projection removes accumulated roundoff in $\Gamma^2 = -I$. Entropy follows from duplicated singular values of restricted covariance matrices.

$$\Gamma' = R \Gamma R$$

Every unitary Majorana gate acts by congruence on the covariance matrix.

Conditional Born sampling and the IID negative control

Xoshiro256++ draws conditional Born outcomes. The Rao-Blackwellized binary entropy is accumulated before each draw. Unbiased IID signs are a deliberately nonphysical diagnostic and never enter physical summaries.




Figure 4. Born sampling compared with the nonphysical IID-sign control.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Parameters, runtime budget, and estimator meanings

Table 1. Parameter settings

Parameter	Setting	Meaning
Widths L	8, 12, 16, 20, 24, 28, 32	finite-size scaling
Independent streams	8	stream-level uncertainty
Ordinary stop	3300	stop launching new work
Hard stop	5100	atomic finalization only

Times are measured from the production simulation and analysis pipeline.




Figure 5. Runtime allocation and effective sample size.

Interpretation limit: Finite widths and declared fit windows limit interpretation.

Independent mathematical and physical oracles

Dense Hilbert-space enumeration at $L=2$ checks joint probabilities, covariances, and entropy independently. Analytic angle limits, the frozen weak-self-dual kernel, exact-Y swaps, and class-D decomposition residuals guard sign and factor conventions.

Table 2. Scientific oracles

Oracle	maximum error
dense Born probability	1.006e-16
dense covariance	1.776e-15
weak self-dual limit	not recorded

Production is gated on these checks.

Reproduction of the known XY transition

The XY scan is a validation gate: its finite-size bracket must overlap the declared reference window before any generic-DIII claim is considered.




Figure 6. XY validation scan and declared reference window.

Interpretation limit: Finite widths and declared fit windows limit interpretation.

Exploratory generic-DIII phase evidence

The generic scan compares area, logarithmic, squared-logarithmic, and Page-augmented arc models across widths. A transition bracket requires persistent opposite phase evidence on adjacent angles.




Figure 7. Generic DIII coarse evidence and selected refinement bracket.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 8. Arc-model coefficients used to distinguish area, logarithmic, and squared-logarithmic behavior.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Casimir amplitude and finite-size correction

$$\gamma(L) = f_{\infty} L - \pi(c_{\text{eff}} \alpha) / (6L) + a/L^3$$

The coefficient of $1/L$ is the directly observed product $c_{\text{eff}} \alpha$; the L^{-3} term absorbs leading finite-size drift.




Figure 9. Casimir finite-size fit with residuals.

Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 10. Hierarchical bootstrap distribution of $c_{\text{eff}} \alpha$.

Interpretation limit: Finite widths and declared fit windows limit interpretation.

Spatial-temporal anisotropy calibration

Spatial parity correlations determine Δ , while the leading temporal Lyapunov gap determines α . Window changes and block deletions must agree before dividing the Casimir amplitude by α .

$$\alpha = gL / (2\pi\Delta)$$

This calibration makes the spacetime conversion explicit instead of assuming isotropy.




Figure 11. Spatial parity decay and temporal Lyapunov gaps.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 12. Anisotropy sensitivity to analysis windows.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Dual effective-central-charge fits and cross-validation

Exploratory number versus universal constant

All finite exploratory estimates are displayed with intervals and failed gates. Only a bracketed transition, stable fits and anisotropy, sufficient streams and blocks, and estimator agreement can promote the value to candidate status.

$$S(\ell, L) = b + (c_{\text{eff}} S(L)/3) \log[(L/\pi) \sin(\pi\ell/L)] + q \cos(2\pi\ell/L)/L^2$$

The coefficient of the periodic chord-length logarithm defines $c_{\text{eff}} S(L)$; the central interval suppresses endpoint effects and the cosine term absorbs the leading oscillatory correction.

The entanglement estimate is extrapolated linearly in $1/L^2$. Independently, the Casimir fit measures c_{eff} and the Lyapunov/spatial calibration determines a . Their agreement is tested after fitting and is never imposed.

Table 3. Effective central charge fits

Estimator	Point estimate	95% interval	Status
Entanglement chord length	3.061e+00	[1.57163, 4.54985]	available
Casimir / anisotropy	1.258e+01	[10.7278, 14.6103]	available
Claim level	1.258e+01	[10.7278, 14.6103]	exploratory

Failed gates: diii_transition_not_bracketed, anisotropy_unstable, estimator_disagreement




Figure 13. Entropy versus periodic chord length at the selected DIII angle.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 14. Finite-size extrapolation of the entanglement effective central charge.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 15. Casimir finite-size fit with residuals.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 16. Residuals of the Casimir finite-size fit.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 17. Anisotropy stability across declared analysis windows.
Interpretation limit: Finite widths and declared fit windows limit interpretation.




Figure 18. Independent entanglement and Casimir effective-central-charge estimates.
Interpretation limit: Finite widths and declared fit windows limit interpretation.

Uncertainty, sensitivity, and claim boundary

Uncertainty is hierarchical: streams are resampled first and complete blocks second. Sensitivity checks change minimum width, corrections, phase brackets, correlation windows, and Lyapunov block deletions.

Claim boundary

This is an exploratory finite-size result, not a final universal constant. Finite estimates remain visible with uncertainty intervals and failed gates; only results passing every gate are labeled candidates.

Reproducibility and code/data inventory

All physics sampling is implemented in Rust; Python only validates frozen artifacts, fits models, bootstraps, plots, and renders reports. Stream JSON, block CSV, refinement requests, and reports are bound by SHA-256.

Table 4. Code and data inventory

Artifact	SHA-256
blocks	f08637095a7084976d9dd58c531f274b12a9b99123e47a43709c2b1c5c c65ca1
stream_set	b9c4115d35f093915e21753b039fbce825c5a7b292b47e0f50b30ef562 3183d4

Hashes bind the report to frozen inputs.