

# LAGRANGE RK4 EXAMPLE

;X acceleration;X acceleration

pro visorbit, inx,iny

nx=n\_elements(inx)

window,xsize=1000,ysize=1000

plot,inx,iny,xrange=[.2,1],yrange=[-.5,.5],ystyle=1,xstyle=1,psym=3

FOR i=0,nx-1 DO BEGIN

oplot,inx[i],iny[i],psym=3,thick=4

wait=.1

ENDFOR

END

FUNCTION xddot, vy,x,u,r1,r2

term1=2.0\*vy+x

term2=-1.0\*u\*(x-1.0+u)/r1^3.

term3=-1.0\*(1.0-u)\*(x+u)/r2^3.

return,term1+term2+term3

END

;Y acceleration

FUNCTION yddot,vx,y,u,r1,r2

term1=y-2.0\*vx

term2=-1.0\*u\*y/r1^3.

term3=-1.0\*(1.0-u)\*y/r2^3.

return, term1+term2+term3

END

```

FUNCTION rad1,x,y,u
return, sqrt((x-1.0+u)^2.+y^2.)
END

```

```

FUNCTION rad2,x,y,u
return, sqrt((x+u)^2.+y^2.)
END

```

```

FUNCTION lpts,a,u
;calculate lagrange points as a function of mu where mu = m1 and m1 < m2
IF a eq 1 THEN BEGIN
alpha=(u/(3d0*(1.0-u)))^(1./3.)
r2=alpha-alpha^2/3d0-alpha^3./9d0-23d0*alpha^3./81d0
return, [(1d0-u)-r2,0d0]
ENDIF

```

```

IF a eq 2 THEN BEGIN
alpha=(u/(3d0*(1.0-u)))^(1./3.)
r2=alpha+alpha^2/3d0-alpha^3./9d0-31d0*alpha^3./81d0
return,[r2+(1d0-u),0d0]
ENDIF

```

```

IF a eq 3 THEN BEGIN
rat=u/(1d0-u)
b=-7d0*rat/12d0+7d0*rat^2./12d0-13223d0*rat^3./20736

```

```
r2=2d0+b
```

```
return,[(1d0-u)-r2,0]
```

```
ENDIF
```

```
IF a eq 4 THEN BEGIN
```

```
Return, [.5d0-u,sqrt(3d0)/2.]
```

```
ENDIF
```

```
IF a eq 5 THEN BEGIN
```

```
Return, [.5d0-u,-1.0*sqrt(3d0)/2.]
```

```
ENDIF
```

```
END
```

```
PRO lagrange,sep,msmall,mbig, values
```

```
;a is separation between m1 and m2 in m
```

```
a=1.49598d11
```

```
a=sep*a
```

```
ms=1.9891d30
```

```
;m2>m1 and on negative x for this formulation to work
```

```
m1=msmall*ms
```

```
m2=mbig*ms
```

```
;radial separation from CM_system
```

```
m=m2/(m1+m2)
```

```
r1=m*a
```

```
r2=(m-1.0)*a
```

```
bounds=[r1,r2]
```

```
kmin=min(bounds)*1.5
```

```

kmax=max(bounds)*1.5
kmin=-2.0*a
kmax=2.0*a
;gravitational constant
G=-6.67428d-11
norm=G*(m1+m2)/a
;array size-keep this even so we kill zeros
n=1000
;make kx and ky and r array where r is distance from center of mass
k = dindgen(n)*(kmax-kmin)/(n-1.)+kmin
kx =k#make_array(n,value=1.0)
ky =make_array(n,value=1.0)#k
kmag2xy = kx*kx+ky*ky
r=sqrt(kmag2xy)
;need s1,s2 = |r-r1|, |r-r2|
;s2=sqrt((kx-r2)^2.+ky^2.)
;s1=sqrt((kx-r1)^2.+ky^2.)
s2=sqrt(r2^2.+r^2.0-2.0*r2*kx)
s1=sqrt(r1^2.+r^2.0-2.0*r1*kx)
pot1= G*m1/s1
pot2= G*m2/s2
pot3= G*.5*(m1+m2)*r^2./a^3.
potential=pot1+pot2+pot3
potnorm=potential/(-1.0*norm)
dx=abs(kx[2]-kx[1])
dy=dx
grad_x=(SHIFT(potential, -1, 0) - SHIFT(potential, 1, 0))/(2 * dx)

```

```

grad_y=(SHIFT(potential, 0, -1) - SHIFT(potential, 0, 1))/(2 * dy)

values =
{r1:r1,r2:r2,kx:kx,ky:ky,kmag2xy:kmag2xy,pot:potential,potnorm:potnorm,m1:m1,
m2:m2, a:a, grad_x:grad_x, grad_y:grad_y}

```

```

END

```

```

PRO l4orbit, t,x,y
;returns x and y orbit of a particle librating about l4
x=(3.54d-4)*sin(.268*t)-(9.85d-5)*sin(.963*t)
y=(6.23d-5)*cos(.268*t)-(4.86d-5)*cos(.963*t)
END

```

```

PRO orbits,x,y,vx,vy,mu,dt,tstop,outarray
;initialize arrays
;l4/l5 y=+-sqrt(3.0)/2. x=.5-mu

```

```

i=0L
dt=double(dt)
t=dt
tarray=t
mu=double(mu)
x0=double(x)
y0=double(y)
vx0=double(vx)
vy0=double(vy)

```

```

xarray=x0
yarray=y0
vxarray=vx0
vyarray=vy0
parray=-mu/rad1(x0,y0,mu)^2.-(1.0-mu)/rad2(x0,y0,mu)^2. -.5*(x0^2.+y0^2.)

```

```

;m1<m2
;mu = M1
;1-mu = M2
;M2+M1=1
;r1=1-mu
;r2=-mu
;G=omega=1

```

```

;MAIN INTEGRATION LOOP

```

```

  WHILE t lt tstop DO BEGIN

```

```

    ;print,"At t = ", t," x0 = ",x0," y0 = ",y0," vx0 = ",vx0," vy0 = ",vy0

```

```

    ;UPDATE x,y,vx,vy - FIRST pass - x1=vx0*dt, v1=a0*dt

```

```

      kx1=vx0*dt ;x1

```

```

      ky1=vy0*dt ;y1

```

```

      kvx1=xddot(vy0,x0,mu,rad1(x0,y0,mu),rad2(x0,y0,mu))*dt ;vx1

```

```

      kvy1=yddot(vx0,y0,mu,rad1(x0,y0,mu),rad2(x0,y0,mu))*dt ;vy1

```

```

    ;print,"At t = ", t," kx1 = ",kx1," ky1 = ",ky1," kvx1 = ",kvx1," kvy1 = ",kvy1

```

;UPDATE x,y,vx,vy - SECOND pass, use at  $x_0+k_1/2$  so new x is  $x_0+v_{x0}dt$ , new v is  $v_{x0}+a*dt$

$kx_2=(v_{x0}+k_{vx1}/2.)*dt$ ;x2

$ky_2=(v_{y0}+k_{vy1}/2.)*dt$ ;y2

$k_{vx2}=xddot(v_{y0}+k_{vy1}/2.,x_0+kx_1/2.,\mu,rad1(x_0+kx_1/2.,y_0+ky_1/2.,\mu),rad2(x_0+kx_1/2.,y_0+ky_1/2.,\mu))*dt$ ;vx2

$k_{vy2}=yddot(v_{x0}+k_{vx1}/2.,y_0+ky_1/2.,\mu,rad1(x_0+kx_1/2.,y_0+ky_1/2.,\mu),rad2(x_0+kx_1/2.,y_0+ky_1/2.,\mu))*dt$ ;vy2

;print,"At t = ",t," kx2 = ",kx2," ky2 = ",ky2," kvx2 = ",kvx2," kvy2 = ",kvy2

;UPDATE x,y,vx,vy - THIRD pass

$kx_3=(v_{x0}+k_{vx2}/2.)*dt$ ;x3

$ky_3=(v_{y0}+k_{vy2}/2.)*dt$ ;y3

$k_{vx3}=xddot(v_{y0}+k_{vy2}/2.,x_0+kx_2/2.,\mu,rad1(x_0+kx_2/2.,y_0+ky_2/2.,\mu),rad2(x_0+kx_2/2.,y_0+ky_2/2.,\mu))*dt$ ;vx3

$k_{vy3}=yddot(v_{x0}+k_{vx2}/2.,y_0+ky_2/2.,\mu,rad1(x_0+kx_2/2.,y_0+ky_2/2.,\mu),rad2(x_0+kx_2/2.,y_0+ky_2/2.,\mu))*dt$ ;vy3

;print,"At t = ",t," kx3 = ",kx3," ky3 = ",ky3," kvx3 = ",kvx3," kvy3 = ",kvy3

;UPDATE x,y,vx,vy - FINAL pass

$kx_4=(v_{x0}+k_{vx3})*dt$ ;x4

$ky_4=(v_{y0}+k_{vy3})*dt$ ;y4

$k_{vx4}=xddot(v_{y0}+k_{vy3},x_0+kx_3,\mu,rad1(x_0+kx_3,y_0+ky_3,\mu),rad2(x_0+kx_3,y_0+ky_3,\mu))*dt$ ;vx4

$k_{vy4}=yddot(v_{x0}+k_{vx3},y_0+ky_3,\mu,rad1(x_0+kx_3,y_0+ky_3,\mu),rad2(x_0+kx_3,y_0+ky_3,\mu))*dt$ ;vy4

```

;print,"At t = ", t," kx4 = ",kx4," ky4 = ",ky4," kvx4 = ",kvx4," kvy4 = ",kvy4

;UPDATE x0,y0, vx0, vy0 to xn,yn,vxn,vyn
  xn = x0+(kx1+2.0*(kx2+kx3)+kx4)/6.
  yn = y0+(ky1+2.0*(ky2+ky3)+ky4)/6.
  vxn = vx0+(kvx1+2.0*(kvx2+kvx3)+kvx4)/6.
  vyn =vy0+(kvy1+2.0*(kvy2+kvy3)+kvy4)/6.

;print,"At t = ", t," xn = ",xn," yn = ",yn," vxn = ",vxn," vyn = ",vyn
  xarray=[[xarray],[xn]]
  yarray=[[yarray],[yn]]
  vxarray=[[vxarray],[vxn]]
  vyarray=[[vyarray],[vyn]]
  x0=xn
  y0=yn
  vx0=vxn
  vy0=vyn
  print, "T = ",t,"  integration loop ", i
  i +=1
  t+=dt
  tarray=[[tarray],[t]]
  potential=-mu/rad1(x0,y0,mu)^2.-(1.0-mu)/rad2(x0,y0,mu)^2. -.5*(x0^2.+y0^2.)
  parray=[[parray],[potential]]
ENDWHILE
  energy=.5*vxarray^2+.5*vyarray^2.
  jacobi=-2.0*(parray+energy)

```



```
momentum= vxarray+vyarray
```

```
angmomentum=vyarray*xarray-vxarray*yarray
```

```
outarray = {ke:energy,pot:parray,jacobi:jacobi, p:momentum, rp:angmomentum,  
x:xarray, y:yarray,vx:vxarray, vy:vyarray, t:tarray}
```

```
END
```

```
PRO lpoints
```

```
END
```

```
pro visorbit, inx,iny
```

```
nx=n_elements(inx)
```

```
window,xsize=1000,ysize=1000
```

```
plot,inx,iny,xrange=[.2,1],yrange=[-.5,.5],ystyle=1,xstyle=1,psym=3
```

```
FOR i=0,nx-1 DO BEGIN
```

```
  oplot,inx[i],iny[i],psym=3,thick=4
```

```
  wait=.1
```

```
ENDFOR
```

```
END
```

```
FUNCTION xddot, vy,x,u,r1,r2
```

```
term1=2.0*vy+x
```

```
term2=-1.0*u*(x-1.0+u)/r1^3.
```

```
term3=-1.0*(1.0-u)*(x+u)/r2^3.
```

```
return,term1+term2+term3
```

END

;Y acceleration

FUNCTION yddot,vx,y,u,r1,r2

term1=y-2.0\*vx

term2=-1.0\*u\*y/r1^3.

term3=-1.0\*(1.0-u)\*y/r2^3.

return, term1+term2+term3

END

FUNCTION rad1,x,y,u

return, sqrt((x-1.0+u)^2.+y^2.)

END

FUNCTION rad2,x,y,u

return, sqrt((x+u)^2.+y^2.)

END

FUNCTION lpts,a,u

;calculate lagrange points as a function of mu where mu = m1 and m1 < m2

IF a eq 1 THEN BEGIN

alpha=(u/(3d0\*(1.0-u)))^(1./3.)

r2=alpha-alpha^2/3d0-alpha^3./9d0-23d0\*alpha^3./81d0

return, [(1d0-u)-r2,0d0]

ENDIF

IF a eq 2 THEN BEGIN

$\alpha = (u / (3d0 * (1.0 - u)))^{(1./3.)}$

$r2 = \alpha + \alpha^2 / 3d0 - \alpha^3 / 9d0 - 31d0 * \alpha^3 / 81d0$

return,[r2+(1d0-u),0d0]

ENDIF

IF a eq 3 THEN BEGIN

$rat = u / (1d0 - u)$

$b = -7d0 * rat / 12d0 + 7d0 * rat^2 / 12d0 - 13223d0 * rat^3 / 20736$

$r2 = 2d0 + b$

return,[(1d0-u)-r2,0]

ENDIF

IF a eq 4 THEN BEGIN

Return, [.5d0-u,sqrt(3d0)/2.]

ENDIF

IF a eq 5 THEN BEGIN

Return, [.5d0-u,-1.0\*sqrt(3d0)/2.]

ENDIF

END

PRO lagrange,sep,msmall,mbig, values

;a is separation between m1 and m2 in m

a=1.49598d11

a=sep\*a

ms=1.9891d30

```

;m2>m1 and on negative x for this formulation to work
m1=m*small*ms
m2=m*big*ms
;radial separation from CM_system
m=m2/(m1+m2)
r1=m*a
r2=(m-1.0)*a
bounds=[r1,r2]
kmin=min(bounds)*1.5
kmax=max(bounds)*1.5
kmin=-2.0*a
kmax=2.0*a
;gravitational constant
G=-6.67428d-11
norm=G*(m1+m2)/a
;array size-keep this even so we kill zeros
n=1000
;make kx and ky and r array where r is distance from center of mass
k = dindgen(n)*(kmax-kmin)/(n-1.)+kmin
kx =k#make_array(n,value=1.0)
ky =make_array(n,value=1.0)#k
kmag2xy = kx*kx+ky*ky
r=sqrt(kmag2xy)
;need s1,s2 = |r-r1|, |r-r2|
;s2=sqrt((kx-r2)^2.+ky^2.)
;s1=sqrt((kx-r1)^2.+ky^2.)
s2=sqrt(r2^2.+r^2.-2.0*r2*kx)

```

```

s1=sqrt(r1^2.+r^2.0-2.0*r1*kx)
pot1= G*m1/s1
pot2= G*m2/s2
pot3= G*.5*(m1+m2)*r^2./a^3.
potential=pot1+pot2+pot3
potnorm=potential/(-1.0*norm)
dx=abs(kx[2]-kx[1])
dy=dx
grad_x=(SHIFT(potential, -1, 0) - SHIFT(potential, 1, 0))/(2 * dx)
grad_y=(SHIFT(potential, 0, -1) - SHIFT(potential, 0, 1))/(2 * dy)

values =
{r1:r1,r2:r2,kx:kx,ky:ky,kmag2xy:kmag2xy,pot:potential,potnorm:potnorm,m1:m1,
m2:m2, a:a, grad_x:grad_x, grad_y:grad_y}

```

END

```

PRO l4orbit, t,x,y
;returns x and y orbit of a particle librating about l4
x=(3.54d-4)*sin(.268*t)-(9.85d-5)*sin(.963*t)
y=(6.23d-5)*cos(.268*t)-(4.86d-5)*cos(.963*t)
END

```

```

PRO orbits,x,y,vx,vy,mu,dt,tstop,outarray
;initialize arrays
;l4/l5 y=+-sqrt(3.0)/2. x=.5-mu

i=0L
dt=double(dt)

```

```
t=dt
```

```
tarray=t
```

```
mu=double(mu)
```

```
x0=double(x)
```

```
y0=double(y)
```

```
vx0=double(vx)
```

```
vy0=double(vy)
```

```
xarray=x0
```

```
yarray=y0
```

```
vxarray=vx0
```

```
vyarray=vy0
```

```
parray=-mu/rad1(x0,y0,mu)^2.-(1.0-mu)/rad2(x0,y0,mu)^2. -.5*(x0^2.+y0^2.)
```

```
;m1<m2
```

```
;mu = M1
```

```
;1-mu = M2
```

```
;M2+M1=1
```

```
;r1=1-mu
```

```
;r2=-mu
```

```
;G=omega=1
```

```
;MAIN INTEGRATION LOOP
```

```
WHILE t lt tstop DO BEGIN
```

```

;print,"At t = ", t," x0 = ",x0," y0 = ",y0," vx0 = ",vx0," vy0 = ",vy0
;UPDATE x,y,vx,vy - FIRST pass - x1=vx0*dt, v1=a0*dt

kx1=vx0*dt ;x1
ky1=vy0*dt ;y1
kvx1=xddot(vy0,x0,mu,rad1(x0,y0,mu),rad2(x0,y0,mu))*dt ;vx1
kvy1=yddot(vx0,y0,mu,rad1(x0,y0,mu),rad2(x0,y0,mu))*dt ;vy1
;print,"At t = ", t," kx1 = ",kx1," ky1 = ",ky1," kvx1 = ",kvx1," kvy1 = ",kvy1

;UPDATE x,y,vx,vy - SECOND pass, use at x0+k1/2 so new x is x0+vx0dt, new v is
vx0+a*dt

kx2=(vx0+kvx1/2.)*dt ;x2
ky2=(vy0+kvy1/2.)*dt ;y2

kvx2=xddot(vy0+kvy1/2.,x0+kx1/2.,mu,rad1(x0+kx1/2.,y0+ky1/2.,mu),rad2(x0+kx
1/2.,y0+ky1/2.,mu))*dt ;vx2

kvy2=yddot(vx0+kvx1/2.,y0+ky1/2.,mu,rad1(x0+kx1/2.,y0+ky1/2.,mu),rad2(x0+kx
1/2.,y0+ky1/2.,mu))*dt ;vy2

;print,"At t = ", t," kx2 = ",kx2," ky2 = ",ky2," kvx2 = ",kvx2," kvy2 = ",kvy2

;UPDATE x,y,vx,vy - THIRD pass

kx3=(vx0+kvx2/2.)*dt ;x3
ky3=(vy0+kvy2/2.)*dt ;y3

kvx3=xddot(vy0+kvy2/2.,x0+kx2/2.,mu,rad1(x0+kx2/2.,y0+ky2/2.,mu),rad2(x0+kx
2/2.,y0+ky2/2.,mu))*dt ;vx3

kvy3=yddot(vx0+kvx2/2.,y0+ky2/2.,mu,rad1(x0+kx2/2.,y0+ky2/2.,mu),rad2(x0+kx
2/2.,y0+ky2/2.,mu))*dt ;vy3

;print,"At t = ", t," kx3 = ",kx3," ky3 = ",ky3," kvx3 = ",kvx3," kvy3 = ",kvy3

```

```
;UPDATE x,y,vx,vy - FINAL pass
```

```
    kx4=(vx0+kvx3)*dt ;x4
```

```
    ky4=(vy0+kvy3)*dt ;y4
```

```
    kvx4=xddot(vy0+kvy3,x0+kx3,mu,rad1(x0+kx3,y0+ky3,mu),rad2(x0+kx3,y0+ky3,mu))*dt ;vx4
```

```
    kvy4=yddot(vx0+kvx3,y0+ky3,mu,rad1(x0+kx3,y0+ky3,mu),rad2(x0+kx3,y0+ky3,mu))*dt ;vy4
```

```
    ;print,"At t = ", t," kx4 = ",kx4," ky4 = ",ky4," kvx4 = ",kvx4," kvy4 = ",kvy4
```

```
;UPDATE x0,y0, vx0, vy0 to xn,yn,vxn,vyn
```

```
    xn = x0+(kx1+2.0*(kx2+kx3)+kx4)/6.
```

```
    yn = y0+(ky1+2.0*(ky2+ky3)+ky4)/6.
```

```
    vxn = vx0+(kvx1+2.0*(kvx2+kvx3)+kvx4)/6.
```

```
    vyn =vy0+(kvy1+2.0*(kvy2+kvy3)+kvy4)/6.
```

```
;print,"At t = ", t," xn = ",xn," yn = ",yn," vxn = ",vxn," vyn = ",vyn
```

```
    xarray=[[xarray],[xn]]
```

```
    yarray=[[yarray],[yn]]
```

```
    vxarray=[[vxarray],[vxn]]
```

```
    vyarray=[[vyarray],[vyn]]
```

```
    x0=xn
```

```
    y0=yn
```

```
    vx0=vxn
```

```
    vy0=vyn
```

```
    print, "T = ",t," integration loop ", i
```



```

i +=1
t+=dt
tarray=[[tarray],[t]]
potential=-mu/rad1(x0,y0,mu)^2.-(1.0-mu)/rad2(x0,y0,mu)^2. -.5*(x0^2.+y0^2.)
parray=[[parray],[potential]]
ENDWHILE

energy=.5*vxarray^2+.5*vyarray^2.
jacobi=-2.0*(parray+energy)
momentum= vxarray+vyarray
angmomentum=vyarray*xarray-vxarray*yarray

outarray = {ke:energy,pot:parray,jacobi:jacobi, p:momentum, rp:angmomentum,
x:xarray, y:yarray,vx:vxarray, vy:vyarray, t:tarray}

END

PRO lpoints

END

```

## PENDULUM WITH SPRING EXAMPLE

```

FUNCTION lddot, l, theta, thetadot, k, m, b
term1 = (l+b)*thetadot^2.
term2=9.8*cos(theta)
term3 = -k*l/m

```

```
return, term1 + term2 + term3
```

```
END
```

```
FUNCTION thetaddot,theta,thetadot,l,ldot,b
```

```
term1=-1.*(9.8*sin(theta)+2*ldot*thetadot)
```

```
term2=l+b
```

```
return, term1/term2
```

```
END
```

```
PRO ch7plot2,l,theta,ldot,thetadot,kin,min,bin,dt,tstop,outarray
```

```
;initialize arrays
```

```
i=0L
```

```
dt=double(dt)
```

```
t=dt
```

```
tarray=t
```

```
k=double(kin)
```

```
m=double(min)
```

```
b=double(bin)
```

```
x0=double(l)
```

```
y0=double(theta)
```

```
vx0=double(ldot)
```

```
vy0=double(thetadot)
```

```
xarray=x0
yarray=y0
vxarray=vx0
vyarray=vy0
```

```
;MAIN INTEGRATION LOOP
```

```
WHILE t lt tstop DO BEGIN
```

```
;print,"At t = ", t," x0 = ",x0," y0 = ",y0," vx0 = ",vx0," vy0 = ",vy0
```

```
;UPDATE x,y,vx,vy - FIRST pass -  $x_1 = vx_0 * dt$ ,  $v_1 = a_0 * dt$ 
```

```
   $kx_1 = vx_0 * dt$  ;x1
```

```
   $ky_1 = vy_0 * dt$  ;y1
```

```
   $kvx_1 = lddot(x_0, y_0, vy_0, k, m, b) * dt$  ;vx1
```

```
   $kvy_1 = thetaddot(y_0, vy_0, x_0, vx_0, b) * dt$  ;vy1
```

```
;print,"At t = ", t,"  $kx_1 =$ ", $kx_1$ ,"  $ky_1 =$ ", $ky_1$ ,"  $kvx_1 =$ ", $kvx_1$ ,"  $kvy_1 =$ ", $kvy_1$ 
```

```
;UPDATE x,y,vx,vy - SECOND pass, use at  $x_0 + k_1/2$  so new x is  $x_0 + vx_0 dt$ , new v is  $vx_0 + a * dt$ 
```

```
   $kx_2 = (vx_0 + kvx_1/2.) * dt$  ;x2
```

```
   $ky_2 = (vy_0 + kvy_1/2.) * dt$  ;y2
```

```
   $kvx_2 = lddot(x_0 + kx_1/2., y_0 + ky_1/2., vy_0 + kvy_1/2., k, m, b) * dt$  ;vx2
```

```
   $kvy_2 = thetaddot(y_0 + ky_1/2., vy_0 + kvy_1/2., x_0 + kx_1/2., vx_0 + kvx_1/2., b) * dt$  ;vy2
```

```
;print,"At t = ", t,"  $kx_2 =$ ", $kx_2$ ,"  $ky_2 =$ ", $ky_2$ ,"  $kvx_2 =$ ", $kvx_2$ ,"  $kvy_2 =$ ", $kvy_2$ 
```

```
;UPDATE x,y,vx,vy - THIRD pass
```

```
   $kx_3 = (vx_0 + kvx_2/2.) * dt$  ;x3
```

```
   $ky_3 = (vy_0 + kvy_2/2.) * dt$  ;y3
```

```

kvx3=lddot(x0+kx2/2.,y0+ky2/2.,vy0+kvy2/2.,k,m,b)*dt ;vx3
kvy3=thetaddot(y0+ky2/2.,vy0+kvy2/2.,x0+kx2/2.,vx0+kvx2/2.,b)*dt;vy3
;print,"At t = ", t," kx3 = ",kx3," ky3 = ",ky3," kvx3 = ",kvx3," kvy3 = ",kvy3

;UPDATE x,y,vx,vy - FINAL pass
kx4=(vx0+kvx3)*dt ;x4
ky4=(vy0+kvy3)*dt ;y4
kvx4=lddot(x0+kx3,y0+ky3,vy0+kvy3,k,m,b)*dt ;vx4
kvy4=thetaddot(y0+ky3,vy0+kvy3,x0+kx3,vx0+kvx3,b)*dt;vy3
;print,"At t = ", t," kx4 = ",kx4," ky4 = ",ky4," kvx4 = ",kvx4," kvy4 = ",kvy4

;UPDATE x0,y0, vx0, vy0 to xn,yn,vxn,vyn
xn = x0+(kx1+2.0*(kx2+kx3)+kx4)/6.
yn = y0+(ky1+2.0*(ky2+ky3)+ky4)/6.
vxn = vx0+(kvx1+2.0*(kvx2+kvx3)+kvx4)/6.
vyn =vy0+(kvy1+2.0*(kvy2+kvy3)+kvy4)/6.

;print,"At t = ", t," xn = ",xn," yn = ",yn," vxn = ",vxn," vyn = ",vyn
xarray=[[xarray],[xn]]
yarray=[[yarray],[yn]]
vxarray=[[vxarray],[vxn]]
vyarray=[[vyarray],[vyn]]
x0=xn
y0=yn
vx0=vxn
vy0=vyn
print, "T = ",t," integration loop ", i

```

```

i +=1
t+=dt
tarray=[[tarray],[t]]
ENDWHILE
energy=.5*vxarray^2+.5*vyarray^2.

outarray = {ke:energy, x:xarray, y:yarray,vx:vxarray, vy:vyarray, t:tarray}

END

```

## SPINNING CYLINDER WITH BEAD EXAMPLE

```

FUNCTION rddot, r,rdot,theta,omega,cons
term1=r*((theta+omega)^2.-2.*9.8*cons)
term2=rdot^2.*4*r*cons^2
term3=1.+4.*cons^2*r^2
return,(term1-term2)/term3
END

FUNCTION thetaddot,thetadot,r,rdot,omega
term1=-2.*rdot*(thetadot+omega)
return, term1/(r+.00001)
END

```

```
PRO ch7plot,r,theta,rdot,thetadot,c,omegaforce,dt,tstop,outarray
```

```
;initialize arrays
```

```
i=0L
```

```
dt=double(dt)
```

```
t=dt
```

```
tarray=t
```

```
omega=double(omegaforce)
```

```
x0=double(r)
```

```
y0=double(theta)
```

```
vx0=double(rdot)
```

```
vy0=double(thetadot)
```

```
cons=double(c)
```

```
xarray=x0
```

```
yarray=y0
```

```
vxarray=vx0
```

```
vyarray=vy0
```

```
;MAIN INTEGRATION LOOP
```

```
WHILE t lt tstop DO BEGIN
```

```
;print,"At t = ", t," x0 = ",x0," y0 = ",y0," vx0 = ",vx0," vy0 = ",vy0
```

;UPDATE x,y,vx,vy - FIRST pass -  $x_1 = vx_0 * dt$ ,  $v_1 = a_0 * dt$

$kx_1 = vx_0 * dt$  ;x1

$ky_1 = vy_0 * dt$  ;y1

$kvx_1 = rddot(x_0, vx_0, y_0, \omega, cons) * dt$  ;vx1

$kvy_1 = thetaddot(vy_0, x_0, vx_0, \omega) * dt$  ;vy1

;print,"At t = ", t," kx1 = ",kx1," ky1 = ",ky1," kvx1 = ",kvx1," kvy1 = ",kvy1

;UPDATE x,y,vx,vy - SECOND pass, use at  $x_0 + k_1/2$  so new x is  $x_0 + vx_0 dt$ , new v is  $vx_0 + a * dt$

$kx_2 = (vx_0 + kvx_1/2.) * dt$  ;x2

$ky_2 = (vy_0 + kvy_1/2.) * dt$  ;y2

$kvx_2 = rddot(x_0 + kx_1/2., vx_0 + kvx_1/2., y_0 + ky_1/2., \omega, cons) * dt$  ;vx2

$kvy_2 = thetaddot(vy_0 + kvy_1/2., x_0 + kx_1/2., vx_0 + kvx_1/2., \omega) * dt$  ;vy2

;print,"At t = ", t," kx2 = ",kx2," ky2 = ",ky2," kvx2 = ",kvx2," kvy2 = ",kvy2

;UPDATE x,y,vx,vy - THIRD pass

$kx_3 = (vx_0 + kvx_2/2.) * dt$  ;x3

$ky_3 = (vy_0 + kvy_2/2.) * dt$  ;y3

$kvx_3 = rddot(x_0 + kx_2/2., vx_0 + kvx_2/2., y_0 + ky_2/2., \omega, cons) * dt$  ;vx3

$kvy_3 = thetaddot(vy_0 + kvy_2/2., x_0 + kx_2/2., vx_0 + kvx_2/2., \omega) * dt$  ;vy3

;print,"At t = ", t," kx3 = ",kx3," ky3 = ",ky3," kvx3 = ",kvx3," kvy3 = ",kvy3

;UPDATE x,y,vx,vy - FINAL pass

$kx_4 = (vx_0 + kvx_3) * dt$  ;x4

$ky_4 = (vy_0 + kvy_3) * dt$  ;y4

$kvx_4 = rddot(x_0 + kx_3, vx_0 + kvx_3, y_0 + ky_3, \omega, cons) * dt$  ;vx4

$kvy_4 = thetaddot(vy_0 + kvy_3, x_0 + kx_3, vx_0 + kvx_3, \omega) * dt$  ;vy4

```

;print,"At t = ", t," kx4 = ",kx4," ky4 = ",ky4," kvx4 = ",kvx4," kvy4 = ",kvy4

;UPDATE x0,y0, vx0, vy0 to xn,yn,vxn,vyn
  xn = x0+(kx1+2.0*(kx2+kx3)+kx4)/6.
  yn = y0+(ky1+2.0*(ky2+ky3)+ky4)/6.
  vxn = vx0+(kvx1+2.0*(kvx2+kvx3)+kvx4)/6.
  vyn =vy0+(kvy1+2.0*(kvy2+kvy3)+kvy4)/6.

;print,"At t = ", t," xn = ",xn," yn = ",yn," vxn = ",vxn," vyn = ",vyn
  xarray=[[xarray],[xn]]
  yarray=[[yarray],[yn]]
  vxarray=[[vxarray],[vxn]]
  vyarray=[[vyarray],[vyn]]
  x0=xn
  y0=yn
  vx0=vxn
  vy0=vyn
  print, "T = ",t,"  integration loop ", i
  i +=1
  t+=dt
  tarray=[[tarray],[t]]
ENDWHILE

  energy=.5*vxarray^2+.5*vyarray^2.

  outarray = {ke:energy, x:xarray, y:yarray,vx:vxarray, vy:vyarray, t:tarray}

END

```