


11.1 Fourier Series

$$f(x) = \begin{cases} -k & \text{if } -\pi < x < 0 \\ k & \text{if } 0 < x < \pi \end{cases} \quad \text{and} \quad f(x + 2\pi) = f(x).$$



$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$






Fig. 261. First three partial sums of the corresponding Fourier series

Orthogonality of the Trigonometric System (3)

The trigonometric system (3) is orthogonal on the interval $-\pi \leq x \leq \pi$ (hence also on $0 \leq x \leq 2\pi$ or any other interval of length 2π because of periodicity); that is, the integral of the product of any two functions in (3) over that interval is 0, so that for any integers n and m ,

$$(a) \quad \int_{-\pi}^{\pi} \cos nx \cos mx \, dx = 0 \quad (n \neq m)$$

$$(b) \quad \int_{-\pi}^{\pi} \sin nx \sin mx \, dx = 0 \quad (n \neq m)$$


$$(c) \quad \int_{-\pi}^{\pi} \sin nx \cos mx \, dx = 0 \quad (n \neq m \text{ or } n = m).$$

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \, dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx$$



Periodic Rectangular Wave (Fig. 260)

Find the Fourier coefficients of the periodic function $f(x)$ in Fig. 260. The formula is

$$(7) \quad f(x) = \begin{cases} -k & \text{if } -\pi < x < 0 \\ k & \text{if } 0 < x < \pi \end{cases} \quad \text{and} \quad f(x + 2\pi) = f(x).$$

Functions of this kind occur as external forces acting on mechanical systems, electromotive forces in electric circuits, etc. (The value of $f(x)$ at a single point does not affect the integral; hence we can leave $f(x)$ undefined at $x = 0$ and $x = \pm\pi$.)





Fig. 260. Given function $f(x)$ (Periodic rectangular wave)

EXAMPLE 1

$$f(x) = \begin{cases} -k & \text{if } -\pi < x < 0 \\ k & \text{if } 0 < x < \pi \end{cases} \quad \text{and} \quad f(x + 2\pi) = f(x).$$



$$\begin{aligned} a_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 (-k) \cos nx \, dx + \int_0^{\pi} k \cos nx \, dx \right] \\ &= \frac{1}{\pi} \left[-k \frac{\sin nx}{n} \Big|_{-\pi}^0 + k \frac{\sin nx}{n} \Big|_0^{\pi} \right] = 0 \end{aligned}$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) \, dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx \, dx$$


$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx$$

$$\begin{aligned} b_n &= \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx \, dx = \frac{1}{\pi} \left[\int_{-\pi}^0 (-k) \sin nx \, dx + \int_0^{\pi} k \sin nx \, dx \right] = \frac{1}{\pi} \left[k \frac{\cos nx}{n} \Big|_{-\pi}^0 - k \frac{\cos nx}{n} \Big|_0^{\pi} \right] \\ &= \frac{k}{n\pi} [\cos 0 - \cos(-n\pi) - \cos n\pi + \cos 0] = \frac{2k}{n\pi} (1 - \cos n\pi). \end{aligned}$$

Since the a_n are zero, the Fourier series of $f(x)$ is

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx) = \frac{4k}{\pi} \left(\sin x + \frac{1}{3} \sin 3x + \frac{1}{5} \sin 5x + \cdots \right).$$

$$\cos n\pi = \begin{cases} -1 & \text{for odd } n, \\ 1 & \text{for even } n, \end{cases} \quad \text{and thus} \quad 1 - \cos n\pi = \begin{cases} 2 & \text{for odd } n, \\ 0 & \text{for even } n. \end{cases}$$



Since the a_n are zero, the Fourier series of $f(x)$ is

$$(8) \quad \frac{4k}{\pi} (\sin x + \frac{1}{3} \sin 3x + \frac{1}{5} \sin 5x + \cdots).$$

The partial sums are

$$S_1 = \frac{4k}{\pi} \sin x, \quad S_2 = \frac{4k}{\pi} \left(\sin x + \frac{1}{3} \sin 3x \right), \quad \text{etc.}$$

Their graphs in Fig. 261 seem to indicate that the series is convergent and has the sum $f(x)$, the given function. We notice that at $x = 0$ and $x = \pi$, the points of discontinuity of $f(x)$, all partial sums have the value zero, the arithmetic mean of the limits $-k$ and k of our function, at these points. This is typical.

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$




Fig. 261. First three partial sums of the corresponding Fourier series

Representation by a Fourier Series

Let $f(x)$ be periodic with period 2π and piecewise continuous (see Sec. 6.1) in the interval $-\pi \leq x \leq \pi$. Furthermore, let $f(x)$ have a left-hand derivative and a right-hand derivative at each point of that interval. Then the Fourier series (5) of $f(x)$ [with coefficients (6)] converges. Its sum is $f(x)$, except at points x_0 where $f(x)$ is discontinuous. There the sum of the series is the average of the left- and right-hand limits² of $f(x)$ at x_0 .

$$f(x) = a_0 + \sum_{n=1}^{\infty} (a_n \cos nx + b_n \sin nx)$$

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos nx dx$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin nx dx$$




Fig. 261. First three partial sums of the corresponding Fourier series

THEOREM 2

Ch11.1 Fourier Series: In Class Examples



Helpful Formulas/Results

① $\cos(-x) = \cos(x)$

② $\sin(-x) = -\sin(x)$

③ For $A\sin(Bx)$ & $A\cos(Bx)$, $A = \text{amplitude}$, $\text{period} = \frac{2\pi}{B}$, $\text{freq} = \frac{B}{2\pi}$

④




Examples

- ① Observe from the graphs of $\sin x$ & $\cos x$ that they are periodic with period 2π . It can also be said that they are periodic with period 4π , since they repeat after 4π units. Thus the fundamental period is 2π , since it is the smallest period.

- ② $\sin(3x)$ has fundamental period $\frac{2\pi}{3}$ (graph repeats after $[0, \frac{2\pi}{3}]$).
 $\cos(2\pi x)$ has fundamental period $\frac{2\pi}{2\pi} = 1$ (graph repeats after $[0, 1]$).

(Note: Answer in back of book for #1 has errors. See if you can correct them.)

$$(3) f(x) = \begin{cases} 0, & -\pi < x < 0 \\ x, & 0 < x < \pi \end{cases}$$



(a) Find a_0 , a_n , and b_n .

$$a_0 = \frac{1}{2\pi} \int_{-\pi}^{\pi} f(x) dx = \frac{1}{2\pi} \int_0^{\pi} x dx = \frac{1}{2\pi} \left(\frac{1}{2} x^2 \right) \Big|_0^{\pi} = \frac{1}{4\pi} [\pi^2 - 0^2] = \frac{\pi}{4}$$

$$a_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \cos(nx) dx = \frac{1}{\pi} \int_0^{\pi} x \cos(nx) dx$$

$\begin{cases} u=x & dv=\cos(nx)dx \\ du=dx & v=\frac{1}{n}\sin(nx) \end{cases}$

$$= \frac{1}{\pi} \left[\frac{x}{n} \sin(nx) \Big|_0^{\pi} - \frac{1}{n} \int_0^{\pi} \sin(nx) dx \right]$$

$$= \frac{1}{\pi} \left[\frac{1}{n^2} \cos(nx) \Big|_0^{\pi} \right]$$

$$= \frac{1}{n^2\pi} [\cos(n\pi) - 1]$$

$$= \begin{cases} 0, & \text{if } n=2k \text{ (even)} \\ -\frac{2}{n^2\pi}, & \text{if } n=2k+1 \text{ (odd)} \end{cases}$$

$$b_n = \frac{1}{\pi} \int_{-\pi}^{\pi} f(x) \sin(nx) dx = \frac{1}{\pi} \int_0^{\pi} x \sin(nx) dx = \frac{1}{\pi} \left[-\frac{x}{n} \cos(nx) \Big|_0^{\pi} + \frac{1}{n} \int_0^{\pi} \cos(nx) dx \right]$$

$\begin{cases} u=x & dv=\sin(nx)dx \\ du=dx & v=-\frac{1}{n}\cos(nx) \end{cases}$

$$= \frac{1}{\pi} \left[-\frac{\pi}{n} \cos(n\pi) + \frac{1}{n^2} \sin(nx) \Big|_0^{\pi} \right]$$

$$= -\frac{1}{n} \cos(n\pi) = -\frac{1}{n} (-1)^n = \frac{(-1)^{n+1}}{n}$$

(b) Find Fourier series for $f(x)$.

$$f(x) = \frac{\pi}{4} - \frac{2}{\pi} \left(\cos(x) + \frac{1}{9} \cos(3x) + \frac{1}{25} \cos(5x) + \dots \right)$$

$$+ \left(\sin(x) - \frac{1}{2} \sin(2x) + \frac{1}{3} \sin(3x) - \frac{1}{4} \sin(4x) + \dots \right)$$

(c) Graph partial sums up to that including $\cos 5x$ and $\sin 5x$.

- Use Maple, Desmos.com or graphing calculator.

Math 360
Ch11.1 Fourier Series

Methods of Applied Math

In-Class Example

