Taylor Larrechea
Dr. Gustafson
MATH 361 HW 2.1

// / x / / #- 1,2,35,8,14,17

Problem 1  $f(x) = \sqrt{x} - \cos(x)$  on [0,1]

use bisection method to find P3

f(0)=-1, f(1)=0.459 : f(0) = f(6) < 0 /

f(0.5) = -0.170: f(0.5) < 0

f(0.75) = 0.134 : f(0.75) > 0

f(0.625) = -0.0203 : f(0.625) < 0

P3(x) = 0.625

f(x)= 3(x+1)(x-1/2)(x-1)=0

a.) [-2,1.5]

		ρ			
- 2	1.5	-0.25 -1.125 -0.6875	- 1	+ 1	+ 1
- ೩	-0.25	-1.125	-1	+ (	- 1
-1.125	-0.25	-0.6875	-1	+1	+ 1

 $P_3(x) = -0.6875$ 

b.) [-1.25, a.5]

a	Ь	ρ	fan	िरक	F(P)
		0.625		<b>+</b> 1	-1
		1.5625		+	+1
0.625	1.5625	1.09375	<b>-</b> l	+1	+1

 $P_3(x) = 1.09379$ 

$$f(x) = x^3 - 7x^2 + 14x - 6$$

b.) [1,3.2]

a	<b> </b>	ρ	flas	fw	F(P)
1	3.2	<b>2</b> .1	+ 1	-1	+1
ર્ગ. ા	3.2	2.65	+ (	-1	+1
2.65	3.2	2.925	<b>+</b> l	-1	+ l
2.925	3.Z	3.0625	+ l	- l	~ 1
2.925	3.0625	<i>2.</i> 99375	+ 1	-1	+ l
2.99375	3.0625	3.028125	+ (	-1	-1
3.023125	3.0625	3.0453125	+1	-(	- (
3.028125	3-0463125	3.63671875	+ (	-1	- 1

 $P_8(x) = 3.03671875$ 

 $f(x) = X^2 - 3$  [1,2]  $\sqrt{3} \approx 1.73205$ 

_ a	_ b	ρ	far	fw	F(P)
1	a	1.5	-1	+1	-1
1.5	a	1.75	-1	+1	+1
1.5	1.75	1.625	-1	+ (	-1
1.625	1.75	1.6875	-1	+1	-1
1.6875	1.75	1.71875	-1	<b>+</b> 1	-1
1.71875	1.75	1.734375	~1	+1	+1
1.71875	1.734375	1.7265625	-1	1+	-1
1.7265625	1.734375	1.73046875	-1	+ l	-1
1.73046875	1.734375	1.73.2421875	-1	+1	+ 1
1.73046875	1.732421875	1.7314453125	-1	+ 1	-1
1.7314453125	1.732421875	1.73193359375	-1	+1	-1
1.73193359375	1.732421875	J.73217773	-1	+1	+1

√3 ≈ 1.7321

## Problem 17

$$f(x) = x^{3} - x - 1 = 0 , \quad [1,2]$$

$$\frac{b - a}{a^{n}} < 10^{-4} - b \quad \frac{b - a}{10^{-4}} < a^{n} \rightarrow J_{010}(a^{n}) > J_{010}(\frac{b - a}{10^{-4}})$$

$$n \cdot J_{010}(a) > J_{010}(\frac{b - a}{10^{-4}}) \rightarrow n > \frac{1}{J_{010}(a)} \cdot J_{010}(\frac{b - a}{10^{-4}})$$

$$n > \frac{1}{J_{010}(a)} \cdot J_{010}(\frac{a - 1}{10^{-4}}) \rightarrow n > 13.2877$$

n ≥ 14 iterations