Fundamental Mechanics: Quiz 9

1 November 2016

Name: Taylor Larrachea

Total:

1.5 /5

Formulae:

$$K = \frac{1}{2}mv^2$$

$$K = \frac{1}{2}mv^2$$
 $W = \vec{\mathbf{F}} \cdot \Delta \vec{\mathbf{r}} = F\Delta r \cos \theta$ $W_{\rm net} = \Delta K$ $g = 9.80 \,\mathrm{m/s^2}$

$$W_{\rm net} = \Delta K$$

$$g = 9.80 \,\mathrm{m/s^2}$$

$$U_{\rm g}=mgy$$

$$U_{\rm g} = mgy$$
 $U_{\rm spring} = \frac{1}{2}k(\Delta s)^2$ $E = K + U_{\rm g} + U_{\rm spring}$ $\Delta E = W_{\rm nc}$


$$E = K + U_{\rm g} + U_{\rm spring}$$

$$\Delta E = W_{\rm nc}$$

$$F_x = -\frac{dU}{dx}$$

A particle with the illustrated total mechanical energy moves subject to the illustrated potential. Indicate the locations at which the speed is a maximum and the force on the particle is zero.

At spots 1.5 and 15 (-0.5) the speeds are maximums because this is where all of the energy is kinetic with no potential.



Where X=0.25 and X=7.25 is where the force is 0 because on of the energy is potential.

F= slope...