

Spectral decomposition, purification and superdense coding

Ryuhei Mori

Tokyo Institute of Technology

Pauli matrices in bracket notation

-

$$Z = \begin{bmatrix} 1 & 0 \\ 0 & -1 \end{bmatrix} = |0\rangle\langle 0| - |1\rangle\langle 1|$$

- $|+\rangle := (|0\rangle + |1\rangle)/\sqrt{2}, \quad |-\rangle := (|0\rangle - |1\rangle)/\sqrt{2}.$

$$X = \begin{bmatrix} 0 & 1 \\ 1 & 0 \end{bmatrix} = |+\rangle\langle +| - |-\rangle\langle -|$$

- $|a\rangle := (|0\rangle + i|1\rangle)/\sqrt{2}, \quad |b\rangle := (|0\rangle - i|1\rangle)/\sqrt{2}.$

$$Y = \begin{bmatrix} 0 & -i \\ i & 0 \end{bmatrix} = |a\rangle\langle a| - |b\rangle\langle b|$$

Spectral decomposition theorem

Definition (Normal operator)

$A \in \mathcal{L}(\mathcal{X})$ is said to be **normal** if $AA^\dagger = A^\dagger A$.

Hermitian matrix ($H^\dagger = H$) and unitary matrix ($UU^\dagger = I$) are normal.

Theorem (Spectral decomposition theorem)

$A \in \mathcal{L}(\mathbb{C}^n)$ is **normal** if and only if there exist orthonormal basis $\{|\psi_j\rangle\}$ of \mathbb{C}^n and complex numbers $\{\lambda_j\}$ such that

$$A = \sum_j \lambda_j |\psi_j\rangle \langle \psi_j|.$$

Any complex matrix has an eigenvalue

For any $A \in \mathcal{L}(\mathbb{C}^n)$ and non-zero $|\psi\rangle \in \mathbb{C}^n$,

$$|\psi\rangle, A|\psi\rangle, A^2|\psi\rangle, \dots, A^n|\psi\rangle$$

are linearly **dependent**. There exist a_0, \dots, a_n that are not all-zero satisfying

$$\begin{aligned} 0 &= a_0 |\psi\rangle + a_1 A |\psi\rangle + \dots + a_n A^n |\psi\rangle \\ &= a_m (A - \lambda_1 I)(A - \lambda_2 I) \dots (A - \lambda_m I) |\psi\rangle \end{aligned}$$

where m is the largest i such that $a_i \neq 0$.

This means that there exist $i \in \{1, 2, \dots, m\}$ and non-zero $|\varphi\rangle \in \mathbb{C}^n$ such that

$$(A - \lambda_i I) |\varphi\rangle = 0.$$

Orthogonal projection

For linear space V and its subspace W , the **orthogonal projection** onto W is defined by

$$P = \sum_j |\psi_j\rangle \langle \psi_j|$$

where $(|\psi_j\rangle)_j$ forms an orthonormal basis of W .

- P is Hermitian
- $P^2 = P$
- $P|\psi\rangle \in W$ for any $|\psi\rangle \in V$
- $P|\psi\rangle = |\psi\rangle$ for any $|\psi\rangle \in W$
- $I - P$ is the orthogonal projection onto W_\perp

Any normal matrix has a spectral decomposition

Induction on the dimension n . Spectral decomposition theorem obviously holds for $n = 1$. A has a eigenvalue λ and corresponding eigenspace W . Let P be the orthogonal projection onto W . Let $Q = I - P$.

$$A = (P + Q)A(P + Q) = PAP + PAQ + QAP + QAQ$$

- $PAP = \lambda P$.
- $QAP = Q\lambda P = 0$.
- For $|\psi\rangle \in W$, $AA^\dagger |\psi\rangle = A^\dagger A |\psi\rangle = \lambda A^\dagger |\psi\rangle$ which means $A^\dagger |\psi\rangle \in W$. This implies $(PAQ)^\dagger = QA^\dagger P = 0$.

Hence, $A = \lambda P + QAQ$. Since $QA = QA(P + Q) = QAQ$ and $QA^\dagger = QA^\dagger(P + Q) = QA^\dagger Q$,

$$\begin{aligned}(QAQ)(QA^\dagger Q) &= QAA^\dagger Q \\ &= QA^\dagger AQ = QA^\dagger QQAQ\end{aligned}$$

Hence, QAQ can be regarded as a normal linear operator on W_\perp . From the hypothesis of induction, QAQ has a spectral decomposition.

Terminology

- **Density matrix**, density operator: A Hermitian matrix ρ that represents a state, i.e., $\rho \succeq 0$, $\text{Tr}(\rho) = 1$.
- **Pure state**: A state that cannot be written as a convex combination of other states. Equivalently, its a density operator with rank one.
- **State vector**: A complex unit vector $|\psi\rangle$ that represents a pure state $\rho = |\psi\rangle \langle\psi|$.
- **Mixed state**: A state that is not a pure state.
- Positive operator-valued measurement (**POVM**): A tuple $\{P_j\}$ of Hermitian matrices that represents a measurement, i.e., $P_j \succeq 0$ and $\sum_j P_j = I$.

Ensemble of states

Let ρ_1, \dots, ρ_k be density matrices. If ρ_i is prepared with probability p_i , and POVM $\{P_j\}$ is applied, outcome j is obtained with probability

$$\sum_{i=1}^k p_i \text{Tr}(\rho_i P_j) = \text{Tr} \left(\sum_{i=1}^k p_i \rho_i P_j \right).$$

Hence, this ensemble of states is represented by $\rho := \sum_i p_i \rho_i$.

Ensemble of pure states

Any quantum state

$$\rho = \sum_i \lambda_i |\psi_i\rangle \langle \psi_i|$$

for $(\lambda_i \geq 0)_i$ can be regarded as an **ensemble** $(\lambda_i, |\psi_i\rangle \langle \psi_i|)_i$ of pure states.

$$\begin{aligned}\rho &= \frac{3}{4} |0\rangle \langle 0| + \frac{1}{4} |1\rangle \langle 1| \\ &= \frac{1}{2} |a\rangle \langle a| + \frac{1}{2} |b\rangle \langle b|\end{aligned}$$

for

$$\begin{aligned}|a\rangle &:= \sqrt{\frac{3}{4}} |0\rangle + \sqrt{\frac{1}{4}} |1\rangle \\ |b\rangle &:= \sqrt{\frac{3}{4}} |0\rangle - \sqrt{\frac{1}{4}} |1\rangle .\end{aligned}$$

Observable

Let $\{P_j\}$ be a POVM. If we assign real value a_j for each outcome j , its expectation is

$$\mathbb{E}[a] := \sum_j a_j \text{Tr}(\rho P_j) = \text{Tr} \left(\rho \sum_j a_j P_j \right) = \text{Tr}(\rho A).$$

Here, Hermitian operator $A := \sum_j a_j P_j$ is called a **observable**.

If $\{P_j\}$ is a **projective measurement**, i.e., $P_j P_k = \delta_{j,k} P_j$,

$$\mathbb{E}[a^n] := \sum_j a_j^n \text{Tr}(\rho P_j) = \text{Tr} \left(\rho \sum_j a_j^n P_j \right) = \text{Tr}(\rho A^n).$$

For example, X and Z are observables for POVMs $\{|+\rangle\langle +|, |-\rangle\langle -|\}$ and $\{|0\rangle\langle 0|, |1\rangle\langle 1|\}$ with the assignments ± 1 , respectively.

Decoherence

For orthonormal basis $\{|\psi_i\rangle\}$, POVM $\{|\psi_i\rangle\langle\psi_i|\}$ is performed to a quantum state ρ . If outcome is i , the quantum state ρ is **transformed** into $|\psi_i\rangle\langle\psi_i|$. If we **don't see** the measurement outcome, the state after the measurement is

$$\sum_i \text{Tr}(\rho |\psi_i\rangle\langle\psi_i|) |\psi_i\rangle\langle\psi_i| = \sum_i \langle\psi_i|\rho|\psi_i\rangle |\psi_i\rangle\langle\psi_i|$$

$$\rho = \sum_{i,j} \rho_{i,j} |\psi_i\rangle\langle\psi_j| \mapsto \sum_i \rho_{i,i} |\psi_i\rangle\langle\psi_i|$$

This phenomenon is called **decoherence**.

Purification

Theorem

For any density matrix ρ on V , there exists a pure state $|\psi\rangle$ of a composite system on $V \otimes W$ for some W such that $\text{Tr}_W(|\psi\rangle\langle\psi|) = \rho$.

Proof.

For a spectral decomposition of ρ

$$\rho = \sum_i \lambda_i |\psi_i\rangle_V \langle\psi_i|_V$$

let

$$|\psi\rangle_{V \otimes W} := \sum_i \sqrt{\lambda_i} |\psi_i\rangle_V |i\rangle_W$$

where $\{|i\rangle\}_i$ is an arbitrary orthonormal basis of W . Then,

$$\text{Tr}_W(|\psi\rangle_{V \otimes W} \langle\psi|_{V \otimes W}) = \rho.$$



$|\psi\rangle_{V \otimes W}$ is called a **purification** of ρ .

Quantum states discrimination

Alice encodes her classical information $\{1, 2, \dots, n\}$ into quantum states $\rho_1, \rho_2, \dots, \rho_n \in \mathcal{H}(\mathbb{C}^m)$, and send it to Bob. Bob performs a POVM $\{P_1, \dots, P_n\}$ for estimating $i \in \{1, \dots, n\}$ that Alice encoded. Assume Bob could estimate i without error. Then,

$$\text{Tr}(\rho_i P_j) = \delta_{ij}.$$

$$\begin{aligned}\text{Tr}(\rho_i P_j) &= \text{Tr} \left(\sum_k \lambda_k^{(i)} |\psi_k^{(i)}\rangle \langle \psi_k^{(i)}| P_j \right) \\ &= \sum_k \lambda_k^{(i)} \langle \psi_k^{(i)}| P_j |\psi_k^{(i)}\rangle\end{aligned}$$

$\langle \psi_k^{(i)}| P_j |\psi_k^{(i)}\rangle = \delta_{ij}$ implies $|\psi_k^{(i)}\rangle$ is an eigenvector of P_j with eigenvalue δ_{ij} . Hence, $\langle \psi_k^{(i)}| \psi_\ell^{(j)}\rangle = 0$ for $i \neq j$ which implies $n \leq m$.

Superdense coding

Alice can send **two** bits to Bob by sending a single qubit and using a shared Bell state.

$$\begin{aligned} |\Phi_{00}\rangle &= \frac{1}{\sqrt{2}}(|0\rangle_A |0\rangle_B + |1\rangle_A |1\rangle_B) \\ |\Phi_{01}\rangle &= \frac{1}{\sqrt{2}}(|0\rangle_A |1\rangle_B + |1\rangle_A |0\rangle_B), && \text{by } X \\ |\Phi_{10}\rangle &= \frac{1}{\sqrt{2}}(|0\rangle_A |0\rangle_B - |1\rangle_A |1\rangle_B), && \text{by } Z \\ |\Phi_{11}\rangle &= \frac{1}{\sqrt{2}}(|0\rangle_A |1\rangle_B - |1\rangle_A |0\rangle_B), && \text{by } ZX \end{aligned}$$

These are **orthogonal**.

Assignments

- ① Show the reduced density matrix $\rho_V \in \mathcal{H}(V)$ of

$$\rho = \sum_{i,j} \rho_{i,j} |i\rangle \langle j| \otimes |i\rangle \langle j| \in \mathcal{H}(V \otimes W)$$

where $\{|i\rangle\}$ is a orthonormal basis of V and W .

- ② For a single-qubit density matrix $\rho = \sum_{i,j=0}^1 \rho_{i,j} |i\rangle \langle j|$, show the density matrix of an ensemble of quantum states ρ and $Z\rho Z$ chosen with probabilities $1/2$.
- ③ Show a purification of $\rho = \frac{3}{4} |0\rangle \langle 0| + \frac{1}{4} |1\rangle \langle 1|$.