

# Parametric Matrix Models

Model Emulation, Model Discovery,  
and General Machine Learning

**Patrick Cook**

Danny Jammooa, Morten Hjorth-Jensen, Daniel D. Lee, Dean Lee

Facility for Rare Isotope Beams  
Michigan State University

Theory Seminar  
Institute for Nuclear Physics  
TU Darmstadt



# Parametric Matrix Models

Model Emulation, Model Discovery,  
~~and General Machine Learning~~

**Patrick Cook**

Danny Jammooa, Morten Hjorth-Jensen, Daniel D. Lee, Dean Lee

Facility for Rare Isotope Beams  
Michigan State University

Theory Seminar  
Institute for Nuclear Physics  
TU Darmstadt



- ▶ Your system explores a complicated phase space



- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough



- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough
- ▶ Linear models are better, but not great, and also slower




- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough
- ▶ Linear models are better, but not great, and also slower
- ▶ Neural networks are powerful...



- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough
- ▶ Linear models are better, but not great, and also slower
- ▶ Neural networks are powerful... too powerful...

# NEURAL NETWORKS



- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough
- ▶ Linear models are better, but not great, and also slower
- ▶ Neural networks are powerful... too powerful...



- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough
- ▶ Linear models are better, but not great, and also slower
- ▶ Neural networks are powerful... too powerful...

# NEURAL NETWORKS



- ▶ Your system explores a complicated phase space
- ▶ Simple fits are fast but rough
- ▶ Linear models are better, but not great, and also slower
- ▶ Neural networks are powerful... too powerful...
- ▶ **Parametric Matrix Models** are your system!

# NEURAL NETWORKS



# Build-a-PMM<sup>TM</sup>

# Build-a-PMM<sup>TM</sup>

## Your Underlying Equations

$$\begin{cases} H(c) = H_0 + cH_1 \\ H(c) |\psi(c)\rangle = E(c) |\psi(c)\rangle \\ \langle\psi(c)|O|\psi(c)\rangle = f(c) \end{cases}$$

# Build-a-PMM<sup>TM</sup>

## Your Underlying Equations

$$\begin{cases} H(c) = H_0 + cH_1 \\ H(c) |\psi(c)\rangle = E(c) |\psi(c)\rangle \\ \langle\psi(c)|O|\psi(c)\rangle = f(c) \end{cases}$$

---

$$\begin{cases} \frac{\partial \vec{v}}{\partial t} = A(\vec{c})\vec{v} + f(t, c, \vec{v}) \\ \vec{v}(t = 0) = \vec{v}_0 \\ v_i(t) = \text{b.c.} \end{cases}$$

# Build-a-PMM<sup>TM</sup>

## Your Underlying Equations

$$\begin{cases} H(c) = H_0 + cH_1 \\ H(c)|\psi(c)\rangle = E(c)|\psi(c)\rangle \\ \langle\psi(c)|O|\psi(c)\rangle = f(c) \end{cases}$$

## Your PMM

$$\begin{cases} M(c) = \underline{M}_0 + c\underline{M}_1 \\ M(c)|\tilde{\psi}(c)\rangle = \tilde{E}(c)|\tilde{\psi}(c)\rangle \\ \langle\tilde{\psi}(c)|\underline{X}|\tilde{\psi}(c)\rangle = \tilde{f}(c) \end{cases}$$

---


$$\begin{cases} \frac{\partial \vec{v}}{\partial t} = A(\vec{c})\vec{v} + f(t, c, \vec{v}) \\ \vec{v}(t=0) = \vec{v}_0 \\ v_i(t) = \text{b.c.} \end{cases}$$

# Build-a-PMM™

## Your Underlying Equations

$$\begin{cases} H(c) = H_0 + cH_1 \\ H(c)|\psi(c)\rangle = E(c)|\psi(c)\rangle \\ \langle\psi(c)|O|\psi(c)\rangle = f(c) \end{cases}$$

$$\begin{cases} \frac{\partial \vec{v}}{\partial t} = A(\vec{c})\vec{v} + f(t, c, \vec{v}) \\ \vec{v}(t=0) = \vec{v}_0 \\ v_i(t) = \text{b.c.} \end{cases}$$

## Your PMM

$$\begin{cases} M(c) = \underline{M_0} + c\underline{M_1} \\ M(c)|\tilde{\psi}(c)\rangle = \tilde{E}(c)|\tilde{\psi}(c)\rangle \\ \langle\tilde{\psi}(c)|\underline{X}|\tilde{\psi}(c)\rangle = \tilde{f}(c) \end{cases}$$

Okay it's not always *that* simple...

# Build-a-PMM™

## Your Underlying Equations

$$\begin{cases} H(c) = H_0 + cH_1 \\ H(c)|\psi(c)\rangle = E(c)|\psi(c)\rangle \\ \langle\psi(c)|O|\psi(c)\rangle = f(c) \end{cases}$$

$$\begin{cases} \frac{\partial \vec{v}}{\partial t} = A(\vec{c})\vec{v} + f(t, c, \vec{v}) \\ \vec{v}(t=0) = \vec{v}_0 \\ v_i(t) = \text{b.c.} \end{cases}$$

## Your PMM

$$\begin{cases} M(c) = \underline{M_0} + c\underline{M_1} \\ M(c)|\tilde{\psi}(c)\rangle = \tilde{E}(c)|\tilde{\psi}(c)\rangle \\ \langle\tilde{\psi}(c)|\underline{X}|\tilde{\psi}(c)\rangle = \tilde{f}(c) \end{cases}$$

Okay it's not always *that* simple...


**Then fit to whatever data you have!**

# Example: Anharmonic Lipkin Model

$$H(\xi; \alpha) = (1 - \xi)\hat{n} + \frac{2\xi}{S} \left( S^2 - \hat{S}_x^2 \right) + \frac{\alpha}{2S} \hat{n}(\hat{n} + 1)$$

$$\hat{n} = \left( S + \hat{S}_z \right)$$

# Example: Anharmonic Lipkin Model



Gamito, Khalouf-Rivera, et al. 2022

# Example: Anharmonic Lipkin Model

$$H_\alpha(\xi) = H_0 + \xi H_1$$

$$\langle E_0(\xi) | \hat{n} | E_0(\xi) \rangle$$

# Example: Anharmonic Lipkin Model

$$H_\alpha(\xi) = H_0 + \xi H_1$$


$$\langle E_0(\xi) | \hat{n} | E_0(\xi) \rangle$$

**PMM**


$$M_\alpha(\xi) = \underline{M_0} + \xi \underline{M_1}$$

$$\langle \tilde{E}_0(\xi) | \underline{X} | \tilde{E}_0(\xi) \rangle$$

# Example: Anharmonic Lipkin Model



# Example: Anharmonic Lipkin Model



# Example: Anharmonic Lipkin Model




# Example: Anharmonic Lipkin Model



Neural Network:  
“Sometimes,  $E_0 > E_1$ ”


# Example: Anharmonic Lipkin Model

## WHAT



Neural Network:  
“Sometimes,  $E_0 > E_1$ ”

# Example: Anharmonic Lipkin Model



# Isn't this just Eigenvector Continuation?

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

Pros:

- ▶ Exponential convergence

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

Cons:

- ▶ Poor computational scaling

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

Cons:

- ▶ Poor computational scaling
- ▶ Limited in scope

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

Cons:

- ▶ Poor computational scaling
- ▶ Limited in scope
- ▶ Curse of dimensionality

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$

$$H(c) = \frac{1}{2N} \sum_i^N (\sigma_i^z + c\sigma_i^x)$$

Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

Cons:

- ▶ Poor computational scaling
- ▶ Limited in scope
- ▶ Curse of dimensionality

# Isn't this just Eigenvector Continuation?

## Eigenvector Continuation

Requires:

- ▶ Snapshots of full eigenvectors
- ▶ The evaluation of  $H(c)$


Pros:

- ▶ Exponential convergence
- ▶ Constructed rather than trained

Cons:

- ▶ Poor computational scaling
- ▶ Limited in scope
- ▶ Curse of dimensionality

$$H(c) = \frac{1}{2N} \sum_i^N (\sigma_i^z + c\sigma_i^x)$$



# Time evolution on a quantum computer of a non-commuting Hamiltonian

$$H = H_0 + H_1 + \cdots + H_l$$

# Time evolution on a quantum computer of a non-commuting Hamiltonian

$$H = H_0 + H_1 + \cdots + H_l$$

$$U(dt) = e^{-iHdt} \approx e^{-iH_0dt} e^{-iH_1dt} \cdots e^{-iH_ldt}$$

# Time evolution on a quantum computer of a non-commuting Hamiltonian

$$H = H_0 + H_1 + \cdots + H_l$$

$$U(dt) = e^{-iHdt} \approx e^{-iH_0dt} e^{-iH_1dt} \cdots e^{-iH_ldt}$$

**PMM**


$$\tilde{U}(dt) = e^{-i\underline{M}_0dt} e^{-i\underline{M}_1dt} \cdots e^{-i\underline{M}_ldt}$$

## 1D Heisenberg Model with Dzyaloshinskii-Moriya Interaction

$$H = B \sum_i^N r_i \sigma_i^z + J \sum_{u \in \{x,y,z\}} \sum_i^N \sigma_i^u \sigma_{i+1}^u + D \sum_i^N (\sigma_i^x \sigma_{i+1}^y - \sigma_i^y \sigma_{i+1}^x)$$


# 1D Heisenberg Model with Dzyaloshinskii-Moriya Interaction

$$H = B \sum_i^N r_i \sigma_i^z + J \sum_{u \in \{x, y, z\}} \sum_i^N \sigma_i^u \sigma_{i+1}^u + D \sum_i^N (\sigma_i^x \sigma_{i+1}^y - \sigma_i^y \sigma_{i+1}^x)$$




# 1D Heisenberg Model with Dzyaloshinskii-Moriya Interaction

$$H = B \sum_i^N r_i \sigma_i^z + J \sum_{u \in \{x, y, z\}} \sum_i^N \sigma_i^u \sigma_{i+1}^u + D \sum_i^N (\sigma_i^x \sigma_{i+1}^y - \sigma_i^y \sigma_{i+1}^x)$$




# 1D Heisenberg Model with Dzyaloshinskii-Moriya Interaction

$$H = B \sum_i^N r_i \sigma_i^z + J \sum_{u \in \{x, y, z\}} \sum_i^N \sigma_i^u \sigma_{i+1}^u + D \sum_i^N (\sigma_i^x \sigma_{i+1}^y - \sigma_i^y \sigma_{i+1}^x)$$



# 1D Heisenberg Model with Dzyaloshinskii-Moriya Interaction

$$H = B \sum_i^N r_i \sigma_i^z + J \sum_{u \in \{x,y,z\}} \sum_i^N \sigma_i^u \sigma_{i+1}^u + D \sum_i^N (\sigma_i^x \sigma_{i+1}^y - \sigma_i^y \sigma_{i+1}^x)$$



# Practical Summary

# Practical Summary

**The PMM for your system *is* your system!**

# Practical Summary

**The PMM for your system *is* your system!**

- ▶ Use the underlying equations of your system

# Practical Summary

**The PMM for your system *is* your system!**

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system

# Practical Summary

## The PMM for your system *is* your system!

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system
- ▶ Low-order equations are perfectly fine

# Practical Summary

## The PMM for your system *is* your system!

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system
- ▶ Low-order equations are perfectly fine
- ▶ Due to low dimensionality, exact solutions are possible

# Practical Summary

## The PMM for your system *is* your system!

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system
- ▶ Low-order equations are perfectly fine
- ▶ Due to low dimensionality, exact solutions are possible

# Practical Summary

**The PMM for your system *is* your system!**

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system
- ▶ Low-order equations are perfectly fine
- ▶ Due to low dimensionality, exact solutions are possible

**Two outstanding questions:**

# Practical Summary

## The PMM for your system *is* your system!

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system
- ▶ Low-order equations are perfectly fine
- ▶ Due to low dimensionality, exact solutions are possible

## Two outstanding questions:

- ▶ *What if I don't know the underlying equations?*

# Practical Summary

## The PMM for your system *is* your system!

- ▶ Use the underlying equations of your system
- ▶ The PMM will be a low-dimension analog of your system
- ▶ Low-order equations are perfectly fine
- ▶ Due to low dimensionality, exact solutions are possible

## Two outstanding questions:

- ▶ *What if I don't know the underlying equations?*
- ▶ *What if my problem is nonlinear?*

**arXiv:2404.11566**

# Emulators for scarce and noisy data: application to auxiliary field diffusion Monte Carlo for the deuteron

Rahul Somasundaram,<sup>1, 2,\*</sup> Cassandra L. Armstrong,<sup>3</sup>  
Pablo Giuliani,<sup>4, 5</sup> Kyle Godbey,<sup>4</sup> Stefano Gandolfi,<sup>1</sup> and Ingo Tews<sup>1</sup>

<sup>1</sup>*Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

<sup>2</sup>*Department of Physics, Syracuse University, Syracuse, NY 13244, USA*

<sup>3</sup>*Intelligence and Space Research Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

<sup>4</sup>*Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>5</sup>*Department of Statistics and Probability, Michigan State University, East Lansing, Michigan 48824, USA*

arXiv:2404.11566

# Emulators for scarce and noisy data: application to auxiliary field diffusion Monte Carlo for the deuteron

Rahul Somasundaram,<sup>1, 2,\*</sup> Cassandra L. Armstrong,<sup>3</sup>  
Pablo Giuliani,<sup>4, 5</sup> Kyle Godfrey,<sup>4</sup> Stefano Gandolfi,<sup>1</sup> and Ingo Tews<sup>1</sup>


<sup>1</sup> Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>2</sup> Department of Physics, Syracuse University, Syracuse, NY 13244, USA

<sup>3</sup> Intelligence and Space Research Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>4</sup> Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA

<sup>5</sup> Department of Statistics and Probability, Michigan State University, East Lansing, Michigan 48824, USA



arXiv:2404.11566

# Emulators for scarce and noisy data: application to auxiliary field diffusion Monte Carlo for the deuteron

Rahul Somasundaram,<sup>1, 2,\*</sup> Cassandra L. Armstrong,<sup>3</sup>  
Pablo Giuliani,<sup>4, 5</sup> Kyle Godfrey,<sup>4</sup> Stefano Gandolfi,<sup>1</sup> and Ingo Tews<sup>1</sup>



<sup>1</sup> Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>2</sup> Department of Physics, Syracuse University, Syracuse, NY 13244, USA

<sup>3</sup> Intelligence and Space Research Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>4</sup> Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA

<sup>5</sup> Department of Statistics and Probability, Michigan State University, East Lansing, Michigan 48824, USA



**arXiv:2405.20558**

Towards accelerated nuclear-physics parameter estimation from binary neutron star mergers:  
Emulators for the Tolman-Oppenheimer-Volkoff equations

BRENDAN T. REED,<sup>1</sup> RAHUL SOMASUNDARAM,<sup>1,2</sup> SOUMI DE,<sup>1</sup> CASSANDRA L. ARMSTRONG,<sup>3</sup> PABLO GIULIANI,<sup>4</sup>  
COLLIN CAPANO,<sup>2,5</sup> DUNCAN A. BROWN,<sup>2</sup> AND INGO TEWS<sup>1</sup>

<sup>1</sup>*Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

<sup>2</sup>*Department of Physics, Syracuse University, Syracuse, NY 13244, USA*

<sup>3</sup>*Intelligence and Space Research Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA*

<sup>4</sup>*Facility for Rare Isotope Beams, Michigan State University, East Lansing, Michigan 48824, USA*

<sup>5</sup>*Physics Department, University of Massachusetts Dartmouth, North Dartmouth, MA 02747, USA*

**arXiv:2410.00247**

# Inferring three-nucleon couplings from multi-messenger neutron-star observations

Rahul Somasundaram<sup>1,2</sup>, Isak Svensson<sup>3,4,5</sup>, Soumi De<sup>2</sup>,  
Andrew E. Deneris<sup>6</sup>, Yannick Dietz<sup>3,4</sup>, Philippe Landry<sup>7,8</sup>,  
Achim Schwenk<sup>3,4,5</sup>, and Ingo Tews<sup>2</sup>

<sup>1</sup>Department of Physics, Syracuse University, Syracuse, NY 13244, USA

<sup>2</sup>Theoretical Division, Los Alamos National Laboratory, Los Alamos, NM 87545, USA

<sup>3</sup>Technische Universität Darmstadt, Department of Physics, 64289 Darmstadt, Germany

<sup>4</sup>ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH,  
64291 Darmstadt, Germany

<sup>5</sup>Max-Planck-Institut für Kernphysik, Saupfercheckweg 1, 69117 Heidelberg, Germany

<sup>6</sup>Computer, Computational, and Statistical Sciences Division, Los Alamos National  
Laboratory, Los Alamos, NM 87545, USA

<sup>7</sup>Canadian Institute for Theoretical Astrophysics, University of Toronto, Toronto, Ontario M5S  
3H8, Canada

<sup>8</sup>Perimeter Institute for Theoretical Physics, Waterloo, Ontario N2L 2Y5, Canada

# From Data to Discovery


# Anharmonic Lipkin Model

# Anharmonic Lipkin Model

$$\left\{ \begin{array}{l} H_\alpha(\xi) = H_0 + \xi H_1 \\ \langle E_0(\xi) | \hat{n} | E_0(\xi) \rangle \end{array} \right\} \sim \left\{ \begin{array}{l} M_\alpha(\xi) = \underline{M}_0 + \xi \underline{M}_1 \\ \langle \tilde{E}_0(\xi) | \underline{X} | \tilde{E}_0(\xi) \rangle \end{array} \right\}$$


# Anharmonic Lipkin Model

$$\left\{ \begin{array}{l} H_\alpha(\xi) = H_0 + \xi H_1 \\ \langle E_0(\xi) | \hat{n} | E_0(\xi) \rangle \end{array} \right\} \sim \left\{ \begin{array}{l} M_\alpha(\xi) = \underline{M}_0 + \xi \underline{M}_1 \\ \langle \tilde{E}_0(\xi) | \underline{X} | \tilde{E}_0(\xi) \rangle \end{array} \right\}$$




# Generalized Long-Range Spin Model


$$H = - \sum_{i \neq j} J_{ij} (\gamma_x \sigma_i^x \sigma_j^x + \gamma_y \sigma_i^y \sigma_j^y) - B \sum_i \sigma_i^z$$



# Generalized Long-Range Spin Model




$$H = - \sum_{i \neq j} J_{ij} (\gamma_x \sigma_i^x \sigma_j^x + \gamma_y \sigma_i^y \sigma_j^y) - B \sum_i \sigma_i^z$$



$^{171}\text{Yb}^+$


# Generalized Long-Range Spin Model




$$H = - \sum_{i \neq j} J_{ij} (\gamma_x \sigma_i^x \sigma_j^x + \gamma_y \sigma_i^y \sigma_j^y) - B \sum_i \sigma_i^z$$

$$M = - \sum_{i \neq j} \tilde{J}_{ij} (\tilde{\gamma}_x \sigma_i^x \sigma_j^x + \tilde{\gamma}_y \sigma_i^y \sigma_j^y) - \tilde{B} \sum_i \sigma_i^z$$

$^{171}\text{Yb}^+$



# Generalized Long-Range Spin Model





$$H = - \sum_{i \neq j} J_{ij} (\gamma_x \sigma_i^x \sigma_j^x + \gamma_y \sigma_i^y \sigma_j^y) - B \sum_i \sigma_i^z$$


$$M = - \sum_{i \neq j} \tilde{J}_{ij} (\tilde{\gamma}_x \sigma_i^x \sigma_j^x + \tilde{\gamma}_y \sigma_i^y \sigma_j^y) - \tilde{B} \sum_i \sigma_i^z$$


$^{171}\text{Yb}^+$


Supervised and **Unsupervised**




Supervised  $\{(|\Psi_k\rangle, E_k)\}$ 


















$$M = T + \underline{V}(|\vec{r}_{^{12}\text{C}} - \vec{r}_\alpha|)$$



$$M_2 = T + \underline{V}(|\vec{r}_{^{12}\text{C}} - \vec{r}_\alpha|)$$

$$M_3 = T + \underline{V}(\vec{r}_{^8\text{Be}}, \vec{r}_{\alpha_1}, \vec{r}_{\alpha_2})$$

$$M_4 = T + \underline{V}(\vec{r}_{\alpha_1}, \vec{r}_{\alpha_2}, \vec{r}_{\alpha_3}, \vec{r}_{\alpha_4})$$

# Extending PMMs

**Two main requests:**

# Extending PMMs

**Two main requests:**

- ▶ Predicting unseen observables

# Extending PMMs

## Two main requests:

- ▶ Predicting unseen observables
  - ▶ State-vector emulation

# Extending PMMs

## Two main requests:

- ▶ Predicting unseen observables
  - ▶ State-vector emulation
- ▶ Nonlinear problems

# Extending PMMs

## Two main requests:

- ▶ Predicting unseen observables
  - ▶ State-vector emulation
- ▶ Nonlinear problems
  - ▶ Density-dependent Hamiltonians

# Extending PMMs

## Two main requests:

- ▶ Predicting unseen observables
  - ▶ State-vector emulation
- ▶ Nonlinear problems
  - ▶ Density-dependent Hamiltonians

# Extending PMMs

## Two main requests:

- ▶ Predicting unseen observables
  - ▶ State-vector emulation
- ▶ Nonlinear problems
  - ▶ Density-dependent Hamiltonians

**These are the same problem!**

# Proper Orthogonal Decomposition

# Proper Orthogonal Decomposition

**Given:**

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

**Example**

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

**Example**

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

**Example**

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

**POD:**

**Example**

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

# Proper Orthogonal Decomposition

**Given:**

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

**POD:**

- ▶  $r$  most important principal components of the snapshots

**Example**

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

$$u = P^\dagger v$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

$$u = P^\dagger v$$

$$\tilde{A} = P^\dagger AP$$

$$du/dt = \tilde{A}u + P^\dagger f(t, Pu)$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis
- ▶ Solve the projected equations

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

$$u = P^\dagger v$$

$$\tilde{A} = P^\dagger AP$$

$$du/dt = \tilde{A}u + P^\dagger f(t, Pu)$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis
- ▶ Solve the projected equations
- ▶ Reconstruct the full solution

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

$$u = P^\dagger v$$

$$\tilde{A} = P^\dagger AP$$

$$du/dt = \tilde{A}u + P^\dagger f(t, Pu)$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis
- ▶ Solve the projected equations
- ▶ Reconstruct the full solution

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

$$u = P^\dagger v$$

$$\tilde{A} = P^\dagger AP$$

$$du/dt = \tilde{A}u + P^\dagger f(t, Pu)$$

$$v \approx Pu$$

# Proper Orthogonal Decomposition

## Given:

- ▶ A set of state snapshots  $\{\vec{v}_i\}$
- ▶ The governing equations for  $\vec{v}$

## POD:

- ▶  $r$  most important principal components of the snapshots
- ▶ Project the governing equations onto the POD basis
- ▶ Solve the projected equations
- ▶ Reconstruct the full solution

## Example

$$\{v(t_0), v(t_1), \dots, v(t_p)\}$$

$$dv/dt = Av + f(t, v)$$

$$V = [v(t_0) \quad v(t_1) \quad \cdots \quad v(t_p)]$$

$$V \approx P\Sigma_r Q^\dagger \quad \text{SVD}$$

$$u = P^\dagger v$$

$$\tilde{A} = P^\dagger AP$$

$$du/dt = \tilde{A}u + P^\dagger f(t, Pu)$$

$$v \approx Pu$$

# POD for Nonlinear Problems

# POD for Nonlinear Problems

**Consider the simplest nonlinearity:**

# POD for Nonlinear Problems

**Consider the simplest nonlinearity:**

$$f(t, v) = v^2 = v \odot v$$

# POD for Nonlinear Problems

**Consider the simplest nonlinearity:**

$$f(t, v) = v^2 = v \odot v$$

**POD:**

# POD for Nonlinear Problems

**Consider the simplest nonlinearity:**

$$f(t, v) = v^2 = v \odot v$$

**POD:**

$$P^\dagger f(t, Pu) = P^\dagger [(Pu) \odot (Pu)]$$

# POD for Nonlinear Problems

Consider the simplest nonlinearity:

$$f(t, v) = v^2 = v \odot v$$

POD:

$$\begin{aligned} P^\dagger f(t, Pu) &= P^\dagger [(Pu) \odot (Pu)] \\ &\stackrel{i}{=} (P^\dagger)_{i\mu} P_{\mu j} P_{\mu k} u_j u_k \end{aligned}$$

# POD for Nonlinear Problems

Consider the simplest nonlinearity:

$$f(t, v) = v^2 = v \odot v$$

POD:

$$\begin{aligned} P^\dagger f(t, Pu) &= P^\dagger [(Pu) \odot (Pu)] \\ &\stackrel{i}{=} (P^\dagger)_{i\mu} P_{\mu j} P_{\mu k} u_j u_k \\ &\equiv P^\dagger (P \otimes P)(u \otimes u) \end{aligned}$$

# POD for Nonlinear Problems

**Consider the simplest nonlinearity:**

$$f(t, v) = v^2 = v \odot v$$

**POD:**

$$\begin{aligned} P^\dagger f(t, Pu) &= P^\dagger [(Pu) \odot (Pu)] \\ &\stackrel{i}{=} (P^\dagger)_{i\mu} P_{\mu j} P_{\mu k} u_j u_k \\ &\equiv P^\dagger (P \otimes P)(u \otimes u) \end{aligned}$$

**Generalizing to polynomial nonlinearities:**

$$v^n = v \odot^n v$$



$$P^\dagger (P \otimes^n P)(u \otimes^n u) \stackrel{i}{=} (P^\dagger)_{i\mu} P_{\mu j_1} P_{\mu j_2} \cdots P_{\mu j_n} u_{j_1} u_{j_2} \cdots u_{j_n}$$

# Back to PMMs

$$P \otimes^n P \rightarrow \underline{\mathbb{T}^n} \equiv \underline{T_{j_1 j_2 \dots j_n}}$$

# Gross Pitaevskii Equation

$$i\frac{\partial\Psi}{\partial t} = H_0\Psi + g|\Psi|^2\Psi$$

# Gross Pitaevskii Equation

$$i \frac{\partial \Psi}{\partial t} = H_0 \Psi + g |\Psi|^2 \Psi$$

# Gross Pitaevskii Equation

$$i\frac{\partial\Psi}{\partial t} = H_0\Psi + g\Psi \odot \Psi^* \odot \Psi$$

# Gross Pitaevskii Equation


$$i \frac{\partial \Psi}{\partial t} = H_0 \Psi + g \Psi \odot \Psi^* \odot \Psi$$


$$i \frac{\partial \psi}{\partial t} = \tilde{H}_0 \psi + g \mathbb{T}^3 (\psi \otimes \psi^* \otimes \psi)$$


# Gross Pitaevskii Equation


$$i \frac{\partial \Psi}{\partial t} = H_0 \Psi + g \Psi \odot \Psi^* \odot \Psi$$

$$i \frac{\partial \psi}{\partial t} = \tilde{H}_0 \psi + g \underline{T}_{jkl} \psi_j \psi_k^* \psi_l$$









# Beyond POD

**What if the snapshots aren't of the state vector?**

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*)$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*) \approx P_\rho^\dagger (P_\Psi \otimes P_\Psi^*) (\psi \otimes \psi^*)$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*) \approx P_\rho^\dagger (P_\Psi \otimes P_\Psi^*) (\psi \otimes \psi^*)$$

$$\tilde{H}_0 = P_\Psi^\dagger H_0 P_\Psi$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*) \approx P_\rho^\dagger (P_\Psi \otimes P_\Psi^*) (\psi \otimes \psi^*)$$

$$\tilde{H}_0 = P_\Psi^\dagger H_0 P_\Psi$$

$$\frac{d\psi}{dt} = \tilde{H}_0 \psi + g P_\Psi^\dagger (P_\Psi \otimes P_\Psi^* \otimes P_\Psi) (\psi \otimes \psi^* \otimes \psi)$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*) \approx P_\rho^\dagger (P_\Psi \otimes P_\Psi^*) (\psi \otimes \psi^*)$$

$$\tilde{H}_0 = P_\Psi^\dagger H_0 P_\Psi$$

$$\frac{d\psi}{dt} = \tilde{H}_0 \psi + g P_\Psi^\dagger (P_\Psi \otimes P_\Psi^* \otimes P_\Psi) (\psi \otimes \psi^* \otimes \psi)$$

$$\rho \approx P_\rho q$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*) \approx P_\rho^\dagger P_\Psi P_\Psi^\dagger (P_\Psi \otimes P_\Psi^*) (\psi \otimes \psi^*)$$

$$\tilde{H}_0 = P_\Psi^\dagger H_0 P_\Psi$$

$$\frac{d\psi}{dt} = \tilde{H}_0 \psi + g P_\Psi^\dagger (P_\Psi \otimes P_\Psi^* \otimes P_\Psi) (\psi \otimes \psi^* \otimes \psi)$$

$$\rho \approx P_\rho q$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi = P_\Psi^\dagger \Psi$$

$$q = P_\rho^\dagger \rho = P_\rho^\dagger (\Psi \odot \Psi^*) \approx \underbrace{P_\rho^\dagger P_\Psi}_{B_\rho^\Psi} \underbrace{P_\Psi^\dagger (P_\Psi \otimes P_\Psi^*)}_{\mathbb{T}_\Psi^2} (\psi \otimes \psi^*)$$

$$\tilde{H}_0 = P_\Psi^\dagger H_0 P_\Psi$$

$$\frac{d\psi}{dt} = \tilde{H}_0 \psi + g P_\Psi^\dagger (P_\Psi \otimes P_\Psi^* \otimes P_\Psi) (\psi \otimes \psi^* \otimes \psi)$$

$$\rho \approx P_\rho q$$

# Beyond POD

**What if the snapshots aren't of the state vector?**

Taking Gross Pitaevskii as an example, with  $\{\rho(t_i)\} \rightarrow P_\rho$

$$\psi(t_0) = \underline{\psi}_0$$

$$\frac{d\psi}{dt} = \underline{\tilde{H}_0}\psi + g\underline{\mathbb{T}_\Psi^3}(\psi \otimes \psi^* \otimes \psi)$$

$$q = P_\rho^\dagger \rho \approx \underline{B_\rho^\Psi \mathbb{T}_\Psi^2}(\psi \otimes \psi^*)$$

$$\rho \approx P_\rho q$$

# Multiple snapshot sets:

## Multiple snapshot sets:

Navier-Stokes as an example has three velocity fields

$$\mathbf{u} = \begin{bmatrix} u_x \\ u_y \\ u_z \end{bmatrix}$$

## Multiple snapshot sets:

Navier-Stokes as an example has three velocity fields

$$\mathbf{u} = \begin{bmatrix} u_x \\ u_y \\ u_z \end{bmatrix}$$

and the advective term

$$(\mathbf{u} \cdot \nabla) \mathbf{u} = \begin{bmatrix} u_x \odot (\partial_x u_x) + u_y \odot (\partial_y u_x) + u_z \odot (\partial_z u_x) \\ u_x \odot (\partial_x u_y) + u_y \odot (\partial_y u_y) + u_z \odot (\partial_z u_y) \\ u_x \odot (\partial_x u_z) + u_y \odot (\partial_y u_z) + u_z \odot (\partial_z u_z) \end{bmatrix} \stackrel{i}{=} u_j \odot (\partial_j u_i)$$

## Multiple snapshot sets:

Navier-Stokes as an example has three velocity fields

$$\mathbf{u} = \begin{bmatrix} u_x \\ u_y \\ u_z \end{bmatrix}$$

and the advective term

$$(\mathbf{u} \cdot \nabla) \mathbf{u} = \begin{bmatrix} u_x \odot (\partial_x u_x) + u_y \odot (\partial_y u_x) + u_z \odot (\partial_z u_x) \\ u_x \odot (\partial_x u_y) + u_y \odot (\partial_y u_y) + u_z \odot (\partial_z u_y) \\ u_x \odot (\partial_x u_z) + u_y \odot (\partial_y u_z) + u_z \odot (\partial_z u_z) \end{bmatrix} \stackrel{i}{=} u_j \odot (\partial_j u_i)$$

$$\xrightarrow{i} \underline{B_j^{i\dagger}} \underline{\mathbb{T}_j^2} \left( v_j \otimes \underline{D_j B_j^i} v_i \right)$$

## Equation-Free Methods

## Equation-Free Methods

- ▶ **Dynamic  
Mode  
Decomposition**

## Equation-Free Methods

- ▶ **Dynamic  
Mode  
Decomposition**
- ▶ **Sparse  
Identification of  
Nonlinear  
Dynamics**

## Equation-Free Methods

- ▶ **Dynamic  
Mode  
Decomposition**
- ▶ **Sparse  
Identification of  
Nonlinear  
Dynamics**
- ▶ **Neural  
Operators**

## Equation-Free Methods

- ▶ Dynamic Mode Decomposition
- ▶ Sparse Identification of Nonlinear Dynamics
- ▶ Neural Operators

## Equation-Full Methods

## Equation-Free Methods

- ▶ Dynamic Mode Decomposition
- ▶ Sparse Identification of Nonlinear Dynamics
- ▶ Neural Operators

## Equation-Full Methods

- ▶ Proper Orthogonal Decomposition

## Equation-Free Methods

- ▶ Dynamic Mode Decomposition
- ▶ Sparse Identification of Nonlinear Dynamics
- ▶ Neural Operators

## Equation-Full Methods

- ▶ Proper Orthogonal Decomposition
- ▶ Eigenvector Continuation

# Parametric Matrix Models

## Equation-Free Methods

- ▶ Dynamic Mode Decomposition
- ▶ Sparse Identification of Nonlinear Dynamics
- ▶ Neural Operators

## Equation-Full Methods

- ▶ Proper Orthogonal Decomposition
- ▶ Eigenvector Continuation

# Parametric Matrix Models

## Equation-Free Methods

- ▶ Dynamic Mode Decomposition
- ▶ Sparse Identification of Nonlinear Dynamics
- ▶ Neural Operators

## Equation-Full Methods

- ▶ Proper Orthogonal Decomposition
- ▶ Eigenvector Continuation

## Equation-Informed Methods

# Parametric Matrix Models

## Equation-Free Methods

- ▶ Dynamic Mode Decomposition
- ▶ Sparse Identification of Nonlinear Dynamics
- ▶ Neural Operators

## Equation-Full Methods

- ▶ Proper Orthogonal Decomposition
- ▶ Eigenvector Continuation

## Equation-Informed Methods

## Form-Full Methods

# Summary

# Summary

- ▶ Practical high-level overview to making a PMM

$$H(c) = H_0 + cH_1$$



$$M(c) = \underline{M_0} + c\underline{M_1}$$



# Summary

- ▶ Practical high-level overview to making a PMM
- ▶ Highlighted previous results in emulation

$$H(c) = H_0 + cH_1$$



$$M(c) = \underline{M}_0 + c\underline{M}_1$$






# Summary

- ▶ Practical high-level overview to making a PMM
- ▶ Highlighted previous results in emulation
- ▶ Teased ongoing work in model discovery

$$H(c) = H_0 + cH_1$$

↓

$$M(c) = \underline{M}_0 + c\underline{M}_1$$






# Summary

- ▶ Practical high-level overview to making a PMM
- ▶ Highlighted previous results in emulation
- ▶ Teased ongoing work in model discovery
- ▶ Detailed new theory for extending PMMs

$$H(c) = H_0 + cH_1$$

↓

$$M(c) = \underline{M}_0 + c\underline{M}_1$$



$$\underline{T}_{\Psi}^3 \quad \underline{T}_{jkl} \quad \underline{B}_{\rho}^{\Psi} \quad \dots$$

# References

- Cohen, Gregory, Saeed Afshar, et al. (2017). arXiv: 1702.05373 [cs.CV].
- Cook, Patrick, Danny Jammoooa, et al. (2024). <https://arxiv.org/abs/2401.11694>. arXiv: 2401.11694v5 [cs.LG].
- De, Arinjoy, Patrick Cook, et al. (2023). arXiv: 2309.10856 [quant-ph].
- Fein-Ashley, Jacob, Tian Ye, et al. (2024). arXiv: 2402.00564v1 [cs.CV].
- Frame, Dillon, Rongzheng He, et al. (2018). In: *Phys. Rev. Lett.* 121 (3), p. 032501. DOI: 10.1103/PhysRevLett.121.032501.
- Gamito, J., J. Khalouf-Rivera, et al. (2022). In: *Physical Review E* 106.4. DOI: 10.1103/physreve.106.044125.
- Granmo, Ole-Christoffer, Sondre Glimsdal, et al. (2019). arXiv: 1905.09688 [cs.LG].
- Jin, Lauren, Ante Ravlić, et al. (2024). In: *Bulletin of the American Physical Society*.
- Jin, Yingyezhe, Wenrui Zhang, et al. (2018). In: *Proceedings of the 32nd International Conference on Neural Information Processing Systems*. Curran Associates Inc., pp. 7005–7015. DOI: 10.5555/3327757.3327804.
- Kabir, H. M. D., Moloud Abdar, et al. (2020). In: *IEEE Transactions on Artificial Intelligence* 4, pp. 1165–1177.
- König, S., A. Ekström, et al. (2020). In: *Physics Letters B* 810, p. 135814. DOI: <https://doi.org/10.1016/j.physletb.2020.135814>.
- Kramer, Boris and Karen E. Willcox (2019). In: *AIAA Journal* 57.6, pp. 2297–2307. DOI: 10.2514/1.j057791.
- Lecun, Y., L. Bottou, et al. (1998). In: *Proceedings of the IEEE* 86.11, pp. 2278–2324. DOI: 10.1109/5.726791.
- Mazzia, Vittorio, Francesco Salvetti, et al. (2021). In: *Scientific Reports* 11.1. DOI: 10.1038/s41598-021-93977-0.
- Nøkland, Arild and Lars Hiller Eidnes (2019). arXiv: 1901.06656 [stat.ML].
- Pishchik, Evgenii (2023). DOI: 10.20944/preprints202301.0463.v1.
- Reed, Brendan T., Rahul Somasundaram, et al. (2024). arXiv: 2405.20558 [astro-ph.HE].
- Sarkar, Avik and Dean Lee (2021). In: *Phys. Rev. Lett.* 126 (3), p. 032501. DOI: 10.1103/PhysRevLett.126.032501.
- Somasundaram, Rahul, Cassandra L. Armstrong, et al. (2024). arXiv: 2404.11566 [nucl-th].
- Somasundaram, Rahul, Isak Svensson, et al. (2024). arXiv: 2410.00247 [nucl-th].
- Tanveer, M., M. Karim Khan, et al. (2021). In: *2020 25th International Conference on Pattern Recognition (ICPR)*. IEEE Computer Society, pp. 4789–4796. DOI: 10.1109/ICPR48806.2021.9412221.
- Xiao, Han, Kashif Rasul, et al. (2017). arXiv: 1708.07747 [cs.LG].

# General Machine Learning

# General Machine Learning

## Underlying Equations

???

# General Machine Learning

## Underlying Equations

???

## PMM

$$H(\vec{c}) = \underline{H_0} + c_1 \underline{H_1} + \cdots + c_p \underline{H_p}$$

# General Machine Learning

## Underlying Equations

???

## PMM

$$\begin{aligned} H(\vec{c}) &= \underline{H_0} + c_1 \underline{H_1} + \cdots + c_p \underline{H_p} \\ H(\vec{c}) |E_k(\vec{c})\rangle &= E_k(\vec{c}) |E_k(\vec{c})\rangle \end{aligned}$$

# General Machine Learning



## Underlying Equations

???




## PMM

$$\begin{aligned} H(\vec{c}) &= \underline{H_0} + c_1 \underline{H_1} + \cdots + c_p \underline{H_p} \\ H(\vec{c}) |E_k(\vec{c})\rangle &= E_k(\vec{c}) |E_k(\vec{c})\rangle \\ z_i(\vec{c}) &\sim |\langle E_a(\vec{c}) | \underline{O_i} | E_b(\vec{c}) \rangle|^2 \end{aligned}$$





# Discontinuous Regression



# Discontinuous Regression






# Discontinuous Regression



















$$\sum_{nm} a_{nm} \sin(nx) \sin(my)$$





$$\sum_{nm} a_{nm} \sin(nx) \sin(my)$$






$$\sum_{nm} a_{nm} \sin(nx) \sin(my)$$






$$\sum_{nm} a_{nm} \sin(nx) \sin(my)$$



# Image Classification


# Image Classification








97.67% Train Accuracy




97.38% Test Accuracy



# MNIST Digits


Model	Acc. (%)	float
PMM <sup>†</sup>	97.38	4990
ConvPMM	99.10	129 416
DNN-2 <sup>†</sup> [Pishchik 2023]	96.5	~ 311 650
DNN-3 <sup>†</sup> [Pishchik 2023]	97.0	~ 386 718
DNN-5 <sup>†</sup> [Pishchik 2023]	97.2	~ 575 050
GECCO [Fein-Ashley, Ye, et al. 2024]	98.04	~ 19 000
CTM-250 [Granmo, Glimsdal, et al. 2019]	98.82	31 750
CTM-8000 [Granmo, Glimsdal, et al. 2019]	99.4	527 250
Eff.-CapsNet [Mazzia, Salvetti, et al. 2021]	99.84	161 824

† non-convolutional






90.66% Train Accuracy




88.58% Test Accuracy




# Fashion MNIST


Model	Acc. (%)	float
PMM <sup>†</sup>	88.58	16 744
ConvPMM	90.94	278 280
GECCO [Fein-Ashley, Ye, et al. 2024]	88.09	~ 19 000
CTM-250 [Granmo, Glimsdal, et al. 2019]	88.25	31 750
CTM-8000 [Granmo, Glimsdal, et al. 2019]	91.5	527 250
MLP <sup>†</sup> [Nøkland and Eidnes 2019]	91.63	2 913 290
VGG8B [Nøkland and Eidnes 2019]	95.47	~ 7 300 000
F-T DARTS [Tanveer, Khan, et al. 2021]	96.91	~ 3 200 000

† non-convolutional






83.45% Train Accuracy



13792 trainable floats

81.57% Test Accuracy



# EMNIST

Model	Acc. (%)	float
PMM <sup>†</sup>	81.57	13 792
ConvPMM	85.95	349 172
CNN [Kabir, Abdar, et al. 2020]	79.61	21 840
CNN (S-FC) [Kabir, Abdar, et al. 2020]	82.77	13 820
CNN (S-FC) [Kabir, Abdar, et al. 2020]	83.21	16 050
HM2-BP <sup>†</sup> [Y. Jin, Zhang, et al. 2018]	85.57	665 647

† non-convolutional

