
## On Schemes for Exponential Decay

### Hans Petter Langtangen<sup>1,2</sup>

Center for Biomedical Computing, Simula Research Laboratory<sup>1</sup>

Department of Informatics, University of Oslo<sup>2</sup>

Jun 29, 2013



### Goal

The primary goal of this demo talk is to demonstrate how to write talks with doconce and get them rendered in numerous HTML formats.

### Layout.

This version utilizes beamer slides with the theme simula.

# Mathematical problem

$$u'(t) = -au(t), \tag{1}$$

$$u(0) = I, (2)$$

- $t \in (0, T]$
- a, I, and T are prescribed parameters
- ightharpoonup u(t) is the unknown function



## **Numerical solution method**

- ▶ Mesh in time:  $0 = t_0 < t_1 \cdots < t_N = T$
- Assume constant  $\Delta t = t_n t_{n-1}$
- $\triangleright$   $u^n$ : numerical approx to the exact solution at  $t_n$

#### Numerical scheme:

$$u^{n+1} = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}u^n, \quad n = 0, 1, \dots, N - 1$$

### **Numerical solution method**

- ▶ Mesh in time:  $0 = t_0 < t_1 \cdots < t_N = T$
- Assume constant  $\Delta t = t_n t_{n-1}$
- $\triangleright$   $u^n$ : numerical approx to the exact solution at  $t_n$

Numerical scheme:

$$u^{n+1} = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}u^n, \quad n = 0, 1, \dots, N - 1$$

## **Numerical solution method**

- ▶ Mesh in time:  $0 = t_0 < t_1 \cdots < t_N = T$
- Assume constant  $\Delta t = t_n t_{n-1}$
- $\triangleright$   $u^n$ : numerical approx to the exact solution at  $t_n$


Numerical scheme:

$$u^{n+1} = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}u^n, \quad n = 0, 1, \dots, N - 1$$

## **Implementation**

The numerical method is implemented in a Python function:

## The Crank-Nicolson method



Exact solution of the scheme:

$$u^n = A^n$$
,  $A = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}$ 

- Stability: |A| < 1</p>
- No oscillations: A > 0
- Always for Backward Euler ( $\theta = 1$ )
- ▶  $\Delta t < 1/a$  for Forward Euler ( $\theta = 0$ )
- ▶  $\Delta t < 2/a$  for Crank-Nicolson ( $\theta = 1/2$ )

#### Concluding remarks:

Exact solution of the scheme:

$$u^n = A^n$$
,  $A = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}$ .

- Stability: |A| < 1</p>
- No oscillations: A > 0
- ▶ Always for Backward Euler ( $\theta = 1$ )
- ▶  $\Delta t < 1/a$  for Forward Euler ( $\theta = 0$ )
- ▶  $\Delta t < 2/a$  for Crank-Nicolson ( $\theta = 1/2$ )

#### Concluding remarks:

Exact solution of the scheme:

$$u^n = A^n$$
,  $A = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}$ .

- ► Stability: |A| < 1
- ► No oscillations: *A* > 0
- ▶ Always for Backward Euler ( $\theta = 1$ )
- ▶  $\Delta t$  < 1/a for Forward Euler ( $\theta$  = 0)
- ▶  $\Delta t$  < 2/a for Crank-Nicolson ( $\theta$  = 1/2)

### Concluding remarks:

Exact solution of the scheme:

$$u^n = A^n$$
,  $A = \frac{1 - (1 - \theta)a\Delta t}{1 + \theta a\Delta t}$ .

- ► Stability: |A| < 1
- No oscillations: A > 0
- ▶ Always for Backward Euler ( $\theta = 1$ )
- ▶  $\Delta t < 1/a$  for Forward Euler ( $\theta = 0$ )
- $\Delta t < 2/a$  for Crank-Nicolson ( $\theta = 1/2$ )

#### Concluding remarks: