

Dual-lift realizability for Keller-type maps: structural incompleteness, schema, and examples

Daniel Eric Fredriksen
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

22 July 2026

Abstract

We isolate a structural lemma for dual vector fields of local diffeomorphisms $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$: if global dual fields X_j with $DF \cdot X_j = e_j$ are all complete, then F is surjective. Hence any non-surjective dual system has at least one incomplete dual field. We propose a classification schema (B-incomplete / B-ESS / B-Def / B-CCR / B-CP) and populate it on atlas rows A000–A010 (pass and fail baselines, Keller seed A001, family pilot A002, a smooth 2D pair A007/A008 illustrating that the forced incomplete set can be as thin as measure zero, and a polynomial 2D pair A009/A010: Pinchuk’s real-plane map with *open-set* incompleteness, and a polynomial shear automorphism with complete dual fields). We prove restricted dichotomy lemmas: polynomial product local diffeos have complete dual fields; the A007 zero-dimensional asymptotic mechanism is impossible for polynomials (Jelonek); in dimension 2 dual incompleteness is fiber-saturated and falls into empty / open / residual-thin regimes. Residual-thin regime T is excluded on proper, injective, degree-one-component, product, triangular, and graph-type (including all $\deg P \leq 2$) classes, and reduced to incompleteness only on atypical fibers of non-coordinate P of degree ≥ 3 ; full exclusion of T remains open. No gates, channels, or computational advantage are claimed. Companion pack: <https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space/blob/main/docs/notes/COMPANION-PACK.md>.

MSC: 37C10, 47B25, 53D17, 14R15.

Version: B001 draft v0.7 (erratum: real poly local-diffeo language — global C^∞ diffeo / regime E; polynomial inverse only when proved; supersedes freeze 61bf3da automorphism wording).

Software: atlas program_b; CAS anchors:

cas_atlas_A005_A006_B001.json, cas_atlas_A007_A008_B001.json,
cas_atlas_A009_A010_B001.json, cas_atlas_B5_poly_dichotomy_B001.json,
cas_atlas_B6_regime_T_B001.json, cas_atlas_B7_OPEN_T_B001.json.

1 Introduction

Cotangent lifts and dual fields are standard when $\det DF \neq 0$. After the A001 deficiency analysis [1], we classify which maps force incomplete dual fields and how that interacts with ESS / CCR / CP packages [2]. A001 is awaiting math-ph endorsement (code VIPN6B); no arXiv id is assigned yet, but the concept DOI already cites A001.

2 Structural lemma

Lemma 1 (Complete dual fields imply surjectivity). *Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be C^∞ and a local diffeomorphism with dual fields X_j , $DF \cdot X_j = e_j$. If every X_j is complete, then $F(\mathbb{R}^n) = \mathbb{R}^n$.*

Proof. $F(\phi_t^{(j)}(q)) = F(q) + te_j$; iterate; image open and translation-invariant. $\square$

Corollary 1. *Non-surjective dual systems have some incomplete X_j .*

3 Schema

B-incomplete / B-ESS / B-Def / B-CCR / B-CP with values **open** / **fail** / **pass**.

4 Atlas scoreboard (v0.4)

ID	Map (sketch)	B-inc.	Role
A000	id	pass	baseline
A004	shear $(x, y + x, z)$	pass	linear auto
A005	diag $(2, \frac{1}{2}, 1)$	pass	linear stretch (CAS)
A008	$(x, y + \sin x, z)$	pass	2D smooth global diffeo (CAS)
A010	$(x, y + x^2, z)$	pass	2D <i>polynomial</i> auto (CAS)
A003	(x, e^y) pad	fail	smooth B1 example
A006	(e^x, e^y, z)	fail	two incomplete fields, open set (CAS)
A007	$z \mapsto e^z$ pad	fail	incomplete on <i>null set only</i> (CAS)
A009	Pinchuk (P, Q) pad	fail	poly, incomplete on <i>open set</i> (CAS)
A001	Keller seed	fail	$X_1 + H_1$ analytic [1]
A002	family d4 pilot	open	geometry pending

5 Examples (new in v0.2)

Example 1 (A005 — linear stretch, pass-like). $F = \text{diag}(2, \frac{1}{2}, 1)$, $\det DF = 1$. Dual fields $X_0 = \frac{1}{2}\partial_x$, $X_1 = 2\partial_y$, $X_2 = \partial_z$ are complete. CAS: `cas_atlas_A005_A006_B001.json`.

Example 2 (A006 — exp quadrant, fail-like). $F(x, y, z) = (e^x, e^y, z)$, $\det DF = e^{x+y} > 0$, image $(0, \infty)^2 \times \mathbb{R}$. Dual fields $X_0 = e^{-x}\partial_x$, $X_1 = e^{-y}\partial_y$ incomplete backward ($e^x = e^{x_0} + t$). Matches Lemma 1. CAS same file.

6 Examples (new in v0.3): two incompleteness regimes

Example 3 (A007 — complex exponential, thin fail-like). $F(x, y, z) = (e^x \cos y, e^x \sin y, z)$, i.e. $z \mapsto e^z$ padded. Conformal, $\det DF = e^{2x} > 0$; image misses the origin (non-surjective). Writing $w = e^{x+iy}$, the dual field X_0 solves $\dot{w} = 1$, so $w(t) = w(0) + t$. If $\text{Im } w(0) \neq 0$ (full measure), the orbit never reaches $w = 0$: **complete**. If $\text{Im } w(0) = 0$ (a measure-zero pair of rays), $w(t)$ hits 0 at finite t and $x(t) \rightarrow -\infty$: **incomplete**. Thus Lemma 1's corollary is satisfied by an incomplete set of measure zero, sharply contrasting A006/A001 where incompleteness occupies an open set. CAS: `cas_atlas_A007_A008_B001.json`.

Example 4 (A008 — shear + sine, 2D global diffeomorphism). $F(x, y, z) = (x, y + \sin x, z)$, $\det DF = 1$. *Explicit global inverse* $(x, y) \mapsto (x, y - \sin x)$ (verified by CAS roundtrip). Dual fields $X_0 = \partial_x - \cos x \partial_y$, $X_1 = \partial_y$; the X_0 flow $y(t) = y_0 - \sin(x_0 + t) + \sin(x_0)$ is bounded for all t : **complete**. A genuinely nonlinear, non-triangular-affine 2D coupling (unlike A004/A005) that nonetheless has no incompleteness at all.

7 Examples (new in v0.4): polynomial 2D evidence

Example 5 (A009 — Pinchuk real plane map, open fail-like). Let $t = xy - 1$, $h = t(xt + 1)$, $f = (xt + 1)^2(t^2 + y)$, and

$$P = f + h, \quad Q = -t^2 - 6th(h + 1) - u(f, h),$$

with the Campbell/van den Essen auxiliary $u = 170fh + 91h^2 + 195fh^2 + 69h^3 + 75fh^3 + \frac{75}{4}h^4$ (Pinchuk [3], Campbell [4]). Then $F = (P, Q) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ (padded by idle z) is polynomial of degrees (10, 25), with

$$\det DF = t^2 + (t + f(13 + 15h))^2 + f^2 > 0$$

everywhere (sum of squares; $t = f = 0$ is impossible over $\mathbb{R}$). It is not injective (CAS-certified collisions) and not surjective (misses exactly two points [4]). The dual field $X_1 = H(P)/j = (-P_y, P_x)/j$ satisfies $DF \cdot X_1 = e_1$ and $\dot{Q} = 1$. On every level $P = c$ with $c > -1$, the real level curve has poles $h = -1 \pm \sqrt{1 + c}$ at which $(x, y) \rightarrow \infty$ while Q approaches a finite asymptotic value: escape in finite X_1 -time. Hence the incompleteness locus of X_1 contains the open set $\{P > -1\}$.

Open-set incompleteness for a genuinely polynomial 2D Keller-type map, contrasting A007's thin failure. CAS: `cas_atlas_A009_A010_B001.json`.

Example 6 (A010 — polynomial shear automorphism, pass-like). $F(x, y, z) = (x, y + x^2, z)$, $\det DF = 1$, inverse $(x, y - x^2, z)$. Dual fields $X_0 = \partial_x - 2x \partial_y$, $X_1 = \partial_y$; the X_0 flow $y(t) = y_0 - 2x_0t - t^2$ is polynomial in t : **complete**. A polynomial pass baseline paralleling A008.

8 Toward a dichotomy (open)

A007 shows the incomplete locus forced by Lemma 1 can be as thin as measure zero for general smooth local diffeomorphisms. A007 is *not* polynomial. A009 (Pinchuk) supplies the first atlas polynomial 2D non-injective example with *open-set* incompleteness, consistent with A001's open failures in dimension 3 and with A010 as a polynomial complete baseline. The candidate claim — polynomial Keller-type maps never exhibit A007-style thin incompleteness — remains open as a *full* theorem, but is reduced by the restricted lemmas of §9.

9 Restricted dichotomy lemmas (new in v0.5)

Full proofs: PROGRAM-B-B5-poly-dichotomy-lemmas.md. CAS: `cas_atlas_B5_poly_dichotomy_B001.json`.

Lemma 2 (Univariate polynomial local diffeos are global). *If $f \in \mathbb{R}[t]$ satisfies $f'(t) \neq 0$ for all $t \in \mathbb{R}$, then $f : \mathbb{R} \rightarrow \mathbb{R}$ is a global diffeomorphism.*

Proof. f' has constant strict sign, so f is strictly monotone. Even degree is impossible (equal infinite limits $\Rightarrow$ not injective). Odd degree gives opposite infinite limits, hence bijectivity; inverse is C^∞ by IFT. $\square$

Lemma 3 (Polynomial product maps have complete duals). *If $F(x) = (f_1(x_1), \dots, f_n(x_n))$ with each $f_i \in \mathbb{R}[t]$ and $\prod_i f'_i \neq 0$, then F is a global diffeomorphism and every dual field $X_i = (1/f'_i)\partial_{x_i}$ is complete.*

Proof. Apply Lemma 2 coordinatewise; dual ODEs integrate by $f_i(x_i(t)) = f_i(x_i(0)) + t$. $\square$

Lemma 4 (Cylinder criterion). *If $\text{int } \pi_j(\mathbb{R}^n \setminus F(\mathbb{R}^n)) \neq \emptyset$, then the incompleteness locus of X_j contains a nonempty open set.*

Proof. On the open set of domain points whose F -image projects into that interior, the straight-line orbit in the target meets the complement of the image in finite time, so the dual lift is incomplete. $\square$

Lemma 5 (2D fiber saturation and three regimes). *Let $F = (P, Q) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be polynomial with $\det DF \neq 0$. The incompleteness locus U_1 of the dual field with $\dot{Q} = 1$ is a union of connected components of levels of P . The set of bad level values is semi-algebraic in $\mathbb{R}$, hence a finite union of points and open intervals. Thus either $U_1 = \emptyset$ (regime E; e.g. A010), or U_1 contains a nonempty open set (regime O; e.g. A009), or U_1 is a finite union of real algebraic curves (regime T, measure zero) — the only residual thin channel.*

Proof sketch. No critical points of P ; levels are unions of lines; Q is a chart on each component with speed 1; completeness iff Q hits all of $\mathbb{R}$ on that component. Semi-algebraicity of the bad-value set is Tarski–Seidenberg. $\square$

Lemma 6 (Jelonek blocks the A007 asymptotic mechanism). *If $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is polynomial and generically finite and the asymptotic set S_F is nonempty, then $\dim S_F = n - 1$ [5, 6]. In particular S_F cannot be a nonempty finite set of points, so the A007 pattern (single asymptotic value, thin cylindrical projection, null incomplete locus) is unavailable to polynomials.*

Restricted package. Lemmas 2–6 eliminate thin incompleteness for polynomial *product* maps and ban the A007 mechanism for all polynomial local diffeos. Full exclusion of regime T in Lemma 5 remains open; no atlas example realizes T.

10 Regime T partial exclusion (new in v0.6)

Full proofs: PROGRAM-B-B6-regime-T.md. CAS: cas_atlas_B6_regime_T_B001.json.

Lemma 7 (Chamber reduction for regime T). *For polynomial $F = (P, Q)$ as in Lemma 5, Hardt chambers have constant Q -end type. Hence regime T occurs if and only if every open Hardt chamber of P is good and $\text{Bad}_P(F)$ is a nonempty finite subset of the atypical values of P .*

Lemma 8 (Structured exclusions of regime T). *Regime T is impossible for a polynomial local diffeomorphism $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ in any of the following classes:*

1. F proper (then F is a global C^∞ diffeomorphism; duals complete);
2. F injective (then F is a global C^∞ diffeomorphism; duals complete; polynomial inverse is not automatic over $\mathbb{R}$);
3. one component has total degree 1 (after affine domain change the map is a global C^∞ diffeomorphism);

4. product maps as in Lemma 3, or triangular maps $F = (f(x), \alpha(x)y + \beta(x))$ with f', α never zero.

Proof sketch. (1) Proper local diffeos $\mathbb{R}^n \rightarrow \mathbb{R}^n$ are coverings, hence global diffeos. (2) Injective polynomial local diffeomorphisms of $\mathbb{R}^n$ are bijective global C^∞ diffeomorphisms (IFT + surjectivity of injective real polynomial maps); dual completeness as in the proper case. Over $\mathbb{R}$, the inverse need not be polynomial — e.g. $F(x, y) = (y, x + x^3)$. (Algebraic automorphism theorems over algebraically closed fields do not supply a free poly inverse over $\mathbb{R}$.) (3) Reduce to $P = x$; then $Q_y \neq 0$ forces each univariate $y \mapsto Q(u, y)$ to be a global diffeo by Lemma 2, yielding an explicit global C^∞ inverse. (4) Lemmas 3 and B5.7. $\square$

Lemma 9 (Model atypical submersion is not a coordinate). *Let $P_0 = x + x^2y$. Then $\nabla P_0 \neq 0$ everywhere and 0 is atypical, but there is no $Q \in \mathbb{R}[x, y]$ with $\det D(P_0, Q)$ a nonzero constant. Moreover every Q that is linear in y has $\det D(P_0, Q)$ vanishing somewhere; CAS finds no Q of total degree ≤ 6 with Jacobian equal to any of $\pm 1, x^2 + 1, y^2 + 1, x^2 + y^2 + 1, (x^2 + 1)(y^2 + 1), (x^2 + 1)^2, (y^2 + 1)^2$.*

Partial T package. Lemmas 7–9 exclude T on structured subclasses and block the standard atypical first-component model at low degree. The residual **OPEN-T** asks whether any polynomial local diffeo has Bad_P a nonempty finite subset of $\text{Atyp}(P)$ with all Hardt chambers good. No atlas example realizes T; full dichotomy remains a conjecture.

11 OPEN-T sharpened: graph type and $\deg P \leq 2$ (new in v0.7)

Full proofs: PROGRAM-B-B7-OPEN-T.md. CAS: cas_atlas_B7_OPEN_T_B001.json.

Affine invariance. Affine domain/codomain changes preserve $\det DF \neq 0$, polynomial class, dual completeness (affine conjugacy of flows), and the E/O/T regime labels; the graph-type normal form and $\deg P \leq 2$ reduction are therefore non-vacuous inside the same package (PROGRAM-B-B7-OPEN-T.md, Remark B7.0a).

Lemma 10 (Graph-type first component $\Rightarrow$ global C^∞ diffeo). *Call P graph type if after an affine domain automorphism, $P = y + f(x)$ for some $f \in \mathbb{R}[x]$. If $F = (P, Q)$ is polynomial with $\det DF \neq 0$ and P graph type, then on each fiber $y = c - f(x)$ the restriction $q_c(x) = Q(x, c - f(x))$ is a strictly monotone univariate polynomial (derivative $-j \neq 0$), hence a C^∞ diffeomorphism $\mathbb{R} \rightarrow \mathbb{R}$. Solving fiberwise yields a global C^∞ inverse, so F is a global C^∞ diffeomorphism of $\mathbb{R}^2$ (a polynomial map that is a C^∞ automorphism of $\mathbb{R}^2$) and $\text{Bad}_P = \emptyset$ (regime E). Regimes O and T are impossible. The inverse need not be polynomial in general (e.g. $Q = x + x^3/3 + g(P)$); when fiberwise q_c are degree one or an explicit polynomial inverse is available (e.g. elementary shears), F is a polynomial automorphism — assert that stronger label only then.*

Theorem 1 (Degree ≤ 2 first component). *If $F = (P, Q)$ is a polynomial local diffeomorphism and $\deg P \leq 2$, then after an affine domain change P is graph type (the Hessian of a critical-point-free quadratic is rank one), hence F is a global C^∞ diffeomorphism of $\mathbb{R}^2$ and regime E holds. In particular regime T requires $\deg P \geq 3$. (Polynomial inverse only when separately proved; see Lemma 10.)*

Lemma 11 (P_0 -axis cannot carry T). *For $P_0 = x + x^2y$, any Q with $\{P_0, Q\} \neq 0$ everywhere makes the atypical axis component $\{x = 0\} \subset P_0^{-1}(0)$ good (univariate diffeomorphism in y). Thus T cannot be realized by “only the axis is bad.” Odd vertical degree forces the leading coefficient freeze $a_n = Cx^{2n}$; the linear-in- y case collapses as in Lemma 9. CAS finds no Q of total degree ≤ 8 with Jacobian equal to standard never-zero targets (including $(x^2 + y^2 + 1)^2$).*

Sharpened OPEN-T. Lemmas 10–11 and Theorem 1 exclude T on all graph-type / $\deg P \leq 2$ maps and filter residual candidates to non-coordinate P of degree ≥ 3 with badness on bifurcation-born components. Full exclusion and existence both remain open; no atlas example realizes T.

12 Prior examples (v0.1–v0.2)

A000 identity; A003 half-plane; A001 Keller seed with X_1 incomplete and $H = -iX_1$ deficiency (∞, ∞) in [1] (other X_j incompleteness is recorded in companion validation dossiers, not re-proved here); A002 open; A004 shear pass.

13 Non-claims

No gates, channels, advantage. No dual-flow repair. A003/A005/A006/A007/A008 are not Jacobian counterexamples (A007/A008 are not even polynomial). A009 is a classical strong-real-JC counterexample ($\det > 0$, not injective) but not a complex Jacobian conjecture counterexample ($\det$ not constant; JC(2) over $\mathbb{C}$ remains open). A010 is a polynomial shear automorphism (explicit poly inverse). T4: no dual F -translation package claimed here. The full polynomial-only dichotomy remains a conjecture (§9–§11): regime T of Lemma 5 is excluded on structured classes (Lemma 8) and on graph-type / $\deg P \leq 2$ maps (Theorem 1, as global C^∞ diffeos / regime E); residual OPEN-T is named in §11. Prior freeze 61bf3da wording that concluded “polynomial automorphism” from bijective real poly local diffeos without a poly-inverse proof is superseded; regime-E chains are unchanged. C’s joint q - p sector [2] (v0.9) splits: position CONSTRUCT (Koopman); J2 Joint-Stone-Canonical OBSTRUCTION; J3 full-triple Joint-Stone **G4-conditional** OBSTRUCTION; J4 Joint-Stone-Hom-1 OBSTRUCTION; J5 mere-CP Joint-Stone-CP-1 OBSTRUCTION; J6 Unitary-Image-CP-1 OBSTRUCTION; J6-C Diag-CP- Φ_0 CONSTRUCT; Joint-Form-Core CONSTRUCT / Joint-Form-ESS-1 OBSTRUCT; J7-C Full- ψ -BT-Envelope (bounded dual-momentum transforms + Φ_0 positions; form-level only, not CFC / not Weyl- C^* / not Stinespring) CONSTRUCT; J7 Full- ψ -CFC-SA-1 OBSTRUCTION (via J6); Full- ψ -CP-Weyl- C^* OPEN (narrow).

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., v0.3.6-submit / concept DOI 10.5281/zenodo.21474351, <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.
- [2] D. E. Fredriksen, CP readings of an exotic Weyl endomorphism..., C001 companion v0.9 (J7 SPLIT: Full- ψ -BT-Envelope form-level CONSTRUCT; CFC-SA NO-GO via J6; Weyl- C^* OPEN), <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.
- [3] S. Pinchuk, A counterexample to the strong real Jacobian conjecture, Math. Z. 217 (1994), 1–4.
- [4] L. A. Campbell, The asymptotic variety of a Pinchuk map as a polynomial curve, Appl. Math. Lett. 24 (2011), 62–65; arXiv:1001.3318.
- [5] Z. Jelonek, The set of points at which a polynomial map is not proper, Ann. Polon. Math. 58 (1993), 259–266.
- [6] Z. Jelonek, Testing sets for properness of polynomial mappings, Math. Ann. 315 (1999), 1–12.