

Poisson and Weyl lifts of the Alpöge–Fable Keller
map
and nonunique self-adjoint realizations of a dual
transport operator

Daniel Eric Fredriksen

Quantyra Inc.

<https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

We study the three-dimensional Keller map F announced by Alpöge (with Fable credited): $\det DF \equiv -2$ and a three-point collision. Theorem A restates those finite identities; novelty is Theorems B–F. The cotangent lift $\Phi(q, p) = (F(q), J^{-T}p)$ is a polynomial Poisson map and non-injective. Dual fields $X_j = \sum_k B_{jk} \partial_{q_k}$ commute and are divergence-free (Piola), so $H_j = -iX_j$ are symmetric on $C_c^\infty(\mathbb{R}^3)$. The field X_1 is incomplete. In flow-box coordinates $(a, s, c) = F(q)$ one has $X_1 = \partial_s$ and $dq = \frac{1}{2} da ds dc$. Open forward and backward escape walls yield whole-orbit deficiency vectors in $\text{Dom}(H^*)$ (transverse cutoffs only; no interior s -indicators), hence

$$(n_+, n_-) = (\infty, \infty)$$

for $H = -iX_1$. Infinitely many self-adjoint extensions exist; algebraic CCR/Poisson data select none. No gates, channels, computational advantage, or seed discovery claim.

MSC 2020: 47B25, 81S05, 14R15, 37C10, 53D17.

Software: v0.3.3; concept DOI 10.5281/zenodo.21474351 (version DOI on Zenodo ingest).

Errata. v0.2.2 pair $(0, \infty)$ withdrawn. v0.3.0 interior s -indicator deficiency proof withdrawn ($\notin \text{Dom}(H^*)$).

1 Introduction

On 19–20 July 2026, Levent Alpöge announced $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ with $\det DF \equiv -2$ and a three-point collision (credits Fable) [2]. **We do not claim discovery of F .** We ask whether algebraic Poisson/CCR lifts select a unique self-adjoint dual transport generator on $C_c^\infty(\mathbb{R}^3)$. Answer: no for $H = -iX_1$; deficiency indices (∞, ∞) .

2 Setup

$$\begin{aligned} F_0 &= (1 + xy)^3 z + y^2(1 + xy)(4 + 3xy), \\ F_1 &= y + 3x(1 + xy)^2 z + 3xy^2(4 + 3xy), \\ F_2 &= 2x - 3x^2 y - x^3 z. \end{aligned}$$

$$B = J^{-T}, \quad X_j = \sum_k B_{jk} \partial_{q_k}, \quad H := -iX_1.$$

3 Theorems A–D (summary)

Theorem 1 (A). $\det DF \equiv -2$; *three-point collision*; F *noninjective*. (*Restatement; Lean+CAS.*)

Theorem 2 (B–C). Φ *is polynomial Poisson and noninjective*. $X_j(F_i) = \delta_{ij}$, $[X_i, X_j] = 0$, $\operatorname{div} X_j = 0$ (*Piola*), $H_j = -iX_j$ *symmetric*, $[F_i, H_j] = i\delta_{ij}$ *on C_c^∞* .

Theorem 3 (D). X_1 *incomplete*: *explicit γ through $(1, 0, 0)$ with $F(\gamma(t)) = (0, t, 2)$ escapes as $t \rightarrow \frac{1}{2}^-$* .

4 Flow-box coordinates

On local inverse sheets, $(a, s, c) = F(q)$ gives

$$dq = \frac{1}{2} da \, ds \, dc, \quad X_1 = \partial_s, \quad H = -i\partial_s. \quad (7.1)$$

Elimination yields $A(s; a, c)q_0^3 + B(s; c)q_0 + C(c) = 0$ with $A = -cs^3 + s^2 + 18acs - 27a^2c^2 - 16a$, $B = 3cs - 4$, $C = 2c$.

Reconstruction. If $x = q_0$ solves the cubic and denominators are nonzero,

$$q_1 = \frac{9acx^2 - 3csx - 3c - sx^2 + 6x}{x((3cs - 4)x + 3c)}, \quad q_2 = \frac{2x - 3x^2q_1 - c}{x^3},$$

then $F(x, q_1, q_2) = (a, s, c)$.

5 Theorems E–F

Dictionary. On (ℓ, r) : finite upper end $\Rightarrow n_- = 1$ (model e^{s-r}); finite lower end $\Rightarrow n_+ = 1$ (model $e^{-(s-\ell)}$).

Proposition 1 (Forward wall). At $(0, \frac{1}{2}, 2)$, $A = 0$, $\partial_s A = -\frac{1}{2}$, $B = -1$. IFT gives wall β . Set $\tau = \sqrt{\beta - s}$, $q_0 = r/\tau$, and

$$G_+ = \frac{A(\beta - \tau^2)}{\tau^2} r^3 + B(\beta - \tau^2)r + C\tau,$$

extended at $\tau = 0$ by $(-A_s)r^3 + Br$. Simple roots $r = \pm\sqrt{2}$ ($\partial_r G_+ = -2B \neq 0$) continue by IFT; reconstruct by the formulas above.

Proposition 2 (Backward wall). At $a = \frac{1}{54}$, $c = 2$, wall α with $A_s = \frac{1}{6}$, $B = -1$. Set $\tau = \sqrt{s - \alpha}$ and analogous G_- extended by $A_s r^3 + Br$. Simple roots $r = \pm\sqrt{6}$ continue by IFT on an open transverse set.

Lemma 1 (Saturated maximal sheets). Cross-section the selected branch below/above the wall; saturate under the flow of X_1 . Since $X_1 F_1 = 1$, s is time. ODE theory yields a C^∞ diffeomorphism Ψ_+ from

$$D_+ = \{(a, c, s) : (a, c) \in W_+ \subseteq U_+, \ell_+(a, c) < s < \beta(a, c)\}$$

onto an open invariant set $\Omega_+ \subset \mathbb{R}^3$, with $F \circ \Psi_+ = (a, s, c)$. The end $s \rightarrow \beta^-$ is at spatial infinity. If ℓ_+ is finite, maximality forces escape as $s \downarrow \ell_+$ as well. Analogous Ψ_-, Ω_- for the lower wall.

Lemma 2 (Deficiency vectors in $\text{Dom}(H^*)$). For $\chi \in C_c^\infty(W_+)$ define on the entire maximal interval

$$u_-(\Psi_+(a, c, s)) = \chi(a, c) e^{s-\beta(a, c)},$$

and $u_- = 0$ off Ω_+ . No interior s -cutoff. Then

$$\|u_-\|_2^2 = \frac{1}{2} \int_{W_+} |\chi|^2 \int_{\ell_+}^{\beta} e^{2(s-\beta)} ds da dc \leq \frac{1}{4} \|\chi\|_{L^2}^2.$$

For $\varphi \in C_c^\infty(\mathbb{R}^3)$, $\text{supp } \varphi$ meets orbits in compact s -segments strictly inside (ℓ_+, β) . Integration by parts with $\text{div } X_1 = 0$ gives no jump and no endpoint terms, so $u_- \in \text{Dom}(H^*)$ and $(H^* + i)u_- = 0$. Thus $n_- = \infty$. Likewise

$$u_+(\Psi_-(a, c, s)) = \chi(a, c) e^{-(s - \alpha(a, c))}$$

on (α, r_-) yields $n_+ = \infty$.

Theorem 4 (F). $(n_+, n_-) = (\infty, \infty)$.

Proof. The two walls, saturation, and deficiency lemmas; separability. $\square$

Theorem 5 (E). H is not essentially self-adjoint on $C_c^\infty(\mathbb{R}^3)$.

Proof. $n_\pm \geq 1$. $\square$

Corollary 1. Infinitely many self-adjoint extensions exist. Core algebraic Poisson/CCR relations do not distinguish among them.

6 Discussion

The example separates two layers that are often collapsed in practice. On the algebraic side, the Poisson cotangent lift and the dual-field/CCR identities are clean: brackets close, rows of B are divergence-free, and $H_j = -iX_j$ are symmetric on $C_c^\infty(\mathbb{R}^3)$. On the analytic side, incompleteness of X_1 produces open families of finite orbit ends, and the whole-orbit deficiency construction yields

$$(n_+, n_-) = (\infty, \infty).$$

Thus a working algebraic CCR/Poisson presentation does *not* select a unique self-adjoint dual momentum for $H = -iX_1$.

This is not a claim that essential self-adjointness fails for every incomplete vector field in full generality, nor that the phenomenon is new in the abstract: incomplete flows and half-line models are classical [1]. The point is geometric and structural. The same map that is a noninjective Keller seed (constant nonzero Jacobian, restated from Alpöge–Fable) induces dual transport whose incompleteness is explicit and of positive transverse measure in both time directions. The algebraic lift therefore underdetermines quantization for this dual momentum: many self-adjoint extensions exist, and the core polynomial relations do not name one.

What the paper does *not* establish is equally important. We do not claim discovery of F ; unitary gates, channels, or computational advantage;

essential-self-adjointness failure for H_0 or H_2 ; strong CCR after extensions; or a preferred physical boundary condition at the incomplete ends. Lean certification covers the seed identities (Theorem A) only.

Natural next steps are the ESS status of H_0 and H_2 , the residual CCR question after choosing extensions, finer spectral theory of the extension family, and Lean coverage beyond Gate 0.

7 Non-claims

No unitary gate, channel, CP map, or computational advantage. No ESS claim for H_0, H_2 . No strong CCR after extensions. No discovery claim for F . No preferred physical extension. Lean covers Theorem A only.

References

- [1] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II*, Academic Press, 1975.
- [2] L. Alpöge, announcement, July 2026 (credits Fable).
- [3] D. E. Fredriksen, EXOTIC-CCR A001, GitHub Quantyra/jacobian-weyl-quantum-phase-space v0.3.3.
- [4] D. E. Fredriksen, exotic-ccr-lean, GitHub Quantyra/exotic-ccr-lean.