

Completely positive readings of an exotic Weyl endomorphism: algebraic correspondence and standard-package obstructions

Daniel Eric Fredriksen
Quantyra Inc.

<https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>

22 July 2026

Abstract

Let ψ be the polynomial Weyl endomorphism of the Alpöge–Fable seed (restated): $Q = F(q)$, $P = J^{-T}p$. With the physics normalization $[q_i, p_j] = i\delta_{ij}$ and $q_i^* = q_i$, $p_j^* = p_j$, ψ is a unital $*$ -endomorphism (algebraic SOS positivity). We construct its algebraic endomorphism correspondence and obstruct: (i) unital/nondegenerate $*$ -endomorphisms of $C_0(\mathbb{R}^3)$ extending composition with non-proper F ; (ii) Bogoliubov/quasifree endomorphisms of the CCR algebra (nonlinear F). A former “Free-Strict-Regular-v2” nondegeneracy obstruction is **withdrawn** as incorrect. We then **construct** a genuine normal unital $*$ -homomorphism (Koopman pull-back) on the bounded-Borel position sector, valid for any local-diffeomorphism F regardless of properness or surjectivity. For the joint q – p sector we record: no choice-free Joint-Stone-Canonical package at the H_1 leg (A001 deficiency (∞, ∞)); under the G4 companion H_0, H_2 deficiency pairs (not an A001 claim)—no full-triple Joint-Stone package; and a new **obstruction** for multiplicative single-generator Joint-Stone-Hom-1 (Theorem J4: Stone–von Neumann product structure vs. s -dependent preimage count of F). Bare Joint-Form-Core is constructed on C_c^∞ ; Joint-Form with ESS uniqueness fails for H_1 . Residual plain-CP choice-dependent Joint-Stone remains open. No dual F -translations, gates, or advantage. Companion to [1]. Pack: COMPANION-PACK.md.

MSC: 46L07, 81S05, 47L90, 46L08.

Version: C001 companion v0.6 (J4 Hom obstruction; Joint-Form split).

T4: no dual F -translations on $L^2(\mathbb{R}^3)$.

Erratum. Free-Strict-Regular-v2 (projection off $\overline{\text{im } F}$) is **withdrawn**: $q \notin \overline{\text{im } F}$ does not imply $f(F(q)) = 0$, and $M_f \mapsto M_{f \circ F}$ on $C_b(Q)$ is a valid unital $*$ -homomorphism extendable by Arveson. Do not cite v0.2 Regular-v2 as a theorem.

1 Domain

Definition 1. $\mathcal{W}$: polynomial Weyl algebra over $\mathbb{C}$ with $[q_i, p_j] = i\delta_{ij}$, $[q_i, q_j] = [p_i, p_j] = 0$, and involution $q_i^* = q_i$, $p_j^* = p_j$. (Consistent: $[q, p]^* = -[q, p]$ and $(i\delta)^* = -i\delta$.) With real $B = J^{-T}$,

$$\psi(q_i) = F_i(q), \quad \psi(p_j) = \sum_k B_{jk}(q) p_k$$

(coefficients left of momenta) extends to a unital algebra endomorphism.

Remark 1 (Normalization). *Earlier drafts used $[q, p] = \delta$ with self-adjoint generators, which is inconsistent under $*$. This note uses the standard physics normalization $[q, p] = i$.*

Theorem 1 (Algebraic $*$ -SOS). *Assume F, B real and the Piola/dual-field identity $\operatorname{div} B = 0$ (row divergence free; true for the A001 cotangent lift). Then each Schrödinger operator $H_j = -iX_j$ is symmetric on $C_c^\infty(\mathbb{R}^3)$ (not claimed essentially self-adjoint—cf. [1] for H_1). At the algebraic level the formal adjoint correction $-i \sum_k \partial_k B_{jk}$ vanishes, so*

$$P_j^* = P_j \quad \text{in } \mathcal{W}$$

with the physics involution. Hence ψ is a unital $$ -endomorphism and $\operatorname{id}_n \otimes \psi$ preserves hermitian-square cones.*

Definition 2 (Algebraic correspondence). *Let $E = \mathcal{W}$ as a right $\mathcal{W}$ -module by $\xi \cdot b = \xi b$, with left action $a \cdot \xi = \psi(a)\xi$ and $\langle \xi, \eta \rangle_{\mathcal{W}} = \xi^* \eta$.*

2 Geometry

Theorem 2 (Non-proper). *F is not proper along the incomplete X_1 -orbit of [1] (CAS `cas.F_not_proper.C1c.A001`).*

3 Obstructions (retained)

Theorem 3 (C_0 composition endo). *There is no $*$ -homomorphism $\alpha : C_0(\mathbb{R}^3) \rightarrow C_0(\mathbb{R}^3)$ (nondegenerate/approximately unital in the usual C_0 sense) with $\alpha(f) = f \circ F$ for all $f \in C_0(\mathbb{R}^3)$, because F is not proper so $f \circ F \notin C_0$ for some $f \in C_0$.*

Theorem 4 (Bogoliubov / quasifree). *No Bogoliubov (linear quasifree) endomorphism of the CCR C^* -algebra—i.e. a map induced by a linear symplectic contraction on the one-particle space, acting on Weyl unitaries by $W(z) \mapsto W(Tz)$ —can reproduce the Heisenberg picture dynamics of ψ on the Weyl one-parameter groups associated with the field coordinates in a regular representation: those Bogoliubov maps act linearly on one-particle fields, while $q \mapsto F(q)$ is a nonlinear polynomial map (total degree > 1 ; A001 seed has terms up to total degree 7).*

Remark 2 (Not claimed). *Polynomial q, p are not themselves elements of the CCR C^* -algebra; the comparison is through regular representations / affiliated fields / Weyl groups. We do not claim every continuous unital $*$ -endo of the CCR algebra fails—only Bogoliubov/quasifree. We do not claim $f \mapsto M_{f \circ F}$ fails as a map $C_b \rightarrow B(L^2)$. Arveson extensions of the abelian composition map need not implement the momentum part of ψ .*

4 Withdrawn

Free-Strict-Regular-v2 nondegeneracy NO-GO: **withdrawn** (see erratum).

5 Construct: Koopman position sector

Theorem 5 (Koopman pullback). *Let $\mathcal{B}_b(\mathbb{R}^3)$ be the abelian von Neumann algebra of bounded Borel functions of Q . For F a local diffeomorphism ($\det DF \neq 0$ everywhere, in particular the A001 seed), the pullback*

$$\Phi_0(f) := f \circ F, \quad f \in \mathcal{B}_b(\mathbb{R}^3),$$

is a well-defined **unital normal $*$ -homomorphism** $\mathcal{B}_b(\mathbb{R}^3) \rightarrow \mathcal{B}_b(\mathbb{R}^3)$ (hence a normal unital CP map), for any such F —regardless of properness or surjectivity.

Proof. F continuous $\Rightarrow f \circ F$ Borel and bounded whenever f is. Local diffeomorphism $\Rightarrow F$ is Lebesgue-nonsingular (preimages of null sets are null, by the inverse function theorem on a countable cover of charts), so Φ_0 descends to a.e.-equivalence classes. $(f \circ F)^* = \overline{f \circ F}$ and $(fg) \circ F = (f \circ F)(g \circ F)$ pointwise, hence a.e. A unital $*$ -homomorphism of von Neumann algebras is automatically normal and completely positive. $\square$

Remark 3. *This resolves, at the position-sector level, the intuition behind the withdrawn Regular-v2: non-surjectivity of F does not obstruct the bounded-Borel Koopman pullback. The C0 obstruction (Theorem 3) is specific to the vanishing-at-infinity algebra C_0 , not to bounded Borel functions.*

6 Joint q - p extension: five-slice status

Φ_0 only captures ψ on the position generators. The joint sector is no longer a single vague “open”; it splits into five slices as follows (full write-up: `PROGRAM-C-Free-Strict-Abstract-Joint.md`; J4 detail: `PROGRAM-C-residual-J4-Joint-Stone-Hom.md`).

(i) **Position sector — CONSTRUCT.** The Koopman pullback Φ_0 is a normal unital $*$ -homomorphism on bounded Borel functions of Q (previous section).

(ii) **Joint-Stone-Canonical — CONDITIONAL-OBSTRUCTION.** A *Joint-Stone* package asks for self-adjoint extensions $\tilde{H}_j \supset H_j = -iX_j$ together with a unital normal CP map Φ extending Φ_0 and implementing $\Phi(e^{isP_j^{\text{Sch}}}) = e^{is\tilde{H}_j}$. *Joint-Stone-Canonical* further requires the package to be uniquely fixed by the algebraic data (ψ, F, B) alone.

- **Theorem J2** (A001 H_1 only): since $(n_+, n_-) = (\infty, \infty)$ for $H_1 = -iX_1$ [1], von Neumann’s theorem yields a continuum of inequivalent SA extensions; (ψ, F, B) does not select one. Hence no Joint-Stone-Canonical package exists. This does *not* rule out every choice-dependent Joint-Stone package.
- **Theorem J3** (G4 companion, not A001): the companion note `G4-H0-H2-deficiency-bounds.md` records $(n_+, n_-) = (\infty, 0)$ for H_0 and $(0, \infty)$ for H_2 , so those minimal operators admit *no* SA extensions. Under those pairs, no full-triple Joint-Stone package $(\Phi; \tilde{H}_0, \tilde{H}_1, \tilde{H}_2)$ exists at all.

(iii) **Joint-Stone-Hom-1 — OBSTRUCTION (new).** A *Joint-Stone-Hom-1* package asks for a self-adjoint extension $\tilde{H}_1 \supset H_1$ and a unital normal $*$ -homomorphism Φ on the vNa generated by bounded Borel functions of Q and $\{e^{isP_1^{\text{Sch}}}\}$ with $\Phi|_{\mathcal{B}_b(Q)} = \Phi_0$ and $\Phi(e^{isP_1^{\text{Sch}}}) = e^{is\tilde{H}_1}$.

Theorem 6 (J4). *No Joint-Stone-Hom-1 package exists for the A001 seed.*

Proof sketch. Multiplicativity yields dual- F_1 Heisenberg covariance $e^{-is\tilde{H}_1}(f \circ F)e^{is\tilde{H}_1} = f(F_0, F_1 + s, F_2)$. Hence $(F_1, \tilde{H}_1)$ is a regular Schrödinger couple and F_0, F_2 act on the multiplicity space only. The joint spectral measure of (F_0, F_1, F_2) must then have s -independent Lebesgue density. But $|\det DF| \equiv 2$, so that density equals $m/2$ with $m = \#F^{-1}$. For A001, m takes the values 3 and 1 on two positive-measure open slabs near $(a, c) = (0, 2)$. Contradiction. Full write-up: `PROGRAM-C-residual-J4-Joint-Stone-Hom.md`. $\square$

J4 uses multiplicativity in an essential way; it does *not* rule out merely completely positive (non-homomorphic) choice-dependent Joint-Stone.

(iv) **Joint-Form split.**

- **Joint-Form-Core — CONSTRUCT:** sesquilinear forms of the minimal dual momenta $H_j = -iX_j$ on C_c^∞ realize $\psi(p_j)$ together with Φ_0 on positions (no SA/Stone requirement).
- **Joint-Form-ESS-1 — OBSTRUCT:** requiring ESS uniqueness for the $j = 1$ form fails by A001 Theorems E–F.

(v) **Residual route — OPEN.** Choice-dependent single-extension Joint-Stone with a *mere CP* (non-multiplicative) map extending Φ_0 remains **open**. See `PROGRAM-C-C1d-Koopman-position.md` and `PROGRAM-C-Free-Strict-Abstract-Joint.md`.

7 Non-claims

No channel on $B(H)$ implementing the full ψ . No gates/advantage. No dual F -translations. No “all quantization fails.” No claim that every joint CP construction is impossible—only Joint-Stone-Canonical (J2), full-triple Joint-Stone under G4 pairs (J3), multiplicative Joint-Stone-Hom-1 (J4), and Joint-Form-ESS-1. No promotion of G4 H_0/H_2 pairs to A001 theorems.

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts. . . , v0.3.6-submit, <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.