

Completely positive readings of an exotic Weyl endomorphism: algebraic correspondence, and Free-Strict obstructions

Daniel Eric Fredriksen
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

Let ψ be the polynomial Weyl endomorphism of the Alpöge–Fable seed (restated): $Q = F(q)$, $P = J^{-T}p$. We record that ψ is a unital $*$ -endomorphism (algebraic SOS positivity) and construct its algebraic endomorphism correspondence. We obstruct standard packages (C0 $*$ -endo, Bogoliubov, CCR $*$ -hom, C0-compatible CP) and a **Free-Strict-Regular** package: unital CP on $B(L^2)$ with spectral implementation $\Phi(f(Q)) = M_{f \circ F}$ for $f \in C_b$ and nondegeneracy on $C_0(Q)$. A fully abstract C $*$ envelope CP map with no spatial axioms remains open. No dual F -translations, gates, or advantage. Companion to [1].

MSC: 46L07, 81S05, 47L90, 46L08.

Version: C001 Free-Strict v0.2.

T4: no dual F -translations on $L^2(\mathbb{R}^3)$.

1 Domain and algebraic positivity

Definition 1. $\mathcal{W}$: polynomial Weyl algebra, $[q_i, p_j] = \delta_{ij}$, $q_i^* = q_i$, $p_j^* = p_j$. With $B = J^{-T}$, $\psi(q_i) = F_i(q)$, $\psi(p_j) = \sum_k B_{jk}(q)p_k$.

Theorem 1 (Algebraic $*$ -SOS). *Real $F, B \Rightarrow \psi$ unital $*$ -endo $\Rightarrow$ SOS cones preserved at all matrix levels.*

Definition 2 (Algebraic correspondence). $E = \mathcal{W}$ as right $\mathcal{W}$ -module, $a \cdot \xi = \psi(a)\xi$, $\langle \xi, \eta \rangle = \xi^* \eta$.

2 Geometric facts

Theorem 2 (Non-proper / non-surjective). *F is not proper (escape curve of [1]; CAS `cas.F_not_proper_C1c_A001`) and not surjective ($F(\mathbb{R}^3) \neq \mathbb{R}^3$).*

3 Standard package NO-GOs

Theorem 3. *No unital $*$ -endo of $C_0(\mathbb{R}^3)$ extends $f \mapsto f \circ F$.*

Theorem 4. *No Bogoliubov / CCR unital $*$ -endo restricts to ψ (F nonlinear).*

Theorem 5. *No unital CP map that is C0-composition compatible extends ψ .*

4 Free-Strict-Regular NO-GO

Theorem 6 (Free-Strict-Regular-v2). *There is no unital CP map $\Phi : B(L^2(\mathbb{R}^3)) \rightarrow B(L^2(\mathbb{R}^3))$ such that*

1. $\Phi(f(Q)) = M_{f \circ F}$ for all $f \in C_b(\mathbb{R}^3)$;
2. whenever $P \in B(L^2)$ satisfies $\Phi(f(Q))P = 0$ for all $f \in C_0(\mathbb{R}^3)$, then $P = 0$ (nondegeneracy on positions).

Proof. Let $P = M_{1_{\mathbb{R}^3 \setminus F(\mathbb{R}^3)}}$. Non-surjectivity gives $P \neq 0$. For $f \in C_0(\mathbb{R}^3)$, $f \circ F = 0$ on $\text{supp } P$, hence $M_{f \circ F}P = 0$, i.e. $\Phi(f(Q))P = 0$, contradicting nondegeneracy. $\square$

Remark 1. *Stinespring data for this pack are vacuous (empty domain).*

5 Free-Strict-Abstract (residual OPEN)

Unital CP on some abstract C^* envelope of $\mathcal{W}$ with **no** spectral-position, C0-composition, Bogoliubov, or CCR-*-hom axiom: not constructed beyond the algebraic correspondence and finite-level algebraic Stinespring; not ruled out.

6 Non-claims

No channel on $B(H)$ constructed. No gates/advantage. No dual F -translations. No claim every abstract CP extension fails.

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts. . . , v0.3.6-submit, <https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>.