

Dual-lift realizability for Keller-type maps: structural incompleteness, schema, and examples

Daniel Eric Fredriksen
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

We isolate a structural lemma for dual vector fields of local diffeomorphisms $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$: if global dual fields X_j with $DF \cdot X_j = e_j$ are all complete, then F is surjective. Hence any non-surjective dual system has at least one incomplete dual field. We propose a classification schema (B-incomplete / B-ESS / B-Def / B-CCR / B-CP) and populate it on atlas rows A000–A008 (pass and fail baselines, Keller seed A001, family pilot A002, and a genuinely 2D pair A007/A008 illustrating that the forced incomplete set can be as thin as measure zero, contrasting the open-set failures of A001/A006). No gates, channels, or computational advantage are claimed. Companion pack: <https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space/blob/main/docs/notes/COMPANION-PACK.md>.

MSC: 37C10, 47B25, 53D17, 14R15.

Version: B001 draft v0.3.

Software: atlas program.b; CAS cas_atlas_A005_A006_B001.json, cas_atlas_A007_A008_B001.json.

1 Introduction

Cotangent lifts and dual fields are standard when $\det DF \neq 0$. After the A001 deficiency analysis [1], we classify which maps force incomplete dual fields and how that interacts with ESS / CCR / CP packages [2]. A001 is awaiting math-ph endorsement (code VIPN6B); no arXiv id is assigned yet, but the concept DOI already cites A001.

2 Structural lemma

Lemma 1 (Complete dual fields imply surjectivity). *Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be C^∞ and a local diffeomorphism with dual fields X_j , $DF \cdot X_j = e_j$. If every X_j is complete, then $F(\mathbb{R}^n) = \mathbb{R}^n$.*

Proof. $F(\phi_t^{(j)}(q)) = F(q) + te_j$; iterate; image open and translation-invariant. □

Corollary 1. *Non-surjective dual systems have some incomplete X_j .*

3 Schema

B-incomplete / B-ESS / B-Def / B-CCR / B-CP with values open / fail / pass.

4 Atlas scoreboard (v0.3)

ID	Map (sketch)	B-inc.	Role
A000	id	pass	baseline
A004	shear $(x, y + x, z)$	pass	linear auto
A005	diag $(2, \frac{1}{2}, 1)$	pass	linear stretch (CAS)
A008	$(x, y + \sin x, z)$	pass	2D nonlinear global diffeo (CAS)
A003	(x, e^y) pad	fail	smooth B1 example
A006	(e^x, e^y, z)	fail	two incomplete fields, open set (CAS)
A007	$z \mapsto e^z$ pad	fail	incomplete on <i>null set only</i> (CAS)
A001	Keller seed	fail	$X_1 + H_1$ analytic [1]
A002	family d4 pilot	open	geometry pending

5 Examples (new in v0.2)

Example 1 (A005 — linear stretch, pass-like). $F = \text{diag}(2, \frac{1}{2}, 1)$, $\det DF = 1$. Dual fields $X_0 = \frac{1}{2}\partial_x$, $X_1 = 2\partial_y$, $X_2 = \partial_z$ are complete. CAS: `cas_atlas_A005_A006_B001.json`.

Example 2 (A006 — exp quadrant, fail-like). $F(x, y, z) = (e^x, e^y, z)$, $\det DF = e^{x+y} > 0$, image $(0, \infty)^2 \times \mathbb{R}$. Dual fields $X_0 = e^{-x}\partial_x$, $X_1 = e^{-y}\partial_y$ incomplete backward ($e^x = e^{x_0} + t$). Matches Lemma 1. CAS same file.

6 Examples (new in v0.3): a genuine dichotomy

Example 3 (A007 — complex exponential, thin fail-like). $F(x, y, z) = (e^x \cos y, e^x \sin y, z)$, i.e. $z \mapsto e^z$ padded. Conformal, $\det DF = e^{2x} > 0$; image misses the origin (non-surjective). Writing $w = e^{x+iy}$, the dual field X_0 solves $\dot{w} = 1$, so $w(t) = w(0) + t$. If $\text{Im } w(0) \neq 0$ (full measure), the orbit never reaches $w = 0$: **complete**. If $\text{Im } w(0) = 0$ (a measure-zero pair of rays), $w(t)$ hits 0 at finite t and $x(t) \rightarrow -\infty$: **incomplete**. Thus Lemma 1's corollary is satisfied by an incomplete set of measure zero, sharply contrasting A006/A001 where incompleteness occupies an open set. CAS: `cas_atlas_A007_A008_B001.json`.

Example 4 (A008 — shear + sine, 2D global diffeomorphism). $F(x, y, z) = (x, y + \sin x, z)$, $\det DF = 1$. Explicit global inverse $(x, y) \mapsto (x, y - \sin x)$ (verified by CAS roundtrip). Dual fields $X_0 = \partial_x - \cos x \partial_y$, $X_1 = \partial_y$; the X_0 flow $y(t) = y_0 - \sin(x_0 + t) + \sin(x_0)$ is bounded for all t : **complete**. A genuinely nonlinear, non-triangular-affine 2D coupling (unlike A004/A005) that nonetheless has no incompleteness at all.

7 Toward a dichotomy (open)

A007 shows the incomplete locus forced by Lemma 1 can be as thin as measure zero for general smooth local diffeomorphisms. A007 is *not* polynomial, so it does not settle whether **polynomial** Keller maps specifically always force an *open* incomplete set (as in A001/A006) or could also exhibit thin failure. We record this as an open classification question (see PROGRAM-B-B4-2D-dichotomy.md) rather than claim a theorem.

8 Prior examples (v0.1–v0.2)

A000 identity; A003 half-plane; A001 Keller seed with X_1 incomplete and $H = -iX_1$ deficiency (∞, ∞) in [1] (other X_j incompleteness is recorded in companion validation dossiers, not re-proved here); A002 open; A004 shear pass.

9 Non-claims

No gates, channels, advantage. No dual-flow repair. A003/A005/A006/A007/A008 are not Jacobian counterexamples (A007/A008 are not even polynomial). T4: no dual F -translation package claimed here. No theorem is claimed for the polynomial-only dichotomy question (§7, open). C Free-Strict-Abstract-Joint remains open [2] (position sector now constructed via Koopman pull-back).

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., v0.3.6-submit / concept DOI 10.5281/zenodo.21474351, <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.
- [2] D. E. Fredriksen, CP readings of an exotic Weyl endomorphism..., C001 companion v0.4 (Koopman position construct), <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.