

Poisson and Weyl lifts of the Alpöge–Fable Keller
map
and nonunique self-adjoint realizations of a dual
transport operator

Daniel Eric Fredriksen

Quantyra Inc.

<https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

We study the three-dimensional Keller map F announced by Alpöge (with Fable credited): $\det DF \equiv -2$ and a three-point collision. Theorem A restates those finite identities; novelty is Theorems B–F. The cotangent lift $\Phi(q, p) = (F(q), J^{-T}p)$ is a polynomial Poisson map and non-injective. Dual fields $X_j = \sum_k B_{jk} \partial_{q_k}$ commute and are divergence-free (Piola), so $H_j = -iX_j$ are symmetric on $C_c^\infty(\mathbb{R}^3)$. The field X_1 is incomplete. In flow-box coordinates $(a, s, c) = F(q)$ one has $X_1 = \partial_s$ and $dq = \frac{1}{2} da ds dc$. Open forward and backward escape walls yield whole-orbit deficiency vectors in $\text{Dom}(H^*)$ (transverse cutoffs only; no interior s -indicators), hence

$$(n_+, n_-) = (\infty, \infty)$$

for $H = -iX_1$. Infinitely many self-adjoint extensions exist; algebraic CCR/Poisson data select none. No gates, channels, computational advantage, or seed discovery claim.

MSC 2020: 47B25, 81S05, 14R15, 37C10, 53D17.

Software: v0.3.2; concept DOI 10.5281/zenodo.21474351 (version DOI on Zenodo ingest).

Errata. v0.2.2 pair $(0, \infty)$ withdrawn. v0.3.0 interior s -indicator deficiency proof withdrawn ($\notin \text{Dom}(H^*)$).

1 Introduction

On 19–20 July 2026, Levent Alpöge announced $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ with $\det DF \equiv -2$ and a three-point collision (credits Fable) [2]. **We do not claim discovery of F .** We ask whether algebraic Poisson/CCR lifts select a unique self-adjoint dual transport generator on $C_c^\infty(\mathbb{R}^3)$. Answer: no for $H = -iX_1$; deficiency indices (∞, ∞) .

2 Setup

$$\begin{aligned} F_0 &= (1 + xy)^3 z + y^2(1 + xy)(4 + 3xy), \\ F_1 &= y + 3x(1 + xy)^2 z + 3xy^2(4 + 3xy), \\ F_2 &= 2x - 3x^2 y - x^3 z. \end{aligned}$$

$$B = J^{-T}, \quad X_j = \sum_k B_{jk} \partial_{q_k}, \quad H := -iX_1.$$

3 Theorems A–D (summary)

Theorem 1 (A). $\det DF \equiv -2$; *three-point collision*; F *noninjective*. (*Restatement; Lean+CAS.*)

Theorem 2 (B–C). Φ *is polynomial Poisson and noninjective*. $X_j(F_i) = \delta_{ij}$, $[X_i, X_j] = 0$, $\operatorname{div} X_j = 0$ (*Piola*), $H_j = -iX_j$ *symmetric*, $[F_i, H_j] = i\delta_{ij}$ *on C_c^∞* .

Theorem 3 (D). X_1 *incomplete*: *explicit γ through $(1, 0, 0)$ with $F(\gamma(t)) = (0, t, 2)$ escapes as $t \rightarrow \frac{1}{2}^-$.*

4 Flow-box coordinates

On local inverse sheets, $(a, s, c) = F(q)$ gives

$$dq = \frac{1}{2} da \, ds \, dc, \quad X_1 = \partial_s, \quad H = -i\partial_s. \quad (7.1)$$

Elimination yields $A(s; a, c)q_0^3 + B(s; c)q_0 + C(c) = 0$ with $A = -cs^3 + s^2 + 18acs - 27a^2c^2 - 16a$, $B = 3cs - 4$, $C = 2c$.

Reconstruction. If $x = q_0$ solves the cubic and denominators are nonzero,

$$q_1 = \frac{9acx^2 - 3csx - 3c - sx^2 + 6x}{x((3cs - 4)x + 3c)}, \quad q_2 = \frac{2x - 3x^2q_1 - c}{x^3},$$

then $F(x, q_1, q_2) = (a, s, c)$.

5 Theorems E–F

Dictionary. On (ℓ, r) : finite upper end $\Rightarrow n_- = 1$ (model e^{s-r}); finite lower end $\Rightarrow n_+ = 1$ (model $e^{-(s-\ell)}$).

Proposition 1 (Forward wall). At $(0, \frac{1}{2}, 2)$, $A = 0$, $\partial_s A = -\frac{1}{2}$, $B = -1$. IFT gives wall β . Set $\tau = \sqrt{\beta - s}$, $q_0 = r/\tau$, and

$$G_+ = \frac{A(\beta - \tau^2)}{\tau^2} r^3 + B(\beta - \tau^2)r + C\tau,$$

extended at $\tau = 0$ by $(-A_s)r^3 + Br$. Simple roots $r = \pm\sqrt{2}$ ($\partial_r G_+ = -2B \neq 0$) continue by IFT; reconstruct by the formulas above.

Proposition 2 (Backward wall). At $a = \frac{1}{54}$, $c = 2$, wall α with $A_s = \frac{1}{6}$, $B = -1$. Set $\tau = \sqrt{s - \alpha}$ and analogous G_- extended by $A_s r^3 + Br$. Simple roots $r = \pm\sqrt{6}$ continue by IFT on an open transverse set.

Lemma 1 (Saturated maximal sheets). Cross-section the selected branch below/above the wall; saturate under the flow of X_1 . Since $X_1 F_1 = 1$, s is time. ODE theory yields a C^∞ diffeomorphism Ψ_+ from

$$D_+ = \{(a, c, s) : (a, c) \in W_+ \subseteq U_+, \ell_+(a, c) < s < \beta(a, c)\}$$

onto an open invariant set $\Omega_+ \subset \mathbb{R}^3$, with $F \circ \Psi_+ = (a, s, c)$. The end $s \rightarrow \beta^-$ is at spatial infinity. If ℓ_+ is finite, maximality forces escape as $s \downarrow \ell_+$ as well. Analogous Ψ_-, Ω_- for the lower wall.

Lemma 2 (Deficiency vectors in $\text{Dom}(H^*)$). For $\chi \in C_c^\infty(W_+)$ define on the entire maximal interval

$$u_-(\Psi_+(a, c, s)) = \chi(a, c) e^{s-\beta(a, c)},$$

and $u_- = 0$ off Ω_+ . No interior s -cutoff. Then

$$\|u_-\|_2^2 = \frac{1}{2} \int_{W_+} |\chi|^2 \int_{\ell_+}^{\beta} e^{2(s-\beta)} ds da dc \leq \frac{1}{4} \|\chi\|_{L^2}^2.$$

For $\varphi \in C_c^\infty(\mathbb{R}^3)$, $\text{supp } \varphi$ meets orbits in compact s -segments strictly inside (ℓ_+, β) . Integration by parts with $\text{div } X_1 = 0$ gives no jump and no endpoint terms, so $u_- \in \text{Dom}(H^*)$ and $(H^* + i)u_- = 0$. Thus $n_- = \infty$. Likewise

$$u_+(\Psi_-(a, c, s)) = \chi(a, c) e^{-(s - \alpha(a, c))}$$

on (α, r_-) yields $n_+ = \infty$.

Theorem 4 (F). $(n_+, n_-) = (\infty, \infty)$.

Proof. The two walls, saturation, and deficiency lemmas; separability. $\square$

Theorem 5 (E). H is not essentially self-adjoint on $C_c^\infty(\mathbb{R}^3)$.

Proof. $n_\pm \geq 1$. $\square$

Corollary 1. Infinitely many self-adjoint extensions exist. Core algebraic Poisson/CCR relations do not distinguish among them.

6 Non-claims

No unitary gate, channel, CP map, or computational advantage. No ESS claim for H_0, H_2 . No strong CCR after extensions. No discovery claim for F . No preferred physical extension. Lean covers Theorem A only.

References

- [1] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II*, Academic Press, 1975.
- [2] L. Alpöge, announcement, July 2026 (credits Fable).
- [3] D. E. Fredriksen, EXOTIC-CCR A001, GitHub Quantyra/jacobian-weyl-quantum-phase-space v0.3.2.
- [4] D. E. Fredriksen, exotic-ccr-lean, GitHub Quantyra/exotic-ccr-lean.