

Poisson and Weyl lifts of the Alpöge–Fable Keller  
map  
and nonunique self-adjoint realizations of a dual  
transport operator

Daniel Eric Fredriksen

Quantyra Inc.

<https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>

21 July 2026

**Abstract**

We study the three-dimensional Keller map  $F$  announced by Alpöge (with Fable credited):  $\det DF \equiv -2$  and a three-point collision. Theorem A restates those finite identities (independently dual-CAS and Lean checked); novelty is Theorems B–F. We construct the cotangent lift  $\Phi(q, p) = (F(q), J^{-T}p)$  and show it is a polynomial Poisson map on generators and non-injective. The matrix  $B = J^{-T}$  defines dual fields  $X_j = \sum_k B_{jk} \partial_{q_k}$ ; by the Piola identity and constant  $\det DF$ ,  $\operatorname{div} B = 0$ , and the fields commute because they are dual to  $dF_i$ . The operators  $H_j = -iX_j$  are symmetric on  $C_c^\infty(\mathbb{R}^3)$ .

We prove  $X_1$  is incomplete by an explicit integral curve escaping as  $t \rightarrow \frac{1}{2}^-$ . In flow-box coordinates  $(a, s, c) = F(q)$  one has  $X_1 = \partial_s$  and  $dq = \frac{1}{2} da ds dc$ . Open transverse families with finite upper and lower  $F_1$ -ends yield square-integrable deficiency functions, so

$$(n_+, n_-) = (\infty, \infty)$$

for  $H = -iX_1$ . Hence  $H$  is not essentially self-adjoint, yet von Neumann’s theorem supplies infinitely many self-adjoint extensions; the algebraic CCR data select none of them.

We claim no unitary gates, channels, computational advantage, or preferred physical extension.

**MSC 2020:** 47B25, 81S05, 14R15, 37C10, 53D17.

**Keywords:** Jacobian conjecture, Keller map, cotangent lift, Weyl algebra,

essential self-adjointness, deficiency indices.

**Software:** concept DOI 10.5281/zenodo.21474351; v0.3.0 DOI 10.5281/zenodo.21477350.

**Erratum.** Release v0.2.2 claimed  $(n_+, n_-) = (0, \infty)$ . An open backward-incomplete family forces  $n_+ = \infty$  as well; the corrected pair is  $(\infty, \infty)$ .

## 1 Introduction

### 1.1 Motivation and attribution

A *Keller map* is a polynomial endomorphism  $F$  of affine space with  $\det DF$  a nonzero constant. The Jacobian conjecture predicts every Keller map is a polynomial automorphism. On 19–20 July 2026, Levent Alpöge announced an explicit map  $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$  with  $\det DF \equiv -2$  and a three-point collision, crediting the AI system Fable [7]; independent checks and Lean formalizations followed promptly (e.g. Cureton [8]; Speyer [9]). **This paper does not claim discovery of  $F$ .** Repository atlas label: A001.

The cotangent (Piola) lift  $\Phi(q, p) = (F(q), J^{-T}p)$  and dual transport fields  $X_j$  are standard when  $\det J$  is a nonzero constant. We ask whether algebraic preservation of Poisson/CCR data selects a unique self-adjoint  $L^2$  realization of a dual transport generator on  $C_c^\infty(\mathbb{R}^3)$ .

### 1.2 Main results

- (A) Restatement:  $\det DF \equiv -2$  and the three-point collision hold.
- (B)  $\Phi$  is a polynomial Poisson map on generators and is non-injective.
- (C) Dual fields commute and are divergence-free;  $H_j = -iX_j$  is symmetric on  $C_c^\infty$ .
- (D)  $X_1$  is incomplete via an explicit escaping integral curve.
- (E)  $H = -iX_1$  is not essentially self-adjoint.
- (F)  $(n_+, n_-) = (\infty, \infty)$ ; infinitely many self-adjoint extensions exist, none selected by the algebraic data.

### 1.3 Related work

Jacobian conjecture background: Keller [4], Bass–Connell–Wright [3]. Deficiency indices: Reed–Simon [2]. Chernoff [1] concerns *sufficiency* criteria for ESS and is not used as a necessity theorem here.

## 2 Setup

$$\begin{aligned} F_0 &= (1 + xy)^3 z + y^2(1 + xy)(4 + 3xy), \\ F_1 &= y + 3x(1 + xy)^2 z + 3xy^2(4 + 3xy), \\ F_2 &= 2x - 3x^2 y - x^3 z. \end{aligned} \tag{2.1}$$

Write  $J = DF$ ,  $B := J^{-T}$ , and

$$X_j := \sum_{k=0}^2 B_{jk}(q) \partial_{q_k}. \tag{2.2}$$

Collision data:

$$F(0, 0, -\tfrac{1}{4}) = F(1, -\tfrac{3}{2}, \tfrac{13}{2}) = F(-1, \tfrac{3}{2}, \tfrac{13}{2}) = (-\tfrac{1}{4}, 0, 0). \tag{2.4}$$

## 3 Theorem A — Seed identities

**Theorem 1** (A). (1)  $\det DF \equiv -2$ . (2) The identities (4) hold in characteristic not 2. (3) Hence  $F$  is not injective.

$$F(0, 0, -\tfrac{1}{4}) = F(1, -\tfrac{3}{2}, \tfrac{13}{2}) = F(-1, \tfrac{3}{2}, \tfrac{13}{2}) = (-\tfrac{1}{4}, 0, 0). \tag{4}$$

*Proof sketch.* Expand  $\det J$ ; evaluate at the three rational points. Dual CAS and Lean formalization [6].

## 4 Theorem B — Poisson lift

**Theorem 2** (B). Let  $\Phi(q, p) = (F(q), B(q)p)$ . Then  $B$  is polynomial,  $JB^T = I$ , the generator Poisson brackets  $\{Q_i, Q_j\} = \{P_i, P_j\} = 0$ ,  $\{Q_i, P_j\} = \delta_{ij}$  hold, and  $\Phi$  is non-injective via (4).

*Proof sketch.* Standard cotangent lift for constant  $\det J$ ; non-injectivity lifts from collisions using  $BJ^T = I$ . CAS regression tests in the software artifact [5].

## 5 Theorem C — Dual fields and Piola

**Theorem 3 (C).** *Let  $X_j = \sum_k B_{jk} \partial_{q_k}$ . Then  $X_j(F_i) = \delta_{ij}$ ,  $[X_i, X_j] = 0$ ,  $\operatorname{div} X_j = 0$ , and  $H_j := -iX_j$  is symmetric on  $C_c^\infty(\mathbb{R}^3)$  with  $[F_i, H_j] = i\delta_{ij}$ .*

*Proof.*  $X_j(F_i) = (JB^T)_{ij} = \delta_{ij}$ . Then  $[X_i, X_j](F_k) = 0$  for all  $k$ ; since  $dF$  is a coframe,  $[X_i, X_j] = 0$ .  $\operatorname{Cof}(DF) = (\det DF)B$ . Piola gives  $\operatorname{Div} \operatorname{Cof}(DF) = 0$ ; constant  $\det DF = -2$  implies  $\operatorname{Div} B = 0$ . Symmetry of  $-iX$  when  $\operatorname{div} X = 0$  is standard.  $\square$

## 6 Theorem D — Incompleteness of $X_1$

**Theorem 4 (D).** *The dual field  $X_1$  is incomplete on  $\mathbb{R}^3$ .*

*Proof.* For  $t \in [0, \frac{1}{2})$ , with continuous extension  $\gamma(0) = (1, 0, 0)$ ,

$$\begin{aligned} q_0(t) &= \frac{-2t - \sqrt{1 - 2t} + 1}{t(2t - 1)}, \\ q_1(t) &= t, \\ q_2(t) &= t^2(2t - 3\sqrt{1 - 2t} - 1). \end{aligned} \tag{6.1}$$

On  $(0, \frac{1}{2})$ ,  $F(\gamma(t)) = (0, t, 2)$  and  $\gamma' = X_1 \circ \gamma$  (CAS-verified identities). As  $t = \frac{1}{2} - \varepsilon$ ,  $q_0 = \sqrt{2/\varepsilon} + O(1) \rightarrow +\infty$ . Thus the forward maximal time from  $(1, 0, 0)$  is at most  $\frac{1}{2} < \infty$ .  $\square$

## 7 Flow-box coordinates

Write  $H := -iX_1$  and  $n_\pm := \dim \ker(H^* \mp i)$  in the usual Reed–Simon convention  $n_+ = \dim \ker(H^* - i)$ . On local inverse sheets of  $F$ , with  $(a, s, c) = F(q)$ ,

$$dq = \frac{1}{2} da \, ds \, dc, \quad X_1 = \partial_s, \quad H = -i\partial_s. \tag{7.1}$$

Eliminating variables yields

$$A(s; a, c) q_0^3 + B(s; c) q_0 + C(c) = 0, \tag{7.2}$$

with  $A = -cs^3 + s^2 + 18acs - 27a^2c^2 - 16a$ ,  $B = 3cs - 4$ ,  $C = 2c$ . If  $A \rightarrow 0$  and  $B \neq 0$ , then some  $|q_0| \rightarrow \infty$ .

## 8 Theorems E–F — Deficiency indices

**Proposition 1** (Forward wall). *At  $(0, \frac{1}{2}, 2)$ ,  $A = 0$  and  $\partial_s A = -\frac{1}{2} \neq 0$ . IFT yields open  $U_+ \ni (0, 2)$  and a finite upper wall  $\beta : U_+ \rightarrow \mathbb{R}$ .*

**Proposition 2** (Backward wall). *At  $a = \frac{1}{54}$ ,  $c = 2$ ,  $A = -\frac{(2s-1)(3s^2-1)}{3}$ , so  $A(\frac{1}{2}) = 0$ ,  $\partial_s A(\frac{1}{2}) = \frac{1}{6} > 0$ ,  $B(\frac{1}{2}) = -1$ . For  $s = \frac{1}{2} + h$ ,  $h \downarrow 0^+$ , large roots  $q_0 = \pm\sqrt{6/h} + O(1)$  give  $\|q\| \rightarrow \infty$ . IFT yields open  $U_-$  of finite lower walls  $\alpha$ .*

**Lemma 1** (Positive measure).  $\text{Leb}_3(\{T_+ < \infty\}) > 0$  and  $\text{Leb}_3(\{T_- > -\infty\}) > 0$ .

*Proof.* Open flow-box slabs below  $\beta$  (resp. above  $\alpha$ ), pushed by local inverses of  $F$ , have positive Lebesgue measure in  $q$ -space.  $\square$

**Theorem 5** (F).  $(n_+, n_-) = (\infty, \infty)$ .

*Proof.* Saturate a transverse cross-section under the  $X_1$ -flow to obtain full maximal orbit intervals  $(\ell_+, \beta)$  and  $(\alpha, r_-)$ . For  $\chi \in C_c^\infty(W_\pm)$  define *whole-orbit* deficiency vectors (no interior  $s$ -cutoff)

$$u_- = \chi(a, c) e^{s-\beta(a,c)} \quad \text{on } (\ell_+, \beta), \quad u_+ = \chi(a, c) e^{-(s-\alpha(a,c))} \quad \text{on } (\alpha, r_-).$$

Then  $u_\pm \in L^2$ ,  $(H^* + i)u_- = 0$ , and  $(H^* - i)u_+ = 0$  (escape ends at spatial infinity; only transverse cutoffs). Interior indicators  $\mathbf{1}_{(\beta-\delta, \beta)}$  are invalid. Separability yields  $(\infty, \infty)$  [2].  $\square$

**Theorem 6** (E).  $H$  is not essentially self-adjoint on  $C_c^\infty(\mathbb{R}^3)$ .

*Proof.* Immediate from  $n_\pm \geq 1$  in Theorem F.  $\square$

**Corollary 1** (Extensions). *Since  $n_+ = n_- = \infty$ , infinitely many self-adjoint extensions exist. Algebraic CCR/Poisson data do not select a preferred extension.*

**Errata.** v0.2.2 claim  $(0, \infty)$  withdrawn. v0.3.0  $s$ -indicator deficiency proof withdrawn (not in  $\text{Dom}(H^*)$ ).

## 9 Non-claims

No unitary gate, channel, CP map, or computational advantage. No ESS claim for  $H_0, H_2$ . No strong CCR after extensions. No discovery claim for  $F$ . No preferred physical extension.

## 10 Conclusion

Starting from the Alpöge–Fable Keller map, Poisson/Weyl lifts exist,  $X_1$  is incomplete, and  $H = -iX_1$  has deficiency indices  $(\infty, \infty)$ : not ESS, infinitely many self-adjoint extensions, none preferred by algebraic data.

## References

- [1] P. R. Chernoff, Essential self-adjointness of powers of generators of hyperbolic equations, *J. Funct. Anal.* **12** (1973), 401–414.
- [2] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II*, Academic Press, 1975.
- [3] H. Bass, E. H. Connell, and D. Wright, The Jacobian conjecture: reduction of degree and formal expansion of the inverse, *Bull. Amer. Math. Soc. (N.S.)* **7** (1982), 287–330.
- [4] O.-H. Keller, Ganze Cremona-Transformationen, *Monatsh. Math. Phys.* **47** (1939), 299–306.
- [5] D. E. Fredriksen, EXOTIC-CCR A001 software artifact, Zenodo (2026), DOI 10.5281/zenodo.21477350.
- [6] D. E. Fredriksen, exotic-ccr-lean,   
texttt<https://github.com/Quantyra/exotic-ccr-lean>.
- [7] L. Alpöge, announcement that the Jacobian conjecture is false in dimension 3, public post, 19–20 July 2026 (credits Fable); see also software provenance in [5].
- [8] D. Cureton, independent Lean 4 formalization,   
texttt<https://github.com/deancureton/jacobian>.
- [9] D. Speyer, The new counterexample to the Jacobian conjecture, Secret Blogging Seminar, 20 July 2026.