

Poisson and Weyl lifts of the Alpöge–Fable Keller map and nonunique self-adjoint realizations of a dual transport operator

Daniel Eric Fredriksen
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

We study the three-dimensional Keller map F announced by Alpöge (with Fable credited): $\det DF \equiv -2$ and a three-point collision. Theorem A restates those finite identities. Theorems B–C record the standard cotangent/Piola/Weyl package for this seed; the novel analytic contribution is Theorems D–F. The cotangent lift $\Phi(q, p) = (F(q), J^{-T}p)$ is a polynomial Poisson map and non-injective. Dual fields $X_j = \sum_k B_{jk} \partial_{q_k}$ commute and are divergence-free (Piola), so $H_j = -iX_j$ are symmetric on $C_c^\infty(\mathbb{R}^3)$. We define the induced polynomial Weyl endomorphism ψ . The field X_1 is incomplete via an explicit integral curve. In flow-box coordinates $(a, s, c) = F(q)$ one has $X_1 = \partial_s$ and $dq = \frac{1}{2} da ds dc$. Open forward and backward escape walls yield whole-orbit deficiency vectors in $\text{Dom}(H^*)$ (transverse cutoffs only; no interior s -indicators), hence

$$(n_+, n_-) = (\infty, \infty)$$

for $H = -iX_1$. Infinitely many self-adjoint extensions exist; the core algebraic CCR/Poisson data alone do not distinguish among them. No gates, channels, computational advantage, or seed discovery claim. Companion code release **v0.3.5-submit**.

MSC 2020: 47B25, 81S05, 14R15, 37C10, 53D17.

Software: v0.3.5-submit; concept DOI 10.5281/zenodo.21474351.

Categories (suggested): math.FA (primary); math-ph, math.SP (secondary).

Errata (prior releases). v0.2.2 pair $(0, \infty)$ for H withdrawn. v0.3.0 interior s -indicator deficiency proof withdrawn ($\notin \text{Dom}(H^*)$). This submission is $H = -iX_1$ **only**: we do *not* claim exact deficiency pairs for H_0, H_2 , nor a theorem-level dual-flow strong CCR no-go (companion notes only).

1 Introduction

On 19–20 July 2026, Levent Alpöge announced an explicit polynomial map $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ with $\det DF \equiv -2$ and a three-point collision, crediting the system Fable [5, 6, 7]. **We do not claim discovery of F .** We re-verified the seed identities by CAS and recorded a Lean 4 formalization of those finite algebraic checks [9].

We ask whether the standard Poisson/CCR lifts of F select a unique self-adjoint dual transport generator on $C_c^\infty(\mathbb{R}^3)$. For $H = -iX_1$ the answer is negative: deficiency indices (∞, ∞) .

Scope. Theorems A–F below concern the algebra of the lifts and the analytic deficiency theory of $H = -iX_1$. Remarks on other dual fields or alternative realizations are deferred to companion validation notes and are not claimed as theorems here.

2 Setup

Let $q = (q_0, q_1, q_2) = (x, y, z)$ and

$$\begin{aligned} F_0 &= (1 + xy)^3 z + y^2(1 + xy)(4 + 3xy), \\ F_1 &= y + 3x(1 + xy)^2 z + 3xy^2(4 + 3xy), \\ F_2 &= 2x - 3x^2 y - x^3 z. \end{aligned}$$

Write $J = DF$ and $B := J^{-T}$. Dual fields:

$$X_j := \sum_{k=0}^2 B_{jk}(q) \partial_{q_k}, \quad j = 0, 1, 2.$$

Normalizations. We use two compatible packages. (i) *Abstract Weyl algebra* $\mathcal{W}$: generators q_k, p_ℓ with $[q_i, p_j] = \delta_{ij}$ (Dixmier convention). (ii) *Schrödinger representation* on $L^2(\mathbb{R}^3)$: multiplication by q_k and $p_k^{\text{Sch}} = -i\partial_{q_k}$, so $[q_i, p_j^{\text{Sch}}] =$

$i\delta_{ij}$. The operators $H_j := -iX_j$ live in package (ii). When $\operatorname{div} X_j = 0$, each H_j is symmetric on $C_c^\infty(\mathbb{R}^3)$. Collision data ($\operatorname{char} \neq 2$):

$$F(0, 0, -\tfrac{1}{4}) = F(1, -\tfrac{3}{2}, \tfrac{13}{2}) = F(-1, \tfrac{3}{2}, \tfrac{13}{2}) = (-\tfrac{1}{4}, 0, 0). \quad (1)$$

3 Theorem A — seed identities

Theorem 1 (A). (1) $\det DF \equiv -2$. (2) The collision identities (1) hold over any field of characteristic not 2. (3) Hence F is not injective.

Proof. Expand $\det J$ as a polynomial identity, or evaluate the formal determinant in a computer algebra system; formalized in Lean 4 in the companion repository (`jacobianDet.F`, `evalMap.F_p0/p1/p2`) [9]. Non-injectivity is immediate from (1). $\square$

Remark 1 (Complex vs real). Over $\mathbb{C}$, constant nonzero Jacobian and non-injectivity would be extraordinary for the Jacobian conjecture [4]. We treat F as an externally announced seed [5, 6, 7]; Theorem A is a restatement with CAS/Lean re-verification, not a priority claim. Our analytic results use the real $L^2(\mathbb{R}^3)$ dual-field package.

4 Theorem B — Poisson cotangent lift

Theorem 2 (B). Let $\Phi(q, p) = (F(q), B(q)p)$ with $B = J^{-T}$. Then:

1. B is polynomial and $JB^T = I$;
2. writing $Q_i = F_i(q)$ and $P_j = \sum_k B_{jk}(q)p_k$,

$$\{Q_i, Q_j\} = 0, \quad \{P_i, P_j\} = 0, \quad \{Q_i, P_j\} = \delta_{ij}$$

for the canonical Poisson structure on $T^*\mathbb{R}^3$;

3. Φ is not injective (lift the three collision sources with matching p).

Proof. (1) follows from $B = J^{-T}$ and $\det J \equiv -2 \neq 0$; polynomiality of B is the classical adjugate formula $B = -(\operatorname{adj} J)^T/2$. (2) The identities $\{Q_i, P_j\} = (JB^T)_{ij} = \delta_{ij}$ are immediate; vanishing of $\{Q_i, Q_j\}$ is clear; vanishing of $\{P_i, P_j\}$ is the standard cotangent-lift calculation for $B = J^{-T}$ (equivalently dual-CAS generator checks in the companion release [8]). (3) If $F(q^{(a)}) = Q_\star$ and $p^{(a)} = J(q^{(a)})^T P_\star$, then $\Phi(q^{(a)}, p^{(a)}) = (Q_\star, P_\star)$ for each collision source in (1). $\square$

5 Theorem C — dual fields, Piola, Weyl endomorphism

Theorem 3 (C). 1. $X_j(F_i) = \delta_{ij}$, $[X_i, X_j] = 0$, and $\operatorname{div} X_j = 0$.

2. Each $H_j = -iX_j$ is symmetric on $C_c^\infty(\mathbb{R}^3)$, and $[M_{F_i}, H_j] = i\delta_{ij}$ on that domain.

3. Let $\mathcal{W}$ be the polynomial Weyl algebra over $\mathbb{C}$ generated by q_k, p_ℓ with $[q_i, p_j] = \delta_{ij}$ (abstract package) and all other generators commuting. The assignment

$$\psi(q_i) = F_i(q), \quad \psi(p_j) = \sum_k B_{jk}(q) p_k$$

extends uniquely to a unital algebra endomorphism $\psi : \mathcal{W} \rightarrow \mathcal{W}$ preserving the Weyl relations on generators (ordering: coefficients $B_{jk}(q)$ to the left of p_k).

Proof. (1) $X_j(F_i) = (JB^T)_{ij} = \delta_{ij}$. Then $[X_i, X_j](F_k) = 0$ for all k ; since dF is a coframe, $[X_i, X_j] = 0$. Piola: $\operatorname{Div} \operatorname{Cof}(DF) = 0$; with $\det DF \equiv -2$, $\operatorname{Div} B = 0$, i.e. $\operatorname{div} X_j = 0$. (2) Divergence-free smooth vector fields give symmetric $-iX$ on $C_c^\infty[1]$; the multiplication commutator is $iX_j(F_i)$ in the Schrödinger package. (3) By the universal property of $\mathcal{W}$ [2] it is enough to check the images of generators satisfy the same relations. Directly (no Poisson-to-commutator dictionary): with left-ordered coefficients,

$$[\psi(q_i), \psi(p_j)] = X_j(F_i) = \delta_{ij}, \quad [\psi(p_i), \psi(p_j)] = \text{the first-order operator associated to } [X_i, X_j] = 0.$$

(The identities are algebraic in the Weyl algebra and match the dual-CAS generator checks in [8].) $\square$

Remark 2. ψ is not claimed to extend to a unitary conjugation on L^2 , a C^* automorphism, or a quantum channel.

6 Theorem D — incompleteness of X_1

Theorem 4 (D). The dual field X_1 is incomplete on $\mathbb{R}^3$.

Proof. For $t \in [0, \frac{1}{2})$, with continuous extension $\gamma(0) = (1, 0, 0)$, set

$$\begin{aligned} q_0(t) &= \frac{-2t - \sqrt{1-2t} + 1}{t(2t-1)}, \\ q_1(t) &= t, \\ q_2(t) &= t^2(2t - 3\sqrt{1-2t} - 1) \end{aligned}$$

on $(0, \frac{1}{2})$, and $\gamma(t) = (q_0(t), q_1(t), q_2(t))$. Then γ is C^∞ on $(0, \frac{1}{2})$ and extends C^1 to $t = 0$. Computer algebra gives the polynomial/algebraic identity $F(\gamma(t)) = (0, t, 2)$ on $(0, \frac{1}{2})$ [8]. Differentiating yields $DF_{\gamma(t)}\gamma'(t) = e_1$. Since $DF \cdot X_1 = e_1$ uniquely, $\gamma' = X_1 \circ \gamma$.

As $t = \frac{1}{2} - \varepsilon$ with $\varepsilon \downarrow 0$,

$$q_0(t) = \sqrt{2/\varepsilon} + O(1) \rightarrow +\infty,$$

so $\gamma(t)$ leaves every compact of $\mathbb{R}^3$ in finite time. Hence X_1 is incomplete. $\square$

7 Flow-box coordinates

Along the X_1 flow (where defined), $X_1 F_0 = X_1 F_2 = 0$ and $X_1 F_1 = 1$. On any open set carrying a smooth local inverse of F (locally everywhere: $\det DF \equiv -2 \neq 0$), the coordinates $(a, s, c) = F(q)$ satisfy

$$dq = \frac{1}{|\det DF|} da ds dc = \frac{1}{2} da ds dc, \quad X_1 = \partial_s, \quad H := -iX_1 = -i\partial_s. \quad (7.1)$$

Indeed, if Ψ is a local inverse sheet, $D\Psi = (DF)^{-1}$ and

$$|\det D\Psi| = \frac{1}{|\det DF|} = \frac{1}{2},$$

so Lebesgue measure pushes forward with factor $\frac{1}{2}$ (orientation chosen so the factor is positive).

Eliminating q_2 from $F_2 = c$ and q_1 from $(F_0, F_1) = (a, s)$ yields

$$A(s; a, c) q_0^3 + B(s; c) q_0 + C(c) = 0, \quad (7.2)$$

with

$$A = -cs^3 + s^2 + 18acs - 27a^2c^2 - 16a, \quad B = 3cs - 4, \quad C = 2c.$$

If $A \rightarrow 0$ with $B \neq 0$, Vieta forces some $|q_0| \rightarrow \infty$.

Reconstruction. If $x = q_0 \neq 0$ solves (7.2) and the denominators below are nonzero,

$$\begin{aligned} q_1 &= \frac{9acx^2 - 3csx - 3c - sx^2 + 6x}{x((3cs - 4)x + 3c)}, \\ q_2 &= \frac{2x - 3x^2q_1 - c}{x^3}, \end{aligned} \tag{8.0}$$

then $F(x, q_1, q_2) = (a, s, c)$ by polynomial identity.

8 Escape walls and deficiency indices

Write

$$n_+ := \dim \ker(H^* - i), \quad n_- := \dim \ker(H^* + i).$$

8.1 Per-orbit dictionary

On $L^2((\ell, r), ds)$ with $h = -i\partial_s$ minimal on $C_c^\infty(\ell, r)$ [1]: finite upper end $\Rightarrow \dim \ker(h^* + i) = 1$ (model e^{s-r}); finite lower end $\Rightarrow \dim \ker(h^* - i) = 1$ (model $e^{-(s-\ell)}$).

8.2 Forward wall

Proposition 1 (Forward wall). *There exist a nonempty open $U_+ \subset \mathbb{R}^2$ and $\beta \in C^\infty(U_+)$ such that for every $(a, c) \in U_+$ a selected real inverse branch of F exists on an s -interval with finite upper end $s = \beta(a, c)$ and $\|q\| \rightarrow \infty$ as $s \rightarrow \beta^-$.*

Proof. At the base $p_\star = (a, s, c) = (0, \frac{1}{2}, 2)$: $A(p_\star) = 0$, $A_s := \partial_s A(p_\star) = -\frac{1}{2} \neq 0$, $B(p_\star) = -1 \neq 0$. By the implicit function theorem there is a neighborhood of $(0, 2)$ and $\beta \in C^\infty$ with $\beta(0, 2) = \frac{1}{2}$ and $A(\beta(a, c); a, c) = 0$.

Set $\tau = \sqrt{\beta - s}$ and $q_0 = r/\tau$. Define

$$G_+(a, c, \tau, r) := \frac{A(\beta - \tau^2; a, c)}{\tau^2} r^3 + B(\beta - \tau^2; c) r + C(c) \tau \quad (\tau \neq 0),$$

and extend at $\tau = 0$ by

$$G_+(a, c, 0, r) = (-A_s(\beta; a, c))r^3 + B(\beta; c)r.$$

Smoothness of the extension follows from $A(\beta) = 0$ and the Taylor expansion $A(\beta - \tau^2) = -A_s(\beta)\tau^2 + O(\tau^4)$ in τ^2 . At the base, $G_+(0, 2, 0, r) = \frac{1}{2}r^3 - r$ has simple roots $r = \pm\sqrt{2}$ because $\partial_r G_+ = \frac{3}{2}r^2 - 1$ and $\partial_r G_+(\pm\sqrt{2}) =$

$-2B = 2 \neq 0$. The IFT continues $r_{\pm}(a, c, \tau)$ on an open set $U_+ \times (0, \tau_0)$. Then $|q_0| = |r_{\pm}|/\tau \rightarrow \infty$ as $\tau \downarrow 0$.

For reconstruction, the denominator in (8.0) behaves as $q_0(Bq_0 + 3c) \sim Bq_0^2$ with $B \rightarrow -1 \neq 0$, hence is nonzero for small τ after shrinking U_+ . The formulas for q_1, q_2 are rational in (q_0, s, a, c) with this nonvanishing denominator; combined with $q_0 \sim r/\tau$ one obtains the spatial escape lower bound

$$\|q\| \geq |q_0| \geq \frac{c_0}{\sqrt{\beta - s}}$$

for some $c_0 > 0$ on a possibly smaller relatively compact $W_+ \Subset U_+$ and s near β . Regression checks: `data/anchor/cas_orbit_measure_IFT_A001.json` [8]. $\square$

8.3 Backward wall

Proposition 2 (Backward wall). *There exist a nonempty open U_- and $\alpha \in C^\infty(U_-)$ such that for $(a, c) \in U_-$ a selected real branch exists for $s > \alpha(a, c)$ with $\|q\| \rightarrow \infty$ as $s \downarrow \alpha^+$.*

Proof. At $a = \frac{1}{54}$, $c = 2$,

$$A(s; \frac{1}{54}, 2) = -\frac{(2s-1)(3s^2-1)}{3},$$

so $A(\frac{1}{2}) = 0$, $A_s(\frac{1}{2}) = \frac{1}{6} > 0$, $B(\frac{1}{2}) = -1 \neq 0$. By the IFT there is a neighborhood of $(\frac{1}{54}, 2)$ and $\alpha \in C^\infty$ with $\alpha(\frac{1}{54}, 2) = \frac{1}{2}$ and $A(\alpha(a, c); a, c) = 0$.

Set $\tau = \sqrt{s - \alpha}$ (so $s = \alpha + \tau^2$) and $q_0 = r/\tau$. Define

$$G_-(a, c, \tau, r) := \frac{A(\alpha + \tau^2; a, c)}{\tau^2} r^3 + B(\alpha + \tau^2; c) r + C(c) \tau \quad (\tau \neq 0),$$

and extend at $\tau = 0$ by

$$G_-(a, c, 0, r) = A_s(\alpha; a, c) r^3 + B(\alpha; c) r.$$

Smoothness of the extension follows from $A(\alpha) = 0$ and $A(\alpha + \tau^2) = A_s(\alpha)\tau^2 + O(\tau^4)$. At the base, $G_-(\frac{1}{54}, 2, 0, r) = \frac{1}{6}r^3 - r$ has simple roots $r = \pm\sqrt{6}$ because $\partial_r G_- = \frac{1}{2}r^2 - 1$ and $\partial_r G_-(\pm\sqrt{6}) = -2B = 2 \neq 0$. The IFT continues $r_{\pm}(a, c, \tau)$ on an open set $U_- \times (0, \tau_0)$, with $r_{\pm} \neq 0$ after shrinking. Then $|q_0| = |r_{\pm}|/\tau \rightarrow \infty$ as $\tau \downarrow 0$.

For reconstruction, the denominator in (8.0) behaves as $q_0(Bq_0 + 3c) \sim Bq_0^2$ with $B \rightarrow -1 \neq 0$, hence is nonzero for small τ after shrinking U_- . Combined with $q_0 \sim r/\tau$ one obtains

$$\|q\| \geq |q_0| \geq \frac{c'_0}{\sqrt{s - \alpha}}$$

on a relatively compact $W_- \Subset U_-$ near the wall. Regression: data/anchor/cas_backward_incomplete. □

8.4 Saturated maximal sheets

Lemma 1 (Saturated maximal sheets). *Let $W_+ \Subset U_+$ be nonempty, open, and relatively compact. On the selected large- q_0 forward branch, pick $\varepsilon_0 > 0$ small and the smooth cross-section*

$$\Sigma_+ = \{q : F(q) = (a, \beta(a, c) - \varepsilon_0, c), (a, c) \in W_+\}.$$

Let ϕ_t be the maximal flow of X_1 . Because $X_1 F_1 = 1$, $F_1(\phi_t(q)) = F_1(q) + t$ on the existence interval. Define

$$D_+ = \{(a, c, s) : (a, c) \in W_+, \ell_+(a, c) < s < \beta(a, c)\},$$

where (ℓ_+, β) is the maximal open s -interval of the selected immersed branch. Then

$$\Psi_+ : D_+ \rightarrow \Omega_+ \subset \mathbb{R}^3, \quad \Psi_+(a, c, s) = \phi_{s-(\beta-\varepsilon_0)}(q_\Sigma(a, c))$$

is a C^∞ diffeomorphism onto an open X_1 -invariant set, $F \circ \Psi_+ = (a, s, c)$, and $s \rightarrow \beta^-$ occurs at spatial infinity. If ℓ_+ is finite, maximality forces escape as $s \downarrow \ell_+$ as well. An analogous Ψ_-, Ω_- exists for the lower wall.

Proof. The maximal-flow domain $\mathcal{D} \subset \mathbb{R} \times \mathbb{R}^3$ of the smooth vector field X_1 is open and the flow ϕ is C^∞ on $\mathcal{D}$ [3]. The set D_+ is (up to the smooth reparameterization $s = F_1$) the pullback of $\mathcal{D}$ through the cross-section Σ_+ ; openness of $\mathcal{D}$ therefore yields openness of D_+ . Uniqueness of ODE solutions gives injectivity of Ψ_+ . Since $F \circ \Psi_+ = (a, s, c)$ identically on D_+ and $\det DF \neq 0$, Ψ_+ is a local diffeomorphism, hence a C^∞ diffeomorphism onto its open image Ω_+ . The upper end is the wall of the forward-wall proposition, hence not attained in $\mathbb{R}^3$. Finite lower maximal time forces escape from compacts by standard ODE maximality. □

8.5 Deficiency vectors in $\text{Dom}(H^*)$

Lemma 2 (Deficiency vectors). *For $\chi \in C_c^\infty(W_+)$ define on the entire maximal interval*

$$u_-(\Psi_+(a, c, s)) := \chi(a, c) e^{s-\beta(a, c)},$$

and $u_- = 0$ off Ω_+ . No interior s -cutoff. Then $u_- \in L^2(\mathbb{R}^3)$, $u_- \in \text{Dom}(H^)$, and $(H^* + i)u_- = 0$. The linear map $\chi \mapsto u_-$ is injective, so $n_- = \infty$. Likewise*

$$u_+(\Psi_-(a, c, s)) := \chi(a, c) e^{-(s-\alpha(a, c))}$$

on (α, r_-) yields $n_+ = \infty$.

Proof. Square-integrability. By (7.1),

$$\|u_-\|_2^2 = \frac{1}{2} \int_{W_+} |\chi|^2 \left(\int_{\ell_+}^\beta e^{2(s-\beta)} ds \right) da dc \leq \frac{1}{4} \|\chi\|_{L^2(W_+)}^2,$$

since the inner integral equals $\frac{1}{2}(1 - e^{2(\ell_+ - \beta)}) \leq \frac{1}{2}$.

Adjoint equation. Let $\varphi \in C_c^\infty(\mathbb{R}^3)$. On Ω_+ , $H = -i\partial_s$ in the flow-box chart with density $\frac{1}{2}$, and $\text{div } X_1 = 0$, so the formal adjoint of $-iX_1$ is again $-iX_1$. There is no interior jump because u_- has no artificial s -cutoff. Off Ω_+ , $u_- = 0$. Lateral boundaries of W_+ contribute nothing because $\chi|_{\partial W_+} = 0$.

For the upper end: $s \rightarrow \beta^-$ occurs at spatial infinity, so φ vanishes in a neighborhood of that end along each orbit it meets. For the lower end: if ℓ_+ is finite then maximality forces escape to infinity as $s \downarrow \ell_+$, and again φ vanishes near that end; if $\ell_+ = -\infty$, the factor $e^{s-\beta}$ decays exponentially as $s \rightarrow -\infty$, so both the integrated boundary term at $-\infty$ and the L^2 tail vanish (no appeal to a compact lower s -cut is needed). Integrating by parts in s for almost every (a, c) (Fubini) therefore yields

$$\langle u_-, H\varphi \rangle = \langle -iu_-, \varphi \rangle,$$

i.e. $u_- \in \text{Dom}(H^*)$ and $(H^* + i)u_- = 0$.

Infinite dimension. If $u_- = 0$ a.e., then $\chi = 0$ a.e. on W_+ because Ψ_+ is a diffeomorphism and the exponential factor never vanishes. Thus $\chi \mapsto u_-$ embeds an infinite-dimensional space into $\ker(H^* + i)$.

The lower-wall argument is symmetric. $\square$

Remark 3 (Withdrawn device). *Interior indicators $\mathbf{1}_{(\beta-\delta, \beta)}(s)$ create δ -masses and exit $\text{Dom}(H^*)$. That construction is withdrawn.*

Theorem 5 (F). $(n_+, n_-) = (\infty, \infty)$.

Proof. Forward and backward walls, saturation, and the deficiency lemma. Separability of $L^2(\mathbb{R}^3)$ makes both dimensions countably infinite. $\square$

Theorem 6 (E). $H = -iX_1$ is not essentially self-adjoint on $C_c^\infty(\mathbb{R}^3)$.

Proof. $n_\pm \geq 1$ by Theorem F. $\square$

Corollary 1. *By von Neumann's theorem, infinitely many self-adjoint extensions of H exist. Every such extension agrees with the same minimal operator on $C_c^\infty(\mathbb{R}^3)$; the core algebraic Poisson/CCR relations recorded in Theorems B–C therefore do not, by themselves, distinguish a unique extension.*

9 Discussion

The example separates algebraic and analytic layers. The Poisson cotangent lift and dual-field/CCR identities are clean, and ψ is a well-defined polynomial Weyl endomorphism. Nonetheless X_1 is incomplete and $H = -iX_1$ has deficiency indices (∞, ∞) . Algebraic CCR/Poisson data therefore do not select a unique self-adjoint dual momentum for this generator.

We do *not* claim here exact deficiency indices for H_0 or H_2 , nor a theorem classifying all strong CCR or dual-translation packages; those topics appear only in companion repository notes and may be incomplete relative to theorem grade.

Classical incomplete flows and half-line models are standard [1]. The contribution is the geometric dual-lift package for this Keller seed together with an explicit two-sided wall construction for $H = -iX_1$.

10 Non-claims

No unitary gate, channel, CP map, or computational advantage. No preferred physical extension. No claim that all quantization routes fail. Discovery of F not claimed. Lean: seed identities and algebraic dual-matrix packaging only [9]. No theorem-level dual-flow strong CCR obstruction in this note.

References

- [1] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*, Academic Press, 1975.
- [2] J. Dixmier, *Sur les algèbres de Weyl*, Bull. Soc. Math. France **96** (1968), 209–242.
- [3] J. M. Lee, *Introduction to Smooth Manifolds*, 2nd ed., Springer, 2013.
- [4] H. Bass, E. H. Connell, and D. Wright, The Jacobian conjecture: reduction of degree and formal expansion of the inverse, Bull. Amer. Math. Soc. **7** (1982), 287–330.
- [5] L. Alpöge, informal public announcement of a Keller map with constant Jacobian and three-point collision (credits Fable), July 2026, https://x.com/__alpoge__/status/2079028340955197566.
- [6] Ulam, *A Counterexample to the Jacobian Conjecture* (public write-up of the July 2026 announcement), <https://www.ulam.ai/research/jacobian.pdf>.
- [7] D. Speyer, *The new counterexample to the Jacobian conjecture*, Secret Blogging Seminar, 20 July 2026, <https://sbseminar.wordpress.com/2026/07/20/the-new-counterexample-to-the-jacobian-conjecture/>.
- [8] D. E. Fredriksen, EXOTIC-CCR A001 companion code and CAS certificates, GitHub `Quantlyra/jacobian-weyl-quantum-phase-space`, release `v0.3.5-submit`, concept DOI 10.5281/zenodo.21474351.
- [9] D. E. Fredriksen, `exotic-ccr-lean`, GitHub `Quantlyra/exotic-ccr-lean` (Lean 4 formalization of the seed identities and dual-matrix algebra for this companion).