

Dual-lift realizability for Keller-type maps: structural incompleteness, schema, and examples

Daniel Eric Fredriksen
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

We isolate a structural lemma for dual vector fields of local diffeomorphisms $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$: if global dual fields X_j with $DF \cdot X_j = e_j$ are all complete, then F is surjective. Hence any non-surjective dual system has at least one incomplete dual field. We propose a classification schema (B-incomplete / B-ESS / B-Def / B-CCR / B-CP) and populate it on atlas rows A000–A006 (pass and fail baselines, Keller seed A001, family pilot A002). No gates, channels, or computational advantage are claimed. Companion pack: <https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space/blob/main/docs/notes/COMPANION-PACK.md>.

MSC: 37C10, 47B25, 53D17, 14R15.

Version: B001 draft v0.2.

Software: atlas program_b; CAS cas_atlas_A005_A006_B001.json.

1 Introduction

Cotangent lifts and dual fields are standard when $\det DF \neq 0$. After the A001 deficiency analysis [1], we classify which maps force incomplete dual fields and how that interacts with ESS / CCR / CP packages [2]. A001 arXiv endorsement may be pending; the concept DOI still cites A001.

2 Structural lemma

Lemma 1 (Complete dual fields imply surjectivity). *Let $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be C^∞ and a local diffeomorphism with dual fields X_j , $DF \cdot X_j = e_j$. If every X_j is complete, then $F(\mathbb{R}^n) = \mathbb{R}^n$.*

Proof. $F(\phi_t^{(j)}(q)) = F(q) + te_j$; iterate; image open and translation-invariant. □

Corollary 1. *Non-surjective dual systems have some incomplete X_j .*

3 Schema

B-incomplete / B-ESS / B-Def / B-CCR / B-CP with values open—(fail—(pass.

4 Atlas scoreboard (v0.2)

ID	Map (sketch)	B-inc.	Role
A000	id	pass	baseline
A004	shear $(x, y + x, z)$	pass	linear auto
A005	diag $(2, \frac{1}{2}, 1)$	pass	linear stretch (CAS)
A003	(x, e^y) pad	fail	smooth B1 example
A006	(e^x, e^y, z)	fail	two incomplete fields (CAS)
A001	Keller seed	fail	maximal analytic [1]
A002	family d4 pilot	open	geometry pending

5 Examples (new in v0.2)

Example 1 (A005 — linear stretch, pass-like). $F = \text{diag}(2, \frac{1}{2}, 1)$, $\det DF = 1$. Dual fields $X_0 = \frac{1}{2}\partial_x$, $X_1 = 2\partial_y$, $X_2 = \partial_z$ are complete. CAS: *cas_atlas_A005_A006_B001.json*.

Example 2 (A006 — exp quadrant, fail-like). $F(x, y, z) = (e^x, e^y, z)$, $\det DF = e^{x+y} > 0$, image $(0, \infty)^2 \times \mathbb{R}$. Dual fields $X_0 = e^{-x}\partial_x$, $X_1 = e^{-y}\partial_y$ incomplete backward ($e^x = e^{x_0} + t$). Matches Lemma 1. CAS same file.

6 Prior examples (v0.1)

A000 identity; A003 half-plane; A001 Keller+ $(n_+, n_-) = (\infty, \infty)$ for $H = -iX_1$ [1]; A002 open; A004 shear pass.

7 Non-claims

No gates, channels, advantage. No dual-flow repair. A003/A005/A006 are not Jacobian counterexamples. T4: no dual F -translation package claimed here. C Free-Strict-Abstract remains open [2].

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., v0.3.6-submit / concept DOI 10.5281/zenodo.21474351, <https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>.
- [2] D. E. Fredriksen, CP readings of an exotic Weyl endomorphism..., v0.2.0-c001-free-strict, <https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>.