

Completely positive readings of an exotic Weyl endomorphism: algebraic correspondence and standard-package obstructions

Daniel Eric Fredriksen
Quanyra Inc.

<https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

Let ψ be the polynomial Weyl endomorphism of the Alpöge–Fable seed (restated): $Q = F(q)$, $P = J^{-T}p$. With the physics normalization $[q_i, p_j] = i\delta_{ij}$ and $q_i^* = q_i$, $p_j^* = p_j$, ψ is a unital $*$ -endomorphism (algebraic SOS positivity). We construct its algebraic endomorphism correspondence and obstruct: (i) unital/nondegenerate $*$ -endomorphisms of $C_0(\mathbb{R}^3)$ extending composition with non-proper F ; (ii) Bogoliubov/quasifree endomorphisms of the CCR algebra (nonlinear F). A former “Free-Strict-Regular-v2” nondegeneracy obstruction is **withdrawn** as incorrect. Fully abstract C^* CP+Stinespring without spatial axioms remains open. No dual F -translations, gates, or advantage. Companion to [1]. Pack: COMPANION-PACK.md.

MSC: 46L07, 81S05, 47L90, 46L08.

Version: C001 companion v0.3 (adversarial REVISE).

T4: no dual F -translations on $L^2(\mathbb{R}^3)$.

Erratum. Free-Strict-Regular-v2 (projection off $\overline{\text{im } F}$) is **withdrawn**: $q \notin \overline{\text{im } F}$ does not imply $f(F(q)) = 0$, and $M_f \mapsto M_{f \circ F}$ on $C_b(Q)$ is a valid unital $*$ -homomorphism extendable by Arveson. Do not cite v0.2 Regular-v2 as a theorem.

1 Domain

Definition 1. $\mathcal{W}$: polynomial Weyl algebra over $\mathbb{C}$ with $[q_i, p_j] = i\delta_{ij}$, $[q_i, q_j] = [p_i, p_j] = 0$, and involution $q_i^* = q_i$, $p_j^* = p_j$. (Consistent: $[q, p]^* = -[q, p]$ and $(i\delta)^* = -i\delta$.) With real $B = J^{-T}$,

$$\psi(q_i) = F_i(q), \quad \psi(p_j) = \sum_k B_{jk}(q) p_k$$

(coefficients left of momenta) extends to a unital algebra endomorphism.

Remark 1 (Normalization). Earlier drafts used $[q, p] = \delta$ with self-adjoint generators, which is inconsistent under $*$. This note uses the standard physics normalization $[q, p] = i$.

Theorem 1 (Algebraic $*$ -SOS). Real $F, B \Rightarrow \psi$ is a unital $*$ -endomorphism of $\mathcal{W}$, hence $\text{id}_n \otimes \psi$ preserves hermitian-square cones.

Definition 2 (Algebraic correspondence). Let $E = \mathcal{W}$ as a right $\mathcal{W}$ -module by $\xi \cdot b = \xi b$, with left action $a \cdot \xi = \psi(a)\xi$ and $\langle \xi, \eta \rangle_{\mathcal{W}} = \xi^* \eta$.

2 Geometry

Theorem 2 (Non-proper). *F is not proper along the incomplete X_1 -orbit of [1] (CAS `cas_F_not_proper_C1c_A001`).*

3 Obstructions (retained)

Theorem 3 (C0 composition endo). *There is no $*$ -homomorphism $\alpha : C_0(\mathbb{R}^3) \rightarrow C_0(\mathbb{R}^3)$ (nondegenerate/approximately unital in the usual C_0 sense) with $\alpha(f) = f \circ F$ for all $f \in C_0(\mathbb{R}^3)$, because F is not proper so $f \circ F \notin C_0$ for some $f \in C_0$.*

Theorem 4 (Bogoliubov / quasifree). *No Bogoliubov (linear quasifree) endomorphism of the CCR C^* -algebra restricts to ψ on the polynomial generators: such maps act linearly on one-particle fields, while F is nonlinear of degree 3.*

Remark 2 (Not claimed). *We do not claim every continuous unital $*$ -endomorphism of the CCR algebra fails—only the Bogoliubov/quasifree class. We do not claim that $f \mapsto M_{f \circ F}$ fails as a map $C_b \rightarrow B(L^2)$.*

4 Withdrawn

Free-Strict-Regular-v2 nondegeneracy NO-GO: **withdrawn** (see erratum).

5 Open

Unital CP+Stinespring on a fully abstract C^* envelope of $\mathcal{W}$ with no C0-composition and no Bogoliubov axioms: not constructed (beyond algebraic correspondence and finite-level algebraic CP), not ruled out.

6 Non-claims

No channel on $B(H)$. No gates/advantage. No dual F -translations. No “all quantization fails.” No full abstract C^* Stinespring construct.

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., v0.3.6-submit, <https://github.com/Quanytyra/jacobian-weyl-quantum-phase-space>.