

# Dual-lift realizability for Keller-type maps: structural incompleteness, schema, and examples

Daniel Eric Fredriksen  
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

21 July 2026

## Abstract

We isolate a structural lemma for dual vector fields of local diffeomorphisms  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ : if global dual fields  $X_j$  with  $DF \cdot X_j = e_j$  are all complete, then  $F$  is surjective. Hence any non-surjective dual system has at least one incomplete dual field. We propose a classification schema (B-incomplete / B-ESS / B-Def / B-CCR / B-CP) and populate it on atlas rows A000–A006 (pass and fail baselines, Keller seed A001, family pilot A002). No gates, channels, or computational advantage are claimed. Companion pack: <https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space/blob/main/docs/notes/COMPANION-PACK.md>.

**MSC:** 37C10, 47B25, 53D17, 14R15.

**Version:** B001 draft v0.2.

**Software:** atlas program\_b; CAS cas\_atlas\_A005\_A006\_B001.json.

## 1 Introduction

Cotangent lifts and dual fields are standard when  $\det DF \neq 0$ . After the A001 deficiency analysis [1], we classify which maps force incomplete dual fields and how that interacts with ESS / CCR / CP packages [2]. A001 arXiv endorsement may be pending; the concept DOI still cites A001.

## 2 Structural lemma

**Lemma 1** (Complete dual fields imply surjectivity). *Let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be  $C^\infty$  and a local diffeomorphism with dual fields  $X_j$ ,  $DF \cdot X_j = e_j$ . If every  $X_j$  is complete, then  $F(\mathbb{R}^n) = \mathbb{R}^n$ .*

*Proof.*  $F(\phi_t^{(j)}(q)) = F(q) + te_j$ ; iterate; image open and translation-invariant. □

**Corollary 1.** *Non-surjective dual systems have some incomplete  $X_j$ .*

## 3 Schema

B-incomplete / B-ESS / B-Def / B-CCR / B-CP with values `open` / `fail` / `pass`.

## 4 Atlas scoreboard (v0.2)

| ID   | Map (sketch)               | B-inc. | Role                        |
|------|----------------------------|--------|-----------------------------|
| A000 | id                         | pass   | baseline                    |
| A004 | shear $(x, y + x, z)$      | pass   | linear auto                 |
| A005 | diag $(2, \frac{1}{2}, 1)$ | pass   | linear stretch (CAS)        |
| A003 | $(x, e^y)$ pad             | fail   | smooth B1 example           |
| A006 | $(e^x, e^y, z)$            | fail   | two incomplete fields (CAS) |
| A001 | Keller seed                | fail   | $X_1 + H_1$ analytic [1]    |
| A002 | family d4 pilot            | open   | geometry pending            |

## 5 Examples (new in v0.2)

**Example 1** (A005 — linear stretch, pass-like).  $F = \text{diag}(2, \frac{1}{2}, 1)$ ,  $\det DF = 1$ . *Dual fields*  $X_0 = \frac{1}{2}\partial_x$ ,  $X_1 = 2\partial_y$ ,  $X_2 = \partial_z$  are complete. *CAS*: `cas_atlas_A005_A006_B001.json`.

**Example 2** (A006 — exp quadrant, fail-like).  $F(x, y, z) = (e^x, e^y, z)$ ,  $\det DF = e^{x+y} > 0$ , *image*  $(0, \infty)^2 \times \mathbb{R}$ . *Dual fields*  $X_0 = e^{-x}\partial_x$ ,  $X_1 = e^{-y}\partial_y$  incomplete backward ( $e^x = e^{x_0} + t$ ). *Matches Lemma 1. CAS same file.*

## 6 Prior examples (v0.1)

A000 identity; A003 half-plane; A001 Keller seed with  $X_1$  incomplete and  $H = -iX_1$  deficiency  $(\infty, \infty)$  in [1] (other  $X_j$  incompleteness is recorded in companion validation dossiers, not re-proved here); A002 open; A004 shear pass.

## 7 Non-claims

No gates, channels, advantage. No dual-flow repair. A003/A005/A006 are not Jacobian counterexamples. T4: no dual  $F$ -translation package claimed here. C Free-Strict-Abstract remains open [2].

## References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., v0.3.6-submit / concept DOI 10.5281/zenodo.21474351, <https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>.
- [2] D. E. Fredriksen, CP readings of an exotic Weyl endomorphism..., v0.3.0-c001-companion, <https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>.