

Poisson and Weyl lifts of an explicit three-dimensional Keller map and nonunique self-adjoint realizations of a dual transport operator

Daniel Eric Fredriksen
Quanyra Inc.

<https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>

29 July 2026

Abstract

We study the three-dimensional Keller map F announced by Alpöge, whose announcement credits Fable for work leading to the map: $\det DF \equiv -2$ and a three-point collision. Theorem A restates those finite identities. Theorems B–C record the standard cotangent/Piola/Weyl package for this seed; the analytic contribution is Theorems D–F. The cotangent lift $\Phi(q, p) = (F(q), J^{-T}p)$ is a polynomial Poisson map and non-injective. Dual fields $X_j = \sum_k B_{jk} \partial_{q_k}$ commute and are divergence-free (Piola), so $H_j = -iX_j$ are symmetric on $C_c^\infty(\mathbb{R}^3)$. We define the induced polynomial Weyl endomorphism ψ , without claiming that ψ is noninvertible or not an automorphism. The field X_1 is incomplete via an explicit integral curve. In flow-box coordinates $(a, s, c) = F(q)$ one has $X_1 = \partial_s$ and $dq = \frac{1}{2} da ds dc$. An open forward escape wall yields whole-orbit vectors in $\ker(H^* + i)$ (transverse cutoffs only; no interior s -indicators); a measure-preserving sign involution transports them isometrically to $\ker(H^* - i)$. Thus the Hilbert deficiency indices are

$$(n_+, n_-) = (\aleph_0, \aleph_0),$$

equivalently (∞, ∞) in conventional deficiency-index notation, for $H = -iX_1$. A continuum-sized family of distinct self-adjoint extensions exists; the core algebraic CCR/Poisson data alone do not distinguish among them. The operator-theory mechanism is classical; the contribution is its explicit realization for this dual field. No gates, channels, computational advantage, or seed discovery claim. Paper-package release target **v0.3.9-referee-revision**.

MSC 2020: 47B25, 81S05, 14R15, 37C10, 53D17.

Paper package: v0.3.9-referee-revision; concept DOI

<https://doi.org/10.5281/zenodo.21474351> (project-level; any version DOI is recorded separately by release ingest).

Categories (suggested): math.SP (primary); math.FA, math-ph (secondary).

Lean coverage boundary. The companion Lean artifact proves the bounded canonical-core declaration `ExoticCCR.theoremE`: the minimal transport core for X_1 is not essentially self-adjoint via a constructed weak $-i$ witness. The historical release tag **v0.1.8-theorem-e** targets commit **be4f330**; **a6bb091** is the reviewed Theorem E source anchor within that release history, not the tag target. The synchronized Lean publication freeze is **2e40c4cab86a1ef97cb3334497d10081dfe33867**; it is untagged and unreleased. This is a provenance-only successor to **fb added034**; theorem and audit source are unchanged. The earlier commit **ff50f4a2a312591c2e5b26e71eb390ade9164b34** is retained only as the historical Theorem F source root. At the synchronized freeze the artifact proves the bounded Theorem F declaration `ExoticCCR.theoremF`: both adjoint eigenspaces

$\ker(H^\dagger \mp i)$ of the canonical minimal core are not finite-dimensional and the bounded reading of $(n_+, n_-) = (\infty, \infty)$. It also proves $\aleph_0$ lower bounds for the corresponding `Module.rank`-valued algebraic/Hamel ranks. The Lean route to the $+i$ side is a measure-preserving sign-involution transport $(\sigma(q_0, q_1, q_2) = (-q_0, -q_1, q_2), X_1 \circ \sigma = -\sigma_* X_1)$ of the forward minus- i families, not this paper’s backward-wall construction; the same SHA separately defines `hilbertDeficiencyIndex` as the cardinality of a chosen Hilbert basis for each closed adjoint eigenspace and proves the exact-index application theorem `hilbertDeficiencyIndex.X1.eq.aleph0`; thus the exact Hilbert-space indices are $\aleph_0$ on both sides. This is distinct from `hamelDeficiencyRank`, which is `Module.rank` (algebraic/Hamel rank) and has lower bounds only. For the specific canonical minimal operator `H.X1.min`, the same SHA proves a bijective von Neumann classification of all `SelfAdjointExtension H.X1.min` witnesses (self-adjoint `LinearPMaps` extending `H.X1.min`) by complex-linear isometric equivalences from the $+i$ adjoint eigenspace to the $-i$ adjoint eigenspace. The declarations `theoremFUnitPhaseExtension` and `theoremFUnitPhaseExtension.injective` prove that unitary complex phases inject into distinct such witnesses. The further conclusion that this is a continuum-sized lower family uses the classical cardinality of the complex unit circle; that cardinal identification is not part of these Lean declarations [16].

The paper’s independent backward-wall proposition, the exact cardinality of the full extension type, preferred extension, arbitrary-operator classification, and algebraic/physical selection claims remain outside the Lean corollary. Formal symmetry of the specific canonical minimal core `H.X1.min` on its C_c^∞ test-function domain is separately Lean-formalized at Stage A SHA `bc679df` in `TheoremFSymmetricCore`, with the corresponding graph-space foundations in `TheoremFVonNeumannGraph`. This is not an arbitrary-operator or arbitrary-transport-realization symmetry theorem.

Errata (prior releases). v0.2.2 pair $(0, \infty)$ for H withdrawn. v0.3.0 interior s -indicator deficiency proof withdrawn ($\notin \text{Dom}(H^*)$). This submission is $H = -iX_1$ **only**: we do *not* claim exact deficiency pairs for H_0, H_2 , nor a theorem-level dual-flow strong CCR no-go (companion notes only).

1 Introduction

On 19–20 July 2026, Levent Alpöge announced an explicit polynomial map $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ with $\det DF \equiv -2$ and a three-point collision. The announcement credits Akhil Mathew with prompting the question and Claude Fable with work leading to the map [8, 10]. **We do not claim discovery of F .** The finite seed identities are *verified* here by dual CAS and by a companion Lean 4 formalization [15, 14]; an independent Isabelle/HOL development verifies the determinant and collision and records the same attribution [10]. Our contribution is the dual-lift deficiency analysis (Theorems D–F), not the Jacobian-conjecture refutation itself.

We ask whether the standard Poisson/CCR lifts of F select a unique self-adjoint dual transport generator on $C_c^\infty(\mathbb{R}^3)$. For $H = -iX_1$ the answer is negative: deficiency indices (∞, ∞) .

Scope. Theorems A–F below concern the algebra of the lifts and the analytic deficiency theory of $H = -iX_1$. Remarks on other dual fields or alternative realizations are deferred to companion validation notes and are not claimed as theorems here.

Algebraic and geometric context

The cotangent, Piola, and Weyl constructions in Theorems B–C are standard once a Keller map is given; they are included to fix conventions and connect the seed to the analytic problem, not claimed

as a new construction. Their Weyl-algebra setting is related to the Dixmier conjecture; for classical Jacobian-conjecture background see [7]. The classical same-rank implication is $\text{DC}_n \Rightarrow \text{JC}_n$, whereas Tsuchimoto and, independently, Belov–Kanel–Kontsevich proved the dimension-doubling converse $\text{JC}_{2n} \Rightarrow \text{DC}_n$; together these give stable equivalence [3, 4, 5]. In particular, one must not replace the latter by a dimension-preserving $\text{JC}_n \Rightarrow \text{DC}_n$. We do not derive or claim here a new rank-three Dixmier counterexample, nor do we claim that the explicitly defined ψ is noninvertible.

The July 2026 follow-on literature is developing rapidly. The Ulam preprint analyzes the seed’s fibers and nonproperness [9]; Shaska studies its mixed-sign grading, quotient geometry, discriminant, and nonproperness mechanism [11]; Speyer records an early public algebraic discussion [13]; and the Isabelle/AFP entry supplies an independent formal check of the finite seed identities [10]. These recent sources are cited as current preprints or formal artifacts, not as completed peer review. More generally, nonproperness sets of generically finite polynomial maps have an established literature [12]. The escape wall below is an operator-theoretic use of explicit behavior at infinity; it does not purport to classify the full nonproperness set.

2 Setup

Let $q = (q_0, q_1, q_2) = (x, y, z)$ and

$$\begin{aligned} F_0 &= (1 + xy)^3 z + y^2(1 + xy)(4 + 3xy), \\ F_1 &= y + 3x(1 + xy)^2 z + 3xy^2(4 + 3xy), \\ F_2 &= 2x - 3x^2 y - x^3 z. \end{aligned}$$

Write $J = DF$ and $B := J^{-T}$. Dual fields:

$$X_j := \sum_{k=0}^2 B_{jk}(q) \partial_{q_k}, \quad j = 0, 1, 2.$$

Normalizations. We use two compatible packages. (i) *Abstract Weyl algebra* $\mathcal{W}$: generators q_k, p_ℓ with $[q_i, p_j] = \delta_{ij}$ (Dixmier convention). (ii) *Schrödinger representation* on $L^2(\mathbb{R}^3)$: multiplication by q_k and $p_k^{\text{Sch}} = -i\partial_{q_k}$, so $[q_i, p_j^{\text{Sch}}] = i\delta_{ij}$. The operators $H_j := -iX_j$ live in package (ii). When $\text{div } X_j = 0$, each H_j is symmetric on $C_c^\infty(\mathbb{R}^3)$. Our complex inner products are conjugate-linear in the first argument and linear in the second. Collision data ($\text{char} \neq 2$):

$$F(0, 0, -\tfrac{1}{4}) = F(1, -\tfrac{3}{2}, \tfrac{13}{2}) = F(-1, \tfrac{3}{2}, \tfrac{13}{2}) = (-\tfrac{1}{4}, 0, 0). \quad (1)$$

3 Theorem A — seed identities

Theorem 1 (A). (1) $\det DF \equiv -2$. (2) The collision identities (1) hold over any field of characteristic not 2. (3) Hence F is not injective.

Proof. Expand $\det J$ as a polynomial identity, or evaluate the formal determinant in a computer algebra system; formalized in Lean 4 in the companion repository (`ExoticCCR/AnchorF.lean: jacobianDet_F, evalMap_F_p*`) [15]. Non-injectivity is immediate from (1). $\square$

Remark 1 (Complex vs real). The seed is stated algebraically over $\mathbb{C}^3$, but all three collision points in (1) are rational and hence real. Thus the same polynomial formulas restrict to a noninjective local diffeomorphism of $\mathbb{R}^3$. We treat F as an externally announced seed [8, 9, 10]; Theorem A is a restatement with CAS/Lean re-verification, not a priority claim. Theorems D–F concern this real restriction and the associated $L^2(\mathbb{R}^3)$ dual-field package.

4 Theorem B — Poisson cotangent lift

Theorem 2 (B). *Let $\Phi(q, p) = (F(q), B(q)p)$ with $B = J^{-T}$. Then:*

1. *B is polynomial and $JB^T = I$;*

2. *writing $Q_i = F_i(q)$ and $P_j = \sum_k B_{jk}(q)p_k$,*

$$\{Q_i, Q_j\} = 0, \quad \{P_i, P_j\} = 0, \quad \{Q_i, P_j\} = \delta_{ij}$$

for the canonical Poisson structure on $T^\mathbb{R}^3$;*

3. *Φ is not injective (lift the three collision sources with matching p).*

Proof. (1) follows from $B = J^{-T}$ and $\det J \equiv -2 \neq 0$; polynomiality of B is the classical adjugate formula $B = -(\text{adj } J)^T/2$. (2) The identities $\{Q_i, P_j\} = (JB^T)_{ij} = \delta_{ij}$ are immediate; vanishing of $\{Q_i, Q_j\}$ is clear; vanishing of $\{P_i, P_j\}$ is the standard cotangent-lift calculation for $B = J^{-T}$ (equivalently dual-CAS generator checks in the companion release [14]). (3) If $F(q^{(a)}) = Q_\star$ and $p^{(a)} = J(q^{(a)})^T P_\star$, then $\Phi(q^{(a)}, p^{(a)}) = (Q_\star, P_\star)$ for each collision source in (1). $\square$

5 Theorem C — dual fields, Piola, Weyl endomorphism

Theorem 3 (C). 1. $X_j(F_i) = \delta_{ij}$, $[X_i, X_j] = 0$, and $\text{div } X_j = 0$.

2. *Each $H_j = -iX_j$ is symmetric on $C_c^\infty(\mathbb{R}^3)$, and $[M_{F_i}, H_j] = i\delta_{ij}$ on that domain.*

3. *Let $\mathcal{W}$ be the polynomial Weyl algebra over $\mathbb{C}$ generated by q_k, p_ℓ with $[q_i, p_j] = \delta_{ij}$ (abstract package) and all other generators commuting. The assignment*

$$\psi(q_i) = F_i(q), \quad \psi(p_j) = \sum_k B_{jk}(q) p_k$$

extends uniquely to a unital algebra endomorphism $\psi : \mathcal{W} \rightarrow \mathcal{W}$ preserving the Weyl relations on generators (ordering: coefficients $B_{jk}(q)$ to the left of p_k).

Proof. (1) $X_j(F_i) = (JB^T)_{ij} = \delta_{ij}$. Then $[X_i, X_j](F_k) = 0$ for all k ; since dF is a coframe, $[X_i, X_j] = 0$. Piola: $\text{Div Cof}(DF) = 0$; with $\det DF \equiv -2$, $\text{Div } B = 0$, i.e. $\text{div } X_j = 0$. (2) Divergence-free smooth vector fields give symmetric $-iX$ on $C_c^\infty[1]$; the multiplication commutator is $iX_j(F_i)$ in the Schrödinger package. (3) By the universal property of $\mathcal{W}$ [2] it is enough to check the images of generators satisfy the same relations. Directly (no Poisson-to-commutator dictionary): with left-ordered coefficients,

$$\begin{aligned} [\psi(q_i), \psi(p_j)] &= X_j(F_i) = \delta_{ij}, \\ [\psi(p_i), \psi(p_j)] &= \text{the first-order operator of } [X_i, X_j] = 0. \end{aligned}$$

(The identities are algebraic in the Weyl algebra and match the dual-CAS generator checks in [14].) $\square$

Lean coverage for Theorem C(2). The all- j analytic symmetry and commutator assertion above is the classical paper-level consequence of the Lean-verified polynomial divergence and dual-field identities. The pinned operator-level Lean theorem `H_X1_min_isFormalAdjoint_H_X1_min` formalizes the specific $j = 1$ compactly supported core used in Theorems E–F; the freeze does not claim a named all- j operator theorem.

Remark 2. ψ is not claimed to extend to a unitary conjugation on L^2 , a C^* automorphism, or a quantum channel.

6 Theorem D — incompleteness of X_1

Theorem 4 (D). *The dual field X_1 is incomplete on $\mathbb{R}^3$.*

Proof. For $t \in [0, \frac{1}{2})$, with continuous extension $\gamma(0) = (1, 0, 0)$, set

$$\begin{aligned} q_0(t) &= \frac{-2t - \sqrt{1-2t} + 1}{t(2t-1)}, \\ q_1(t) &= t, \\ q_2(t) &= t^2(2t - 3\sqrt{1-2t} - 1) \end{aligned}$$

on $(0, \frac{1}{2})$, and $\gamma(t) = (q_0(t), q_1(t), q_2(t))$. Then γ is C^∞ on $(0, \frac{1}{2})$ and extends C^1 to $t = 0$. Computer algebra gives the polynomial/algebraic identity $F(\gamma(t)) = (0, t, 2)$ on $(0, \frac{1}{2})$ [14]. Differentiating yields $DF_{\gamma(t)}\gamma'(t) = e_1$. Since $DF \cdot X_1 = e_1$ uniquely, $\gamma' = X_1 \circ \gamma$.

As $t = \frac{1}{2} - \varepsilon$ with $\varepsilon \downarrow 0$,

$$q_0(t) = \sqrt{2/\varepsilon} + O(1) \rightarrow +\infty,$$

so $\gamma(t)$ leaves every compact of $\mathbb{R}^3$ in finite time. Hence X_1 is incomplete. $\square$

7 Flow-box coordinates

Along the X_1 flow (where defined), $X_1 F_0 = X_1 F_2 = 0$ and $X_1 F_1 = 1$. On any open set carrying a smooth local inverse of F (locally everywhere: $\det DF \equiv -2 \neq 0$), the coordinates $(a, s, c) = F(q)$ satisfy

$$dq = \frac{1}{|\det DF|} da ds dc = \frac{1}{2} da ds dc, \quad X_1 = \partial_s, \quad H := -iX_1 = -i\partial_s. \quad (7.1)$$

Indeed, if Ψ is a local inverse sheet, $D\Psi = (DF)^{-1}$ and

$$|\det D\Psi| = \frac{1}{|\det DF|} = \frac{1}{2},$$

so Lebesgue measure pushes forward with factor $\frac{1}{2}$ (orientation chosen so the factor is positive).

Eliminating q_2 from $F_2 = c$ and q_1 from $(F_0, F_1) = (a, s)$ yields

$$A(s; a, c) q_0^3 + B(s; c) q_0 + C(c) = 0, \quad (7.2)$$

with

$$A = -cs^3 + s^2 + 18acs - 27a^2c^2 - 16a, \quad B = 3cs - 4, \quad C = 2c.$$

If $A \rightarrow 0$ with $B \neq 0$, Vieta forces some $|q_0| \rightarrow \infty$.

Reconstruction. If $x = q_0 \neq 0$ solves (7.2) and the denominators below are nonzero,

$$\begin{aligned} q_1 &= \frac{9acx^2 - 3csx - 3c - sx^2 + 6x}{x((3cs - 4)x + 3c)}, \\ q_2 &= \frac{2x - 3x^2q_1 - c}{x^3}, \end{aligned} \quad (8.0)$$

then $F(x, q_1, q_2) = (a, s, c)$ by polynomial identity.

8 Escape walls and deficiency indices

Write

$$n_+ := \dim_{\mathbb{H}} \ker(H^* - i), \quad n_- := \dim_{\mathbb{H}} \ker(H^* + i),$$

where $\dim_{\mathbb{H}}$ denotes Hilbert dimension, i.e. orthonormal-basis cardinality, not algebraic/Hamel dimension.

8.1 Per-orbit dictionary

On $L^2((\ell, r), ds)$ with $h = -i\partial_s$ minimal on $C_c^\infty(\ell, r)$ [1]: finite upper end $\Rightarrow \dim_{\mathbb{H}} \ker(h^* + i) = 1$ (model e^{s-r}); finite lower end $\Rightarrow \dim_{\mathbb{H}} \ker(h^* - i) = 1$ (model $e^{-(s-\ell)}$).

8.2 Forward wall

Proposition 1 (Forward wall). *There exist a nonempty open $U_+ \subset \mathbb{R}^2$ and $\beta \in C^\infty(U_+)$ such that for every $(a, c) \in U_+$ a selected real inverse branch of F exists on an s -interval with finite upper end $s = \beta(a, c)$ and $\|q\| \rightarrow \infty$ as $s \rightarrow \beta^-$.*

Proof. At the base $p_\star = (a, s, c) = (0, \frac{1}{2}, 2)$: $A(p_\star) = 0$, $A_s := \partial_s A(p_\star) = -\frac{1}{2} \neq 0$, $B(p_\star) = -1 \neq 0$. By the implicit function theorem there is a neighborhood of $(0, 2)$ and $\beta \in C^\infty$ with $\beta(0, 2) = \frac{1}{2}$ and $A(\beta(a, c); a, c) = 0$.

Set $\tau = \sqrt{\beta - s}$ and $q_0 = r/\tau$. Define

$$G_+(a, c, \tau, r) := \frac{A(\beta - \tau^2; a, c)}{\tau^2} r^3 + B(\beta - \tau^2; c) r + C(c) \tau \quad (\tau \neq 0),$$

and extend at $\tau = 0$ by

$$G_+(a, c, 0, r) = (-A_s(\beta; a, c))r^3 + B(\beta; c)r.$$

Smoothness of the extension follows from $A(\beta) = 0$ and the Taylor expansion $A(\beta - \tau^2) = -A_s(\beta)\tau^2 + O(\tau^4)$ in τ^2 . At the base, $G_+(0, 2, 0, r) = \frac{1}{2}r^3 - r$ has simple roots $r = \pm\sqrt{2}$ because $\partial_r G_+ = \frac{3}{2}r^2 - 1$ and $\partial_r G_+(\pm\sqrt{2}) = -2B = 2 \neq 0$. The IFT continues $r_\pm(a, c, \tau)$ on an open set $U_+ \times (0, \tau_0)$. Then $|q_0| = |r_\pm|/\tau \rightarrow \infty$ as $\tau \downarrow 0$.

For reconstruction, the denominator in (8.0) behaves as $q_0(Bq_0 + 3c) \sim Bq_0^2$ with $B \rightarrow -1 \neq 0$, hence is nonzero for small τ after shrinking U_+ . The formulas for q_1, q_2 are rational in (q_0, s, a, c) with this nonvanishing denominator; combined with $q_0 \sim r/\tau$ one obtains the spatial escape lower bound

$$\|q\| \geq |q_0| \geq \frac{c_0}{\sqrt{\beta - s}}$$

for some $c_0 > 0$ on a possibly smaller relatively compact $W_+ \Subset U_+$ and s near β . Regression certificate (companion release [14]): `data/anchor/cas_orbit_measure_IFT_A001.json`. $\square$

8.3 Independent backward-wall geometry (not used in Theorem F)

The following calculation is retained as a geometric comparison with the forward wall. The deficiency proof below does not use it; the $+i$ space is obtained from the forward $-i$ space by the Lean-backed sign involution.

Proposition 2 (Backward wall). *There exist a nonempty open U_- and $\alpha \in C^\infty(U_-)$ such that for $(a, c) \in U_-$ a selected real branch exists for $s > \alpha(a, c)$ with $\|q\| \rightarrow \infty$ as $s \downarrow \alpha^+$.*

Proof. At $a = \frac{1}{54}$, $c = 2$,

$$A(s; \frac{1}{54}, 2) = -\frac{(2s-1)(3s^2-1)}{3},$$

so $A(\frac{1}{2}) = 0$, $A_s(\frac{1}{2}) = \frac{1}{6} > 0$, $B(\frac{1}{2}) = -1 \neq 0$. By the IFT there is a neighborhood of $(\frac{1}{54}, 2)$ and $\alpha \in C^\infty$ with $\alpha(\frac{1}{54}, 2) = \frac{1}{2}$ and $A(\alpha(a, c); a, c) = 0$.

Set $\tau = \sqrt{s - \alpha}$ (so $s = \alpha + \tau^2$) and $q_0 = r/\tau$. Define

$$G_-(a, c, \tau, r) := \frac{A(\alpha + \tau^2; a, c)}{\tau^2} r^3 + B(\alpha + \tau^2; c) r + C(c) \tau \quad (\tau \neq 0),$$

and extend at $\tau = 0$ by

$$G_-(a, c, 0, r) = A_s(\alpha; a, c) r^3 + B(\alpha; c) r.$$

Smoothness of the extension follows from $A(\alpha) = 0$ and $A(\alpha + \tau^2) = A_s(\alpha)\tau^2 + O(\tau^4)$. At the base, $G_-(\frac{1}{54}, 2, 0, r) = \frac{1}{6}r^3 - r$ has simple roots $r = \pm\sqrt{6}$ because $\partial_r G_- = \frac{1}{2}r^2 - 1$ and $\partial_r G_-(\pm\sqrt{6}) = -2B = 2 \neq 0$. The IFT continues $r_\pm(a, c, \tau)$ on an open set $U_- \times (0, \tau_0)$, with $r_\pm \neq 0$ after shrinking. Then $|q_0| = |r_\pm|/\tau \rightarrow \infty$ as $\tau \downarrow 0$.

For reconstruction, the denominator in (8.0) behaves as $q_0(Bq_0 + 3c) \sim Bq_0^2$ with $B \rightarrow -1 \neq 0$, hence is nonzero for small τ after shrinking U_- . Combined with $q_0 \sim r/\tau$ one obtains

$$\|q\| \geq |q_0| \geq \frac{c'_0}{\sqrt{s - \alpha}}$$

on a relatively compact $W_- \Subset U_-$ near the wall. Regression certificate (companion release [14]): `data/anchor/cas_backward_incomplete_wall_A001.json`. $\square$

8.4 Saturated maximal sheets

Lemma 1 (Saturated maximal sheets). *Let $W_+ \Subset U_+$ be nonempty, open, and relatively compact. On the selected large- q_0 forward branch, pick $\varepsilon_0 > 0$ small and the smooth cross-section*

$$\Sigma_+ = \{q : F(q) = (a, \beta(a, c) - \varepsilon_0, c), (a, c) \in W_+\}.$$

Let ϕ_t be the maximal flow of X_1 . Choose $\varepsilon_0 > 0$ small enough that $\varepsilon_0 < \tau_0^2$ for the IFT window of the forward-wall proposition, so that on W_+ the selected large- q_0 branch is a single smooth immersed sheet with nonvanishing reconstruction denominator throughout the open interval $(\beta - \varepsilon_0, \beta)$. Because $X_1 F_1 = 1$, $F_1(\phi_t(q)) = F_1(q) + t$ on the existence interval. Define

$$D_+ = \{(a, c, s) : (a, c) \in W_+, \ell_+(a, c) < s < \beta(a, c)\},$$

where β is the wall value (hence the maximal upper s -time of this sheet) and ℓ_+ is the maximal lower s -time of the immersed branch through Σ_+ . Then

$$\Psi_+ : D_+ \rightarrow \Omega_+ \subset \mathbb{R}^3, \quad \Psi_+(a, c, s) = \phi_{s - (\beta - \varepsilon_0)}(q_\Sigma(a, c))$$

is a C^∞ diffeomorphism onto an open X_1 -invariant set, $F \circ \Psi_+ = (a, s, c)$, and $s \rightarrow \beta^-$ occurs at spatial infinity. If ℓ_+ is finite, maximality forces escape as $s \downarrow \ell_+$ as well.

Proof. The maximal-flow domain $\mathcal{D} \subset \mathbb{R} \times \mathbb{R}^3$ of the smooth vector field X_1 is open and the flow ϕ is C^∞ on $\mathcal{D}$ [6]. The set D_+ is (up to the smooth reparameterization $s = F_1$) the pullback of $\mathcal{D}$ through the cross-section Σ_+ ; openness of $\mathcal{D}$ therefore yields openness of D_+ . Along every orbit, F_0 and F_2 are constant and F_1 is the time coordinate. Hence two points of the parameter domain with

different (a, c) cannot map to the same orbit, while ODE uniqueness handles two parameterizations on a fixed orbit. Equivalently, the identity $F \circ \Psi_+ = (a, s, c)$ immediately gives injectivity of Ψ_+ . Since $\det DF \neq 0$, Ψ_+ is a local diffeomorphism, hence a C^∞ diffeomorphism onto its open image Ω_+ . By the ε_0 -window choice, the selected branch does not degenerate before the wall, so the maximal upper time equals β . Finite lower maximal time forces escape from compacts by standard ODE maximality.

No continuity or semicontinuity of the variable endpoint $\ell_+(a, c)$ is needed below. In particular, the weak-adjoint argument is carried out directly on the open variable domain D_+ , rather than through a global assertion about the finite boundary of Ω_+ . $\square$

Lean coverage for the maximal sheet. The diffeomorphism and X_1 -invariance formulation in this lemma is the paper-level maximal-flow packaging. The pinned Lean route formalizes the ingredients consumed by the deficiency proof, including `injOn_PsiFin3`, `isOpen_image_Psi`, `measurableEmbedding_PsiFin3`, the exact coordinate identity, and `abs_det_fderiv_PsiFin3`; it does not package a named global diffeomorphism or X_1 -invariance theorem.

8.5 Deficiency vectors in $\text{Dom}(H^*)$

Lemma 2 (Deficiency vectors). *For $\chi \in C_c^\infty(W_+)$, so in particular $\text{supp } \chi \Subset W_+$, define on the entire maximal interval*

$$u_-(\Psi_+(a, c, s)) := \chi(a, c) e^{s-\beta(a, c)},$$

and $u_- = 0$ off Ω_+ . No interior s -cutoff. Then $u_- \in L^2(\mathbb{R}^3)$, $u_- \in \text{Dom}(H^)$, and $(H^* + i)u_- = 0$. The linear map $\chi \mapsto u_-$ is injective, so $n_- = \infty$.*

Proof. Square-integrability. By (7.1),

$$\|u_-\|_2^2 = \frac{1}{2} \int_{W_+} |\chi|^2 \left(\int_{\ell_+}^{\beta} e^{2(s-\beta)} ds \right) da dc \leq \frac{1}{4} \|\chi\|_{L^2(W_+)}^2,$$

since the inner integral equals $\frac{1}{2}(1 - e^{2(\ell_+-\beta)}) \leq \frac{1}{2}$.

Adjoint equation. Let $\varphi \in C_c^\infty(\mathbb{R}^3)$ and put $K = \text{supp } \chi \Subset W_+$. On Ω_+ , $H = -i\partial_s$ in the flow-box chart with density $\frac{1}{2}$, and $\text{div } X_1 = 0$. We verify the weak identity for the zero extension directly. This avoids any inference of regularity across $\partial\Omega_+$, and it uses no continuity property of ℓ_+ .

Both terms in the weak-adjoint identity are absolutely integrable. Indeed, $X_1\varphi$ is bounded because φ has compact support (the polynomial growth of X_1 is irrelevant on that compact set), and on D_+

$$\begin{aligned} & \left| \chi(a, c) e^{s-\beta(a, c)} \varphi(\Psi_+(a, c, s)) \right| \\ & + \left| \chi(a, c) e^{s-\beta(a, c)} (X_1\varphi)(\Psi_+(a, c, s)) \right| \leq (\|\varphi\|_\infty + \|X_1\varphi\|_\infty) |\chi(a, c)| e^{s-\beta(a, c)}. \end{aligned}$$

The right side is integrable because $\int_{\ell_+(a, c)}^{\beta(a, c)} e^{s-\beta(a, c)} ds \leq 1$ and χ has compact transverse support.

Change variables on Ω_+ and apply Fubini; the preceding majorant proves absolute integrability of both pairings. Now fix (a, c) . On every compact interval $[r, t] \Subset (\ell_+(a, c), \beta(a, c))$, ordinary one-dimensional integration by parts applies. Let $t \uparrow \beta(a, c)$ and $r \downarrow \ell_+(a, c)$ separately. These are limits on one fixed fiber: the two improper integrals converge absolutely by the same exponential estimate. At the upper endpoint the sheet escapes every compact set, so φ vanishes eventually. At the lower endpoint, either ℓ_+ is finite and maximality again forces spatial escape, or $\ell_+ = -\infty$ and

$e^{s-\beta} \rightarrow 0$. Hence both endpoint terms vanish, and the desired one-dimensional weak identity holds on that fiber.

We then integrate this already established fiberwise equality over (a, c) . Fubini applies to both sides by the global majorant above; there is no transverse limiting process, measurable exhaustion, or endpoint-selection argument. Consequently

$$\langle u_-, H\varphi \rangle = \langle -iu_-, \varphi \rangle,$$

i.e. $u_- \in \text{Dom}(H^*)$ and $(H^* + i)u_- = 0$.

Injectivity with a uniform bound. The collar used to define Σ_+ lies in every selected maximal interval, so $\ell_+(a, c) < \beta(a, c) - \varepsilon_0$. Therefore

$$\|u_-\|_2^2 \geq \frac{1}{4}(1 - e^{-2\varepsilon_0})\|\chi\|_{L^2(W_+)}^2.$$

Thus $\chi \mapsto u_-$ is bounded below, hence injective, and embeds the infinite-dimensional space $C_c^\infty(W_+)$ into $\ker(H^* + i)$. $\square$

Lean coverage for the cutoff family. The displayed bounded-below map and its uniform quantitative constant are paper-level. The pinned Lean proof reaches the same required non-finite-dimensional conclusion through `exists_finite_weakAdjointEigenvector_negI_family`: for every finite cardinality it constructs linearly independent deficiency vectors from disjoint compactly supported transverse cutoffs.

8.6 Sign involution and the opposite deficiency space

Lemma 3 (Sign transport). *Let $\sigma(q_0, q_1, q_2) = (-q_0, -q_1, q_2)$ and $(U_\sigma f)(q) = f(\sigma(q))$. Then U_σ is a complex-linear unitary involution of $L^2(\mathbb{R}^3)$, preserves $C_c^\infty(\mathbb{R}^3)$, and*

$$HU_\sigma = -U_\sigma H \quad \text{on } C_c^\infty(\mathbb{R}^3).$$

*Consequently $H^*U_\sigma = -U_\sigma H^*$, and U_σ restricts to a complex-linear isometric bijection*

$$\ker(H^* + i) \xrightarrow{\cong} \ker(H^* - i).$$

Proof. The linear involution σ preserves Lebesgue measure, so its pullback is unitary and preserves the test-function core. Direct substitution in the explicit polynomial field gives

$$X_1(f \circ \sigma) = -(X_1 f) \circ \sigma,$$

which is the displayed core anticommutation for $H = -iX_1$. For $v \in \text{Dom}(H^*)$ and a core test function, unitarity and core anticommutation transfer the defining adjoint pairing and give $U_\sigma v \in \text{Dom}(H^*)$ with $H^*U_\sigma v = -U_\sigma H^*v$. Since $U_\sigma^2 = I$, this transports the $-i$ eigenspace isometrically and bijectively to the $+i$ eigenspace. This is the route formalized for the specific canonical minimal operator `H_X1_min` in the companion Lean artifact. $\square$

Remark 3 (Withdrawn device). *Interior indicators $\mathbf{1}_{(\beta-\delta, \beta)}(s)$ create δ -masses and exit $\text{Dom}(H^*)$. That construction is withdrawn.*

Theorem 5 (F). *The Hilbert deficiency indices satisfy*

$$n_+ = n_- = \aleph_0,$$

equivalently $(n_+, n_-) = (\infty, \infty)$ in conventional notation.

Proof. The forward-wall deficiency lemma makes $\ker(H^* + i)$ infinite-dimensional. Sign transport gives an isometric bijection with $\ker(H^* - i)$, so the opposite space is infinite-dimensional as well. Both are closed subspaces of the separable Hilbert space $L^2(\mathbb{R}^3)$; their Hilbert dimensions (orthonormal-basis cardinals) are therefore exactly $\aleph_0$. This is a Hilbert-dimension statement, not an assertion that the algebraic Hamel rank is $\aleph_0$. $\square$

Theorem 6 (E). $H = -iX_1$ is not essentially self-adjoint on $C_c^\infty(\mathbb{R}^3)$.

Proof. $n_\pm \geq 1$ by Theorem F. $\square$

Corollary 1. Let $\overline{H}$ denote the closure of H . Von Neumann’s theorem applies to the closed symmetric operator $\overline{H}$. Since $H^* = (\overline{H})^*$, its deficiency spaces are those computed above; moreover, a self-adjoint operator extends H if and only if it extends $\overline{H}$. Fixing one deficiency-space isometry and multiplying it by the unit phases $\lambda \in \mathbb{C}$, $|\lambda| = 1$, gives distinct extensions under the von Neumann bijection. Lean proves the injectivity of this phase parameterization; the classical fact that the complex unit circle has cardinality continuum then gives a continuum-sized family of self-adjoint extensions. Every such extension agrees with the same minimal operator on $C_c^\infty(\mathbb{R}^3)$; the core algebraic Poisson/CCR relations recorded in Theorems B–C therefore do not, by themselves, distinguish a unique extension.

9 Discussion

The example separates algebraic and analytic layers. The Poisson cotangent lift and dual-field/CCR identities are clean, and ψ is a well-defined polynomial Weyl endomorphism. Nonetheless X_1 is incomplete and $H = -iX_1$ has Hilbert deficiency indices $(\aleph_0, \aleph_0)$. Algebraic CCR/Poisson data therefore do not select a unique self-adjoint dual momentum for this generator.

We do *not* claim here exact deficiency indices for H_0 or H_2 , nor a theorem classifying all strong CCR or dual-translation packages; those topics appear only in companion repository notes and may be incomplete relative to theorem grade.

The collision is used in Theorem A and in the noninjectivity clause of Theorem B, but it is not an input to Theorems D–F. Those analytic results use the constant nonzero Jacobian (to define the dual field and its invariant measure) and the explicit incompleteness/escape geometry of X_1 . Thus the logical claim is not “counterexample implies quantization ambiguity.” Classical incomplete flows, half-line deficiency models, and von Neumann extension theory are standard [1]. The contribution is their explicit, Lean-backed realization for this dual transport operator. We make no new general theorem equating surjectivity of a real Keller map with completeness of all its dual fields.

10 Non-claims

No unitary gate, channel, CP map, or computational advantage. No preferred physical extension. No claim that all quantization routes fail. Discovery of F is not claimed. No rank-three Dixmier counterexample is claimed. Lean covers the bounded canonical-core Theorems E–F, the exact Hilbert-basis cardinal result, the specific `H_X1_min` von Neumann classification, and the injective unit-phase family of distinct extension witnesses at the synchronized freeze identified above. The algebraic index `hamelDeficiencyRank`, defined using `Module.rank`, has lower bounds only. The independent backward-wall construction, exact cardinality of the full extension type, arbitrary-operator theorem, strong CCR, preferred extension, and algebraic or physical selection remain

outside the Lean corollary. No theorem-level dual-flow strong CCR obstruction is claimed in this note. In particular, the artifact constructs the displayed Weyl endomorphism ψ but does not establish that ψ is noninvertible.

References

- [1] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II: Fourier Analysis, Self-Adjointness*, Academic Press, 1975.
- [2] J. Dixmier, *Sur les algèbres de Weyl*, Bull. Soc. Math. France **96** (1968), 209–242.
- [3] Y. Tsuchimoto, *Endomorphisms of Weyl algebra and p -curvatures*, Osaka J. Math. **42** (2005), 435–452, <https://doi.org/10.18910/7472>.
- [4] A. Belov–Kanel and M. Kontsevich, *The Jacobian conjecture is stably equivalent to the Dixmier conjecture*, Moscow Math. J. **7** (2007), 209–218, <https://arxiv.org/abs/math/0512171>.
- [5] P. K. Adjagbo and A. van den Essen, *A proof of the equivalence of the Dixmier, Jacobian and Poisson conjectures*, Acta Math. Vietnam. **32** (2007), 205–214, <https://math.ac.vn/uploads/files/0702205.pdf>.
- [6] J. M. Lee, *Introduction to Smooth Manifolds*, 2nd ed., Springer, 2013.
- [7] H. Bass, E. H. Connell, and D. Wright, The Jacobian conjecture: reduction of degree and formal expansion of the inverse, Bull. Amer. Math. Soc. **7** (1982), 287–330.
- [8] L. Alpöge, informal public announcement of a Keller map with constant Jacobian and three-point collision (credits Akhil Mathew for prompting the question and Claude Fable for work leading to the map), July 2026, https://x.com/_alpoge_/status/2079028340955197566.
- [9] Ulam, *A Counterexample to the Jacobian Conjecture*, preprint, July 2026, <https://www.ulam.ai/research/jacobian.pdf>.
- [10] A. Freitas Ramos, D. Barros Hulak, and R. J. G. Barretto de Queiroz, *Formal Verification of an Explicit Counterexample to the Jacobian Conjecture*, Archive of Formal Proofs, 20 July 2026, https://isa-afp.org/entries/Jacobian_Counterexample.html.
- [11] T. Shaska, *Graded Keller maps and the Jacobian Conjecture*, arXiv:2607.20210, 2026, <https://arxiv.org/abs/2607.20210>.
- [12] Z. Jelonek and M. Lason, *Quantitative properties of the non-properness set of a polynomial map*, Manuscripta Math. **156** (2018), 383–397, <https://doi.org/10.1007/s00229-017-0965-0>.
- [13] D. Speyer, *The new counterexample to the Jacobian conjecture*, Secret Blogging Seminar, 20 July 2026, <https://sbseminar.wordpress.com/2026/07/20/the-new-counterexample-to-the-jacobian-conjecture/>.
- [14] D. E. Fredriksen, EXOTIC-CCR A001 companion code and CAS certificates, GitHub Quantyra/jacobian-weyl-quantum-phase-space, paper-package release target v0.3.9-referee-revision; concept DOI <https://doi.org/10.5281/zenodo.21474351> (project-level).
- [15] D. E. Fredriksen, exotic-ccr-lean, GitHub Quantyra/exotic-ccr-lean, released tag v0.1.8-theorem-e, tag target be4f330; reviewed bounded Theorem E source anchor a6bb091, <https://github.com/Quantyra/exotic-ccr-lean/releases/tag/v0.1.8-theorem-e> (Lean 4 formalization of the seed identities, dual-matrix algebra, Piola row divergence, full dual-field commutation, the polynomial Weyl endomorphism ψ , and incompleteness of the dual field X_1).
- [16] Theorem F supplement: synchronized, untagged, unreleased Lean publication freeze 2e40c4cab86a1ef97cb3334497d10081dfe33867 (historical theorem-source root ff50f4a2a312591c2e5b26e71eb390ade9164b34); `ExoticCCR.theoremF`, `aleph0_le_hamelDeficiencyRank_X1`, and `hilbertDeficiencyIndex_X1_eq_aleph0`, `theoremFVonNeumannClassification`, `theoremFUnitPhaseExtension`, and `theoremFUnitPhaseExtension_injective`. Both standard adjoint eigenspaces are not finite-dimensional, and Cardinal.aleph0 is a lower bound for each `hamelDeficiencyRank`. The separate Hilbert-basis cardinal is exactly aleph0 for both closed eigenspaces; `hamelDeficiencyRank` is the algebraic (Hamel) rank and its exact value is not claimed. For the specific canonical minimal operator, all self-adjoint-extension witnesses are classified bijectively by complex-linear isometric equivalences from the $+i$ to the $-i$ adjoint eigenspace, and unitary complex phases inject into distinct witnesses. Together with the classical cardinality of the unit circle, this yields a continuum-sized lower family; the cardinal identification is not asserted as a Lean declaration. No exact cardinality of the full extension type, preferred extension, arbitrary-operator theorem, backward-wall construction, or algebraic/physical selection theorem is claimed.