

Completely positive readings of an exotic Weyl endomorphism: algebraic correspondence, and obstructions for standard completions

Daniel Eric Fredriksen
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

Let ψ be the polynomial Weyl-algebra endomorphism induced by the Alpöge–Fable Keller seed (restated) via $Q = F(q)$ and $P = J^{-T}p$. We prove ψ is a unital $*$ -endomorphism (algebraic SOS positivity), construct its algebraic endomorphism correspondence, and obstruct two standard continuous packages: unital $*$ -endomorphisms of $C_0(\mathbb{R}^3)$ extending composition with F , and Bogoliubov endomorphisms of the CCR C^* -algebra. We also obstruct unital CP maps that are C_0 -composition compatible. A free (non- C_0 , non-Bogoliubov) unital CP extension to some C^* completion remains open. No dual F -translations, gates, or advantage are claimed. Companion to the A001 deficiency note [1].

MSC: 46L07, 81S05, 47L90, 46L08.

Version: C001 obstruction freeze v0.1.

T4: does not implement dual F -translations on $L^2(\mathbb{R}^3)$.

1 Domain

Definition 1. $\mathcal{W}$: polynomial Weyl algebra over $\mathbb{C}$, $[q_i, p_j] = \delta_{ij}$, involution $q_i^* = q_i$, $p_j^* = p_j$. With $B = J^{-T}$,

$$\psi(q_i) = F_i(q), \quad \psi(p_j) = \sum_k B_{jk}(q) p_k$$

extends to a unital algebra endomorphism $\psi : \mathcal{W} \rightarrow \mathcal{W}$.

Theorem 1 (Algebraic $*$ -SOS). *If F, B are real, then ψ is a unital $*$ -endomorphism, hence preserves hermitian-square cones at all matrix levels $\text{id}_n \otimes \psi$.*

2 Non-properness

Theorem 2. $F : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ is not proper: along the explicit incomplete X_1 -orbit $\gamma(t)$ of [1], $F(\gamma(t)) = (0, t, 2)$ stays bounded while $\|\gamma(t)\| \rightarrow \infty$ (CAS `cas_F_not_proper_C1c_A001.json`).

3 Obstructions

Theorem 3 (C0 *-endo). *No unital *-endomorphism of $C_0(\mathbb{R}^3)$ extends $f \mapsto f \circ F$ on polynomials.*

Theorem 4 (Bogoliubov). *No Bogoliubov endomorphism of the CCR C^* -algebra restricts to ψ on $\mathcal{W}$ (F is nonlinear).*

Theorem 5 (C0-compatible CP). *No unital CP map on a C^* algebra containing $C_0(\mathbb{R}^3)$ as position functions can extend ψ while agreeing with composition on a dense subalgebra of that C_0 .*

4 Construct: algebraic correspondence

Definition 2 (Endomorphism correspondence). *Let $E = \mathcal{W}$ as a right $\mathcal{W}$ -module, left action $a \cdot \xi = \psi(a)\xi$, and $\langle \xi, \eta \rangle = \xi^* \eta$.*

This is the standard algebraic Hilbert-module correspondence implementing the unital *-endo ψ . It does *not* by itself yield a continuous CP map on a C^* completion.

Theorem 6 (CCR *-hom extension). *No unital *-endomorphism of the CCR C^* -algebra generated by Weyl unitaries restricts to ψ on $\mathcal{W}$: such endomorphisms are Bogoliubov (linear on one-particle fields), while F is nonlinear.*

5 Open

Strict free unital CP (some abstract C^* envelope of $\mathcal{W}$, with neither C0-composition nor CCR/*-hom/Bogoliubov requirements): not constructed, not ruled out. See companion GenCP-Free progress note.

6 Non-claims

No channel on $B(H)$. No gates/advantage. No dual F -translations. No claim every CP extension is impossible in every axiomatics. Algebraic correspondence $\neq$ physical instrument.

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., v0.3.6-submit, <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.