

# Dual-lift realizability for Keller-type maps: structural incompleteness, schema, and examples

Daniel Eric Fredriksen  
Quantlyra Inc.

<https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space>

22 July 2026

**Current status.** This B001 PDF is an untagged, unreleased cross-companion correction. The historical `v0.2.3-b001-draft` tag does not identify this binary. A001 proposed candidate `v0.3.9-referee-revision` is likewise untagged/unreleased and has no version DOI; its current non-claims and package/metadata re-gates are pending. If those pass, Dan approval and math-ph endorsement/publication actions still remain. The project concept DOI resolves a historical record and does not identify the current A001 candidate.

## Abstract

We isolate a structural lemma for dual vector fields of local diffeomorphisms  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$ : if global dual fields  $X_j$  with  $DF \cdot X_j = e_j$  are all complete, then  $F$  is surjective. Hence any non-surjective dual system has at least one incomplete dual field. We propose a classification schema (B-incomplete / B-ESS / B-Def / B-CCR / B-CP) and populate it on atlas rows A000–A010 (pass and fail baselines, Keller seed A001, family pilot A002, a smooth 2D pair A007/A008 illustrating that the forced incomplete set can be as thin as measure zero, and a polynomial 2D pair A009/A010: Pinchuk’s real-plane map with *open-set* incompleteness, and a polynomial shear automorphism with complete dual fields). We prove restricted dichotomy lemmas: polynomial product local diffeos have complete dual fields; the A007 zero-dimensional asymptotic mechanism is impossible for polynomials (Jelonek); in dimension 2 dual incompleteness is fiber-saturated and falls into empty / open / residual-thin regimes. Residual-thin regime T is excluded on proper, injective, degree-one-component, product, triangular, and graph-type (including all  $\deg P \leq 2$ ) classes, and reduced to incompleteness only on atypical fibers of non-coordinate  $P$  of degree  $\geq 3$ ; full exclusion of T remains open. No gates, channels, or computational advantage are claimed. Companion pack: <https://github.com/Quantlyra/jacobian-weyl-quantum-phase-space/blob/main/docs/notes/COMPANION-PACK.md>.

**MSC:** 37C10, 47B25, 53D17, 14R15.

**Version:** B001 draft v0.7 (erratum: real poly local-diffeo language — global  $C^\infty$  diffeo / regime E; polynomial inverse only when proved; supersedes freeze 61bf3da automorphism wording).

**Software:** atlas `program_b`; CAS anchors:

`cas_atlas_A005_A006_B001.json`, `cas_atlas_A007_A008_B001.json`,  
`cas_atlas_A009_A010_B001.json`, `cas_atlas_B5_poly_dichotomy_B001.json`,  
`cas_atlas_B6_regime_T_B001.json`, `cas_atlas_B7_OPEN_T_B001.json`.

## 1 Introduction

Cotangent lifts and dual fields are standard when  $\det DF \neq 0$ . After the A001 deficiency analysis [1], we classify which maps force incomplete dual fields and how that interacts with ESS / CCR / CP

packages [2]. A001 proposed candidate **v0.3.9-referee-revision** is untagged/unreleased, has no version DOI, and still awaits current non-claims and package/metadata re-gates. If those pass, Dan approval and math-ph endorsement/publication actions remain. The project concept DOI resolves a historical version and does not identify the current A001 candidate.

## 2 Structural lemma

**Lemma 1** (Complete dual fields imply surjectivity). *Let  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  be  $C^\infty$  and a local diffeomorphism with dual fields  $X_j$ ,  $DF \cdot X_j = e_j$ . If every  $X_j$  is complete, then  $F(\mathbb{R}^n) = \mathbb{R}^n$ .*

*Proof.*  $F(\phi_t^{(j)}(q)) = F(q) + te_j$ ; iterate; image open and translation-invariant.  $\square$

**Corollary 1.** *Non-surjective dual systems have some incomplete  $X_j$ .*

## 3 Schema

B-incomplete / B-ESS / B-Def / B-CCR / B-CP with values **open** / **fail** / **pass**.

## 4 Atlas scoreboard (v0.4)

| ID   | Map (sketch)               | B-inc. | Role                                      |
|------|----------------------------|--------|-------------------------------------------|
| A000 | id                         | pass   | baseline                                  |
| A004 | shear $(x, y + x, z)$      | pass   | linear auto                               |
| A005 | diag $(2, \frac{1}{2}, 1)$ | pass   | linear stretch (CAS)                      |
| A008 | $(x, y + \sin x, z)$       | pass   | 2D smooth global diffeo (CAS)             |
| A010 | $(x, y + x^2, z)$          | pass   | 2D <i>polynomial</i> auto (CAS)           |
| A003 | $(x, e^y)$ pad             | fail   | smooth B1 example                         |
| A006 | $(e^x, e^y, z)$            | fail   | two incomplete fields, open set (CAS)     |
| A007 | $z \mapsto e^z$ pad        | fail   | incomplete on <i>null set only</i> (CAS)  |
| A009 | Pinchuk $(P, Q)$ pad       | fail   | poly, incomplete on <i>open set</i> (CAS) |
| A001 | Keller seed                | fail   | $X_1 + H_1$ analytic [1]                  |
| A002 | family d4 pilot            | open   | geometry pending                          |

## 5 Examples (new in v0.2)

**Example 1** (A005 — linear stretch, pass-like).  $F = \text{diag}(2, \frac{1}{2}, 1)$ ,  $\det DF = 1$ . Dual fields  $X_0 = \frac{1}{2}\partial_x$ ,  $X_1 = 2\partial_y$ ,  $X_2 = \partial_z$  are complete. CAS: `cas_atlas_A005_A006_B001.json`.

**Example 2** (A006 — exp quadrant, fail-like).  $F(x, y, z) = (e^x, e^y, z)$ ,  $\det DF = e^{x+y} > 0$ , image  $(0, \infty)^2 \times \mathbb{R}$ . Dual fields  $X_0 = e^{-x}\partial_x$ ,  $X_1 = e^{-y}\partial_y$  incomplete backward ( $e^x = e^{x_0} + t$ ). Matches Lemma 1. CAS same file.

## 6 Examples (new in v0.3): two incompleteness regimes

**Example 3** (A007 — complex exponential, thin fail-like).  $F(x, y, z) = (e^x \cos y, e^x \sin y, z)$ , i.e.  $z \mapsto e^z$  padded. Conformal,  $\det DF = e^{2x} > 0$ ; image misses the origin (non-surjective). Writing  $w = e^{x+iy}$ , the dual field  $X_0$  solves  $\dot{w} = 1$ , so  $w(t) = w(0) + t$ . If  $\text{Im } w(0) \neq 0$  (full measure), the

orbit never reaches  $w = 0$ : **complete**. If  $\text{Im } w(0) = 0$  (a measure-zero pair of rays),  $w(t)$  hits 0 at finite  $t$  and  $x(t) \rightarrow -\infty$ : **incomplete**. Thus Lemma 1's corollary is satisfied by an incomplete set of measure zero, sharply contrasting A006/A001 where incompleteness occupies an open set. CAS: `cas_atlas_A007_A008_B001.json`.

**Example 4** (A008 — shear + sine, 2D global diffeomorphism).  $F(x, y, z) = (x, y + \sin x, z)$ ,  $\det DF = 1$ . Explicit global inverse  $(x, y) \mapsto (x, y - \sin x)$  (verified by CAS roundtrip). Dual fields  $X_0 = \partial_x - \cos x \partial_y$ ,  $X_1 = \partial_y$ ; the  $X_0$  flow  $y(t) = y_0 - \sin(x_0 + t) + \sin(x_0)$  is bounded for all  $t$ : **complete**. A genuinely nonlinear, non-triangular-affine 2D coupling (unlike A004/A005) that nonetheless has no incompleteness at all.

## 7 Examples (new in v0.4): polynomial 2D evidence

**Example 5** (A009 — Pinchuk real plane map, open fail-like). Let  $t = xy - 1$ ,  $h = t(xt + 1)$ ,  $f = (xt + 1)^2(t^2 + y)$ , and

$$P = f + h, \quad Q = -t^2 - 6th(h + 1) - u(f, h),$$

with the Campbell/van den Essen auxiliary  $u = 170fh + 91h^2 + 195fh^2 + 69h^3 + 75fh^3 + \frac{75}{4}h^4$  (Pinchuk [3], Campbell [4]). Then  $F = (P, Q) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  (padded by idle  $z$ ) is polynomial of degrees (10, 25), with

$$\det DF = t^2 + (t + f(13 + 15h))^2 + f^2 > 0$$

everywhere (sum of squares;  $t = f = 0$  is impossible over  $\mathbb{R}$ ). It is not injective (CAS-certified collisions) and not surjective (misses exactly two points [4]). The dual field  $X_1 = H(P)/j = (-P_y, P_x)/j$  satisfies  $DF \cdot X_1 = e_1$  and  $\dot{Q} = 1$ . On every level  $P = c$  with  $c > -1$ , the real level curve has poles  $h = -1 \pm \sqrt{1 + c}$  at which  $(x, y) \rightarrow \infty$  while  $Q$  approaches a finite asymptotic value: escape in finite  $X_1$ -time. Hence the incompleteness locus of  $X_1$  contains the open set  $\{P > -1\}$ . **Open-set incompleteness for a genuinely polynomial 2D Keller-type map**, contrasting A007's thin failure. CAS: `cas_atlas_A009_A010_B001.json`.

**Example 6** (A010 — polynomial shear automorphism, pass-like).  $F(x, y, z) = (x, y + x^2, z)$ ,  $\det DF = 1$ , inverse  $(x, y - x^2, z)$ . Dual fields  $X_0 = \partial_x - 2x \partial_y$ ,  $X_1 = \partial_y$ ; the  $X_0$  flow  $y(t) = y_0 - 2x_0t - t^2$  is polynomial in  $t$ : **complete**. A polynomial pass baseline paralleling A008.

## 8 Toward a dichotomy (open)

A007 shows the incomplete locus forced by Lemma 1 can be as thin as measure zero for general smooth local diffeomorphisms. A007 is *not* polynomial. A009 (Pinchuk) supplies the first atlas polynomial 2D non-injective example with *open-set* incompleteness, consistent with A001's open failures in dimension 3 and with A010 as a polynomial complete baseline. The candidate claim — polynomial Keller-type maps never exhibit A007-style thin incompleteness — remains open as a *full* theorem, but is reduced by the restricted lemmas of §9.

## 9 Restricted dichotomy lemmas (new in v0.5)

Full proofs: PROGRAM-B-B5-poly-dichotomy-lemmas.md. CAS: `cas_atlas_B5_poly_dichotomy_B001.json`.

**Lemma 2** (Univariate polynomial local diffeos are global). *If  $f \in \mathbb{R}[t]$  satisfies  $f'(t) \neq 0$  for all  $t \in \mathbb{R}$ , then  $f : \mathbb{R} \rightarrow \mathbb{R}$  is a global diffeomorphism.*

*Proof.*  $f'$  has constant strict sign, so  $f$  is strictly monotone. Even degree is impossible (equal infinite limits  $\Rightarrow$  not injective). Odd degree gives opposite infinite limits, hence bijectivity; inverse is  $C^\infty$  by IFT.  $\square$

**Lemma 3** (Polynomial product maps have complete duals). *If  $F(x) = (f_1(x_1), \dots, f_n(x_n))$  with each  $f_i \in \mathbb{R}[t]$  and  $\prod_i f'_i \neq 0$ , then  $F$  is a global diffeomorphism and every dual field  $X_i = (1/f'_i)\partial_{x_i}$  is complete.*

*Proof.* Apply Lemma 2 coordinatewise; dual ODEs integrate by  $f_i(x_i(t)) = f_i(x_i(0)) + t$ .  $\square$

**Lemma 4** (Cylinder criterion). *If  $\text{int } \pi_j(\mathbb{R}^n \setminus F(\mathbb{R}^n)) \neq \emptyset$ , then the incompleteness locus of  $X_j$  contains a nonempty open set.*

*Proof.* On the open set of domain points whose  $F$ -image projects into that interior, the straight-line orbit in the target meets the complement of the image in finite time, so the dual lift is incomplete.  $\square$

**Lemma 5** (2D fiber saturation and three regimes). *Let  $F = (P, Q) : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  be polynomial with  $\det DF \neq 0$ . The incompleteness locus  $U_1$  of the dual field with  $\dot{Q} = 1$  is a union of connected components of levels of  $P$ . The set of bad level values is semi-algebraic in  $\mathbb{R}$ , hence a finite union of points and open intervals. Thus either  $U_1 = \emptyset$  (regime E; e.g. A010), or  $U_1$  contains a nonempty open set (regime O; e.g. A009), or  $U_1$  is a finite union of real algebraic curves (regime T, measure zero) — the only residual thin channel.*

*Proof sketch.* No critical points of  $P$ ; levels are unions of lines;  $Q$  is a chart on each component with speed 1; completeness iff  $Q$  hits all of  $\mathbb{R}$  on that component. Semi-algebraicity of the bad-value set is Tarski–Seidenberg.  $\square$

**Lemma 6** (Jelonek blocks the A007 asymptotic mechanism). *If  $F : \mathbb{R}^n \rightarrow \mathbb{R}^n$  is polynomial and generically finite and the asymptotic set  $S_F$  is nonempty, then  $\dim S_F = n - 1$  [5, 6]. In particular  $S_F$  cannot be a nonempty finite set of points, so the A007 pattern (single asymptotic value, thin cylindrical projection, null incomplete locus) is unavailable to polynomials.*

**Restricted package.** Lemmas 2–6 eliminate thin incompleteness for polynomial *product* maps and ban the A007 mechanism for all polynomial local diffeos. Full exclusion of regime T in Lemma 5 remains open; no atlas example realizes T.

## 10 Regime T partial exclusion (new in v0.6)

Full proofs: PROGRAM-B-B6-regime-T.md. CAS: cas\_atlas\_B6\_regime\_T\_B001.json.

**Lemma 7** (Chamber reduction for regime T). *For polynomial  $F = (P, Q)$  as in Lemma 5, Hardt chambers have constant  $Q$ -end type. Hence regime T occurs if and only if every open Hardt chamber of  $P$  is good and  $\text{Bad}_P(F)$  is a nonempty finite subset of the atypical values of  $P$ .*

**Lemma 8** (Structured exclusions of regime T). *Regime T is impossible for a polynomial local diffeomorphism  $F : \mathbb{R}^2 \rightarrow \mathbb{R}^2$  in any of the following classes:*

1.  $F$  proper (then  $F$  is a global  $C^\infty$  diffeomorphism; duals complete);

2.  $F$  injective (then  $F$  is a global  $C^\infty$  diffeomorphism; duals complete; polynomial inverse is not automatic over  $\mathbb{R}$ );
3. one component has total degree 1 (after affine domain change the map is a global  $C^\infty$  diffeomorphism);
4. product maps as in Lemma 3, or triangular maps  $F = (f(x), \alpha(x)y + \beta(x))$  with  $f', \alpha$  never zero.

*Proof sketch.* (1) Proper local diffeos  $\mathbb{R}^n \rightarrow \mathbb{R}^n$  are coverings, hence global diffeos. (2) Injective polynomial local diffeomorphisms of  $\mathbb{R}^n$  are bijective global  $C^\infty$  diffeomorphisms (IFT + surjectivity of injective real polynomial maps); dual completeness as in the proper case. Over  $\mathbb{R}$ , the inverse need not be polynomial — e.g.  $F(x, y) = (y, x + x^3)$ . (Algebraic automorphism theorems over algebraically closed fields do not supply a free poly inverse over  $\mathbb{R}$ .) (3) Reduce to  $P = x$ ; then  $Q_y \neq 0$  forces each univariate  $y \mapsto Q(u, y)$  to be a global diffeo by Lemma 2, yielding an explicit global  $C^\infty$  inverse. (4) Lemmas 3 and B5.7.  $\square$

**Lemma 9** (Model atypical submersion is not a coordinate). *Let  $P_0 = x + x^2y$ . Then  $\nabla P_0 \neq 0$  everywhere and 0 is atypical, but there is no  $Q \in \mathbb{R}[x, y]$  with  $\det D(P_0, Q)$  a nonzero constant. Moreover every  $Q$  that is linear in  $y$  has  $\det D(P_0, Q)$  vanishing somewhere; CAS finds no  $Q$  of total degree  $\leq 6$  with Jacobian equal to any of  $\pm 1, x^2 + 1, y^2 + 1, x^2 + y^2 + 1, (x^2 + 1)(y^2 + 1), (x^2 + 1)^2, (y^2 + 1)^2$ .*

**Partial T package.** Lemmas 7–9 exclude T on structured subclasses and block the standard atypical first-component model at low degree. The residual **OPEN-T** asks whether any polynomial local diffeo has  $\text{Bad}_P$  a nonempty finite subset of  $\text{Atyp}(P)$  with all Hardt chambers good. No atlas example realizes T; full dichotomy remains a conjecture.

## 11 OPEN-T sharpened: graph type and $\deg P \leq 2$ (new in v0.7)

Full proofs: PROGRAM-B-B7-OPEN-T.md. CAS: cas\_atlas\_B7\_OPEN\_T\_B001.json.

**Affine invariance.** Affine domain/codomain changes preserve  $\det DF \neq 0$ , polynomial class, dual completeness (affine conjugacy of flows), and the E/O/T regime labels; the graph-type normal form and  $\deg P \leq 2$  reduction are therefore non-vacuous inside the same package (PROGRAM-B-B7-OPEN-T.md, Remark B7.0a).

**Lemma 10** (Graph-type first component  $\Rightarrow$  global  $C^\infty$  diffeo). *Call  $P$  graph type if after an affine domain automorphism,  $P = y + f(x)$  for some  $f \in \mathbb{R}[x]$ . If  $F = (P, Q)$  is polynomial with  $\det DF \neq 0$  and  $P$  graph type, then on each fiber  $y = c - f(x)$  the restriction  $q_c(x) = Q(x, c - f(x))$  is a strictly monotone univariate polynomial (derivative  $-j \neq 0$ ), hence a  $C^\infty$  diffeomorphism  $\mathbb{R} \rightarrow \mathbb{R}$ . Solving fiberwise yields a global  $C^\infty$  inverse, so  $F$  is a global  $C^\infty$  diffeomorphism of  $\mathbb{R}^2$  (a polynomial map that is a  $C^\infty$  automorphism of  $\mathbb{R}^2$ ) and  $\text{Bad}_P = \emptyset$  (regime E). Regimes O and T are impossible. The inverse need not be polynomial in general (e.g.  $Q = x + x^3/3 + g(P)$ , or degree-one fibers  $Q = (P^2 + 1)x$  with inverse  $x = v/(u^2 + 1)$ ). Assert polynomial automorphism only when an explicit polynomial inverse is exhibited (e.g. elementary shears  $F = (y + f(x), x)$ ).*

**Theorem 1** (Degree  $\leq 2$  first component). *If  $F = (P, Q)$  is a polynomial local diffeomorphism and  $\deg P \leq 2$ , then after an affine domain change  $P$  is graph type (the Hessian of a critical-point-free quadratic is rank one), hence  $F$  is a global  $C^\infty$  diffeomorphism of  $\mathbb{R}^2$  and regime E holds. In particular regime T requires  $\deg P \geq 3$ . (Polynomial inverse only when separately proved; see Lemma 10.)*

**Lemma 11** ( $P_0$ -axis cannot carry T). *For  $P_0 = x + x^2y$ , any  $Q$  with  $\{P_0, Q\} \neq 0$  everywhere makes the atypical axis component  $\{x = 0\} \subset P_0^{-1}(0)$  good (univariate diffeomorphism in  $y$ ). Thus  $T$  cannot be realized by “only the axis is bad.” Odd vertical degree forces the leading coefficient freeze  $a_n = Cx^{2n}$ ; the linear-in- $y$  case collapses as in Lemma 9. CAS finds no  $Q$  of total degree  $\leq 8$  with Jacobian equal to standard never-zero targets (including  $(x^2 + y^2 + 1)^2$ ).*

**Sharpened OPEN-T.** Lemmas 10–11 and Theorem 1 exclude T on all graph-type /  $\deg P \leq 2$  maps and filter residual candidates to non-coordinate  $P$  of degree  $\geq 3$  with badness on bifurcation-born components. Full exclusion and existence both remain open; no atlas example realizes T.

## 12 Prior examples (v0.1–v0.2)

A000 identity; A003 half-plane; A001 Keller seed with  $X_1$  incomplete and  $H = -iX_1$  deficiency  $(\infty, \infty)$  in [1] (other  $X_j$  incompleteness is recorded in companion validation dossiers, not re-proved here); A002 open; A004 shear pass.

## 13 Non-claims

No gates, channels, advantage. No dual-flow repair. A003/A005/A006/A007/A008 are not Jacobian counterexamples (A007/A008 are not even polynomial). A009 is a classical strong-real-JC counterexample ( $\det > 0$ , not injective) but not a complex Jacobian conjecture counterexample ( $\det$  not constant; JC(2) over  $\mathbb{C}$  remains open). A010 is a polynomial shear automorphism (explicit poly inverse). T4: no dual  $F$ -translation package claimed here. The full polynomial-only dichotomy remains a conjecture (§9–§11): regime T of Lemma 5 is excluded on structured classes (Lemma 8) and on graph-type /  $\deg P \leq 2$  maps (Theorem 1, as global  $C^\infty$  diffeos / regime E); residual OPEN-T is named in §11. Prior freeze 61bf3da wording that concluded “polynomial automorphism” from bijective real poly local diffeos without a poly-inverse proof is superseded; regime-E chains are unchanged. C’s joint  $q$ – $p$  sector [2] (v0.9) splits: position CONSTRUCT (Koopman); J2 Joint-Stone-Canonical OBSTRUCTION; historical J3 full-triple Joint-Stone claim **WITHDRAWN** / **OPEN pending Lean**; J4 Joint-Stone-Hom-1 OBSTRUCTION; J5 mere-CP Joint-Stone-CP-1 OBSTRUCTION; J6 Unitary-Image-CP-1 OBSTRUCTION; J6-C Diag-CP- $\Phi_0$  CONSTRUCT; Joint-Form-Core CONSTRUCT / Joint-Form-ESS-1 OBSTRUCT; J7-C Full- $\psi$ -BT-Envelope (bounded dual-momentum transforms +  $\Phi_0$  positions; form-level only, not CFC / not Weyl- $C^*$  / not Stinespring) CONSTRUCT; J7 Full- $\psi$ -CFC-SA-1 OBSTRUCTION (via J6); Full- $\psi$ -CP-Weyl- $C^*$  OPEN (narrow).

## References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., proposed candidate v0.3.9-referee-revision (untagged/unreleased) / concept DOI 10.5281/zenodo.21474351, <https://github.com/Quantytra/jacobian-weyl-quantum-phase-space>.
- [2] D. E. Fredriksen, CP readings of an exotic Weyl endomorphism..., C001 companion v0.9 (J7 SPLIT: Full- $\psi$ -BT-Envelope form-level CONSTRUCT; CFC-SA NO-GO via J6; Weyl- $C^*$  OPEN), <https://github.com/Quantytra/jacobian-weyl-quantum-phase-space>.
- [3] S. Pinchuk, A counterexample to the strong real Jacobian conjecture, Math. Z. 217 (1994), 1–4.

- [4] L. A. Campbell, The asymptotic variety of a Pinchuk map as a polynomial curve, Appl. Math. Lett. 24 (2011), 62–65; arXiv:1001.3318.
- [5] Z. Jelonek, The set of points at which a polynomial map is not proper, Ann. Polon. Math. 58 (1993), 259–266.
- [6] Z. Jelonek, Testing sets for properness of polynomial mappings, Math. Ann. 315 (1999), 1–12.