

Completely positive readings of an exotic Weyl endomorphism: algebraic correspondence and standard-package obstructions

Daniel Eric Fredriksen
Quanytra Inc.

<https://github.com/Quanytra/jacobian-weyl-quantum-phase-space>

22 July 2026

Abstract

Let ψ be the polynomial Weyl endomorphism of the explicit Keller seed announced by Alpöge (with Fable credited in the announcement) (restated): $Q = F(q)$, $P = J^{-T}p$. With the physics normalization $[q_i, p_j] = i\delta_{ij}$ and $q_i^* = q_i$, $p_j^* = p_j$, ψ is a unital $*$ -endomorphism (algebraic SOS positivity). We construct its algebraic endomorphism correspondence and obstruct: (i) unital/nondegenerate $*$ -endomorphisms of $C_0(\mathbb{R}^3)$ extending composition with non-proper F ; (ii) Bogoliubov/quasifree endomorphisms of the CCR algebra (nonlinear F). A former “Free-Strict-Regular-v2” nondegeneracy obstruction is **withdrawn** as incorrect. We then **construct** a genuine normal unital $*$ -homomorphism (Koopman pullback) on the bounded-Borel position sector, valid for any local-diffeomorphism F regardless of properness or surjectivity. For the joint q - p sector we record: no choice-free Joint-Stone-Canonical package at the H_1 leg (A001 deficiency (∞, ∞)); the former J3 full-triple claim based on paper-only H_0, H_2 deficiency assertions is **withdrawn**, and the full-triple question remains **open pending Lean**; **obstruction** for multiplicative single-generator Joint-Stone-Hom-1 (Theorem J4); **obstruction** for mere unital-normal-CP Joint-Stone-CP-1 (Theorem J5); and **obstruction** for the weaker Unitary-Image-CP-1 pack (Theorem J6: unitary image of Schrödinger Stone unitaries without any SA-extension-of- H_1 axiom, same geometry). Bare Joint-Form-Core is constructed on C_c^∞ ; Joint-Form with ESS uniqueness fails for H_1 . Non-Stone CP of the position pullback *splits*: Diag-CP- Φ_0 ($\Phi_0 \circ E$ on the joint vNa) **construct**. Full- ψ abstract C^* without Hom/unitary-image *splits* further (Theorem J7): Full- ψ -BT-Envelope (bounded dual-momentum transforms + Φ_0 positions) **construct**—form-level / bounded-transform content only; *not* Full- ψ -CFC, *not* Weyl- C^* , *not* Stinespring; Full- ψ -CFC-SA-1 **obstruction** (SA continuous functional calculus / unitary Cayley image reduces to J6 on the same joint vNa + normal CP setting); Full- ψ -CP-Weyl- C^* remains open (narrow). No dual F -translations, gates, or advantage. Companion to [1]. Pack: COMPANION-PACK.md.

MSC: 46L07, 81S05, 47L90, 46L08.

Version: C001 companion v0.9 claim-boundary correction (untagged/unreleased; prior v0.3.3 tag is historical). J7 SPLIT: Full- ψ -BT-Envelope form-level CONSTRUCT; CFC-SA NO-GO via J6; Weyl- C^* OPEN).

T4: no dual F -translations on $L^2(\mathbb{R}^3)$.

Erratum. Free-Strict-Regular-v2 (projection off $\overline{\text{im } F}$) is **withdrawn**: $q \notin \overline{\text{im } F}$ does not imply $f(F(q)) = 0$, and $M_f \mapsto M_{f \circ F}$ on $C_b(Q)$ is a valid unital $*$ -homomorphism extendable by Arveson. Do not cite v0.2 Regular-v2 as a theorem.

1 Domain

Definition 1. $\mathcal{W}$: polynomial Weyl algebra over $\mathbb{C}$ with $[q_i, p_j] = i\delta_{ij}$, $[q_i, q_j] = [p_i, p_j] = 0$, and involution $q_i^* = q_i$, $p_j^* = p_j$. (Consistent: $[q, p]^* = -[q, p]$ and $(i\delta)^* = -i\delta$.) With real $B = J^{-T}$,

$$\psi(q_i) = F_i(q), \quad \psi(p_j) = \sum_k B_{jk}(q) p_k$$

(coefficients left of momenta) extends to a unital algebra endomorphism.

Remark 1 (Normalization). Earlier drafts used $[q, p] = \delta$ with self-adjoint generators, which is inconsistent under $*$. This note uses the standard physics normalization $[q, p] = i$.

Theorem 1 (Algebraic *-SOS). Assume F, B real and the Piola/dual-field identity $\operatorname{div} B = 0$ (row divergence free; true for the A001 cotangent lift). Then each Schrödinger operator $H_j = -iX_j$ is symmetric on $C_c^\infty(\mathbb{R}^3)$ (not claimed essentially self-adjoint—cf. [1] for H_1). At the algebraic level the formal adjoint correction $-i \sum_k \partial_k B_{jk}$ vanishes, so

$$P_j^* = P_j \quad \text{in } \mathcal{W}$$

with the physics involution. Hence ψ is a unital *-endomorphism and $\operatorname{id}_n \otimes \psi$ preserves hermitian-square cones.

Definition 2 (Algebraic correspondence). Let $E = \mathcal{W}$ as a right $\mathcal{W}$ -module by $\xi \cdot b = \xi b$, with left action $a \cdot \xi = \psi(a)\xi$ and $\langle \xi, \eta \rangle_{\mathcal{W}} = \xi^* \eta$.

2 Geometry

Theorem 2 (Non-proper). F is not proper along the incomplete X_1 -orbit of [1] (CAS `cas_F_not_proper_C1c_A001.json`).

3 Obstructions (retained)

Theorem 3 (C_0 composition endo). There is no *-homomorphism $\alpha : C_0(\mathbb{R}^3) \rightarrow C_0(\mathbb{R}^3)$ (nondegenerate/approximately unital in the usual C_0 sense) with $\alpha(f) = f \circ F$ for all $f \in C_0(\mathbb{R}^3)$, because F is not proper so $f \circ F \notin C_0$ for some $f \in C_0$.

Theorem 4 (Bogoliubov / quasifree). No Bogoliubov (linear quasifree) endomorphism of the CCR C^* -algebra—i.e. a map induced by a linear symplectic contraction on the one-particle space, acting on Weyl unitaries by $W(z) \mapsto W(Tz)$ —can reproduce the Heisenberg picture dynamics of ψ on the Weyl one-parameter groups associated with the field coordinates in a regular representation: those Bogoliubov maps act linearly on one-particle fields, while $q \mapsto F(q)$ is a nonlinear polynomial map (total degree > 1 ; A001 seed has terms up to total degree 7).

Remark 2 (Not claimed). Polynomial q, p are not themselves elements of the CCR C^* -algebra; the comparison is through regular representations / affiliated fields / Weyl groups. We do not claim every continuous unital *-endo of the CCR algebra fails—only Bogoliubov/quasifree. We do not claim $f \mapsto M_{f \circ F}$ fails as a map $C_b \rightarrow B(L^2)$. Arveson extensions of the abelian composition map need not implement the momentum part of ψ .

4 Withdrawn

Free-Strict-Regular-v2 nondegeneracy NO-GO: **withdrawn** (see erratum).

5 Construct: Koopman position sector

Theorem 5 (Koopman pullback). *Let $\mathcal{B}_b(\mathbb{R}^3)$ be the abelian von Neumann algebra of bounded Borel functions of Q . For F a local diffeomorphism ($\det DF \neq 0$ everywhere, in particular the A001 seed), the pullback*

$$\Phi_0(f) := f \circ F, \quad f \in \mathcal{B}_b(\mathbb{R}^3),$$

*is a well-defined **unital normal *-homomorphism** $\mathcal{B}_b(\mathbb{R}^3) \rightarrow \mathcal{B}_b(\mathbb{R}^3)$ (hence a normal unital CP map), for any such F —regardless of properness or surjectivity.*

Proof. F continuous $\Rightarrow f \circ F$ Borel and bounded whenever f is. Local diffeomorphism $\Rightarrow F$ is Lebesgue-nonsingular (preimages of null sets are null, by the inverse function theorem on a countable cover of charts), so Φ_0 descends to a.e.-equivalence classes. $(f \circ F)^* = \overline{f \circ F}$ and $(fg) \circ F = (f \circ F)(g \circ F)$ pointwise, hence a.e. A unital *-homomorphism of von Neumann algebras is automatically normal and completely positive. $\square$

Remark 3. *This resolves, at the position-sector level, the intuition behind the withdrawn Regular-v2: non-surjectivity of F does not obstruct the bounded-Borel Koopman pullback. The C0 obstruction (Theorem 3) is specific to the vanishing-at-infinity algebra C_0 , not to bounded Borel functions.*

6 Joint q - p extension: eight-slice status

Φ_0 only captures ψ on the position generators. The joint sector splits as follows. Full write-ups:

- Joint: PROGRAM-C-Free-Strict-Abstract-Joint.md
- J4: PROGRAM-C-residual-J4-Joint-Stone-Hom.md
- J5: PROGRAM-C-residual-mere-CP-Joint-Stone.md
- J6: PROGRAM-C-residual-CP-without-Stone.md
- J7: PROGRAM-C-residual-abstract-Cstar-full-psi.md

(i) **Position sector — CONSTRUCT.** The Koopman pullback Φ_0 is a normal unital *-homomorphism on bounded Borel functions of Q (previous section).

(ii) **Joint-Stone-Canonical — OBSTRUCTION (H1 only).** A *Joint-Stone* package asks for self-adjoint extensions $\tilde{H}_j \supset H_j = -iX_j$ together with a unital normal CP map Φ extending Φ_0 and implementing $\Phi(e^{isP_j^{\text{Sch}}}) = e^{is\tilde{H}_j}$. *Joint-Stone-Canonical* further requires the package to be uniquely fixed by the algebraic data (ψ, F, B) alone.

- **Theorem J2** (A001 H_1 only): since $(n_+, n_-) = (\infty, \infty)$ for $H_1 = -iX_1$ [1], von Neumann's theorem and the phase family yield a continuum-sized lower family of distinct SA-extension witnesses; (ψ, F, B) does not select one. Hence no Joint-Stone-Canonical package exists.

- **Historical J3—withdrawn.** Earlier versions imported paper-only G4 assertions about H_0, H_2 deficiency and used them to state a full-triple Joint-Stone no-go. Those inputs are not Lean-backed and are withdrawn as theorem evidence. H0/H2 deficiency, the existence of their self-adjoint extensions, and the full-triple Joint-Stone question remain **open pending Lean**.

(iii) **Joint-Stone-Hom-1 — OBSTRUCTION.** A *Joint-Stone-Hom-1* package asks for a self-adjoint extension $\tilde{H}_1 \supset H_1$ and a unital normal $*$ -homomorphism Φ on the vNa generated by bounded Borel functions of Q and $\{e^{isP_1^{\text{Sch}}}\}$ with $\Phi|_{\mathcal{B}_b(Q)} = \Phi_0$ and $\Phi(e^{isP_1^{\text{Sch}}}) = e^{is\tilde{H}_1}$.

Theorem 6 (J4). *No Joint-Stone-Hom-1 package exists for the A001 seed.*

Proof sketch. Multiplicativity yields dual- F_1 Heisenberg covariance $e^{-is\tilde{H}_1}(f \circ F)e^{is\tilde{H}_1} = f(F_0, F_1 + s, F_2)$. Hence $(F_1, \tilde{H}_1)$ is a regular Schrödinger couple and F_0, F_2 act on the multiplicity space only. The joint spectral measure of (F_0, F_1, F_2) must then have s -independent Lebesgue density. But $|\det DF| \equiv 2$, so that density equals $m/2$ with $m = \#F^{-1}$. For A001, m takes the values 3 and 1 on two positive-measure open slabs near $(a, c) = (0, 2)$. Contradiction. Full write-up: PROGRAM-C-residual-J4-Joint-Stone-Hom.md. $\square$

(iv) **Joint-Stone-CP-1 — OBSTRUCTION.** A *Joint-Stone-CP-1* package weakens Hom to mere unital normal CP: self-adjoint extension $\tilde{H}_1 \supset H_1$, unital normal CP map Φ on the same vNa, $\Phi|_{\mathcal{B}_b(Q)} = \Phi_0$, and $\Phi(e^{isP_1^{\text{Sch}}}) = e^{is\tilde{H}_1}$.

Theorem 7 (J5). *No Joint-Stone-CP-1 package exists for the A001 seed.*

Proof sketch. Unital CP and Kadison–Schwarz: if $U(s) = e^{isP_1^{\text{Sch}}}$ and $\Phi(U(s))$ is unitary, then $U(s)$ lies in the multiplicative domain $\mathcal{D}_\Phi$, so $\Phi(U(s)aU(s)^*) = \Phi(U(s))\Phi(a)\Phi(U(s))^*$ for all a . Taking $a = f(Q)$ recovers the same dual- F_1 Heisenberg covariance as in J4. The SvN product structure and s -dependent A001 sheet count then yield the same contradiction. Full write-up: PROGRAM-C-residual-mere-CP-Joint-Stone.md. $\square$

(v) **Unitary-Image-CP-1 — OBSTRUCTION (new); Diag-CP- Φ_0 — CONSTRUCT.** Drop the SA-extension-of- H_1 axiom entirely. Let M be the vNa generated by $\mathcal{B}_b(Q)$ and $\{U(s) = e^{isP_1^{\text{Sch}}}\}$.

- **Unitary-Image-CP-1:** unital normal CP $\Phi : M \rightarrow B(H)$ with $\Phi|_{\mathcal{B}_b(Q)} = \Phi_0$ and $\Phi(U(s))$ unitary for all s (no $\tilde{H}_1 \supset H_1$ required).
- **Diag-CP- Φ_0 :** $\Phi_{\text{diag}} = \Phi_0 \circ E$ with $E : M \rightarrow \mathcal{B}_b(Q)$ the normal conditional expectation onto positions (crossed-product / injectivity).

Theorem 8 (J6). *No Unitary-Image-CP-1 package exists for the A001 seed.*

Proof sketch. Unitary image $\Rightarrow U(s) \in \mathcal{D}_\Phi \Rightarrow$ same dual- F_1 Heisenberg covariance as J4/J5 $\Rightarrow$ same SvN / sheet-count contradiction. The dual-field SA link is inessential. $\square$

Theorem 9 (J6-C). $\Phi_{\text{diag}} = \Phi_0 \circ E$ is a unital normal CP map on M extending Φ_0 , with $\Phi_{\text{diag}}(U(s)) = 0$ whenever $s \neq 0$ (hence no Stone/SA momentum implementation). Schrödinger–Heisenberg conjugates of position observables pull back by Φ_0 automatically.

Full write-up: PROGRAM-C-residual-CP-without-Stone.md.

(vi) **Joint-Form split.**

- **Joint-Form-Core — CONSTRUCT:** sesquilinear forms of the minimal dual momenta $H_j = -iX_j$ on C_c^∞ realize $\psi(p_j)$ together with Φ_0 on positions (no SA/Stone requirement).
- **Joint-Form-ESS-1 — OBSTRUCT:** requiring ESS uniqueness for the $j = 1$ form fails by A001 Theorems E–F.

(vii) **Full- ψ abstract C^* — SPLIT (J7).** Three named packs (full write-up: `PROGRAM-C-residual-abstract-Cstar-full-psi.md`):

- **Full- ψ -BT-Envelope (bounded dual-momentum transforms + Φ_0 positions):** concrete C^* -algebra $\mathfrak{A}_\psi \subset B(H)$ generated by $\{(f \circ F)(Q) : f \in C_b(\mathbb{R}^3)\}$ and the bounded transforms $Z_j = Z(\overline{H_j})$ of the closed minimal dual momenta ($Z(A) = A(I + A^*A)^{-1/2}$). *Form-level / bounded-transform momentum content only*— **not** Full- ψ -CFC, **not** Weyl- C^* , **not** a Stinespring channel.
- **Full- ψ -CFC-SA-1:** unital normal CP extending Φ_0 that recovers SA continuous functional calculus of some $\tilde{A}_1$ for P_1^{Sch} (equivalently: unitary Cayley image of $C(P_1^{\text{Sch}})$).
- **Full- ψ -CP-Weyl- C^* :** unital CP from a C^* completion of the polynomial Weyl algebra extending algebraic ψ , without SA-CFC/unitary-image axioms.

Theorem 10 (J7-C). *Full- ψ -BT-Envelope (bounded dual-momentum transforms + Φ_0 positions) is a well-defined unital C^* -subalgebra of $B(H)$ carrying position content of ψ via $\Phi_0(C_b)$ and momentum-form content via $\{Z_j\}$. For $j = 1$, Z_1 is symmetric but not self-adjoint (A001: H_1 not ESS). No SA-extension choice, no unitary Stone image, no $*$ -hom of full CCR/Weyl- C^* is used. Not Full- ψ -CFC; not Stinespring.*

Theorem 11 (J7). *No Full- ψ -CFC-SA-1 package exists for the A001 seed.*

Proof sketch. Unitary Cayley image places $C(P_1^{\text{Sch}})$ in the multiplicative domain; continuous functional calculus recovers unitary Stone images $\Phi(e^{isP_1^{\text{Sch}}})$. The forced hypotheses match Unitary-Image-CP-1 on the same joint vNa M + normal CP setting (`PROGRAM-C-residual-CP-without-Stone.md`, J6 §2 checklist in the J7 note), which Theorem J6 obstructs. $\square$

Full- ψ -CP-Weyl- C^* remains **open** (J7-O; narrowed residual). $\text{Diag-CP-}\Phi_0$ is *not* full- ψ . Full- ψ -BT-Envelope is *not* Full- ψ -CFC and *not* Weyl- C^* .

7 Non-claims

No channel on $B(H)$ implementing the full ψ . No gates/advantage. No dual F -translations. No “all quantization fails.” No claim that every joint CP construction is impossible—only Joint-Stone-Canonical (J2), multiplicative Joint-Stone-Hom-1 (J4), mere-CP Joint-Stone-CP-1 (J5), Unitary-Image-CP-1 (J6), Full- ψ -CFC-SA-1 (J7), and Joint-Form-ESS-1. $\text{Diag-CP-}\Phi_0$ is not a momentum channel and is not full- ψ . Full- ψ -BT-Envelope is a concrete form-level C^* envelope (bounded dual-momentum transforms + Φ_0 positions), not a channel, not Full- ψ -CFC, and not Weyl- C^* . Full- ψ -CP-Weyl- C^* without SA-CFC is not claimed impossible. Historical G4 H_0/H_2 claims and J3 are withdrawn; the full-triple question is open pending Lean.

References

- [1] D. E. Fredriksen, Poisson and Weyl lifts..., proposed untagged candidate v0.3.9-referee-revision, <https://github.com/Quanyra/jacobian-weyl-quantum-phase-space>.