

Poisson and Weyl lifts of the Alpöge–Fable Keller
map
and nonunique self-adjoint realizations of a dual
transport operator

Daniel Eric Fredriksen

Quantyra Inc.

<https://github.com/Quantyra/jacobian-weyl-quantum-phase-space>

21 July 2026

Abstract

We study the three-dimensional Keller map F announced by Alpöge (with Fable credited): $\det DF \equiv -2$ and a three-point collision. Theorem A restates those finite identities (independently dual-CAS and Lean checked); novelty is Theorems B–F. We construct the cotangent lift $\Phi(q, p) = (F(q), J^{-T}p)$ and show it is a polynomial Poisson map on generators and non-injective. The matrix $B = J^{-T}$ defines dual fields $X_j = \sum_k B_{jk} \partial_{q_k}$; by the Piola identity and constant $\det DF$, $\operatorname{div} B = 0$, and the fields commute because they are dual to dF_i . The operators $H_j = -iX_j$ are symmetric on $C_c^\infty(\mathbb{R}^3)$.

We prove X_1 is incomplete by an explicit integral curve escaping as $t \rightarrow \frac{1}{2}^-$. In flow-box coordinates $(a, s, c) = F(q)$ one has $X_1 = \partial_s$ and $dq = \frac{1}{2} da ds dc$. Open transverse families with finite upper and lower F_1 -ends yield square-integrable deficiency functions, so

$$(n_+, n_-) = (\infty, \infty)$$

for $H = -iX_1$. Hence H is not essentially self-adjoint, yet von Neumann’s theorem supplies infinitely many self-adjoint extensions; the algebraic CCR data select none of them.

We claim no unitary gates, channels, computational advantage, or preferred physical extension.

MSC 2020: 47B25, 81S05, 14R15, 37C10, 53D17.

Keywords: Jacobian conjecture, Keller map, cotangent lift, Weyl algebra,

essential self-adjointness, deficiency indices.

Software: concept DOI 10.5281/zenodo.21474351; v0.3.0 DOI 10.5281/zenodo.21477350.

Erratum. Release v0.2.2 claimed $(n_+, n_-) = (0, \infty)$. An open backward-incomplete family forces $n_+ = \infty$ as well; the corrected pair is (∞, ∞) .

1 Introduction

1.1 Motivation and attribution

A *Keller map* is a polynomial endomorphism F of affine space with $\det DF$ a nonzero constant. The Jacobian conjecture predicts every Keller map is a polynomial automorphism. On 19–20 July 2026, Levent Alpöge announced an explicit map $F : \mathbb{C}^3 \rightarrow \mathbb{C}^3$ with $\det DF \equiv -2$ and a three-point collision, crediting the AI system Fable [7]; independent checks and Lean formalizations followed promptly (e.g. Cureton [8]; Speyer [9]). **This paper does not claim discovery of F .** Repository atlas label: A001.

The cotangent (Piola) lift $\Phi(q, p) = (F(q), J^{-T}p)$ and dual transport fields X_j are standard when $\det J$ is a nonzero constant. We ask whether algebraic preservation of Poisson/CCR data selects a unique self-adjoint L^2 realization of a dual transport generator on $C_c^\infty(\mathbb{R}^3)$.

1.2 Main results

- (A) Restatement: $\det DF \equiv -2$ and the three-point collision hold.
- (B) Φ is a polynomial Poisson map on generators and is non-injective.
- (C) Dual fields commute and are divergence-free; $H_j = -iX_j$ is symmetric on C_c^∞ .
- (D) X_1 is incomplete via an explicit escaping integral curve.
- (E) $H = -iX_1$ is not essentially self-adjoint.
- (F) $(n_+, n_-) = (\infty, \infty)$; infinitely many self-adjoint extensions exist, none selected by the algebraic data.

1.3 Related work

Jacobian conjecture background: Keller [4], Bass–Connell–Wright [3]. Deficiency indices: Reed–Simon [2]. Chernoff [1] concerns *sufficiency* criteria for ESS and is not used as a necessity theorem here.

2 Setup

$$\begin{aligned} F_0 &= (1 + xy)^3 z + y^2(1 + xy)(4 + 3xy), \\ F_1 &= y + 3x(1 + xy)^2 z + 3xy^2(4 + 3xy), \\ F_2 &= 2x - 3x^2 y - x^3 z. \end{aligned} \tag{2.1}$$

Write $J = DF$, $B := J^{-T}$, and

$$X_j := \sum_{k=0}^2 B_{jk}(q) \partial_{q_k}. \tag{2.2}$$

Collision data:

$$F(0, 0, -\tfrac{1}{4}) = F(1, -\tfrac{3}{2}, \tfrac{13}{2}) = F(-1, \tfrac{3}{2}, \tfrac{13}{2}) = (-\tfrac{1}{4}, 0, 0). \tag{2.4}$$

3 Theorem A — Seed identities

Theorem 1 (A). (1) $\det DF \equiv -2$. (2) The identities (4) hold in characteristic not 2. (3) Hence F is not injective.

$$F(0, 0, -\tfrac{1}{4}) = F(1, -\tfrac{3}{2}, \tfrac{13}{2}) = F(-1, \tfrac{3}{2}, \tfrac{13}{2}) = (-\tfrac{1}{4}, 0, 0). \tag{4}$$

Proof sketch. Expand $\det J$; evaluate at the three rational points. Dual CAS and Lean formalization [6].

4 Theorem B — Poisson lift

Theorem 2 (B). Let $\Phi(q, p) = (F(q), B(q)p)$. Then B is polynomial, $JB^T = I$, the generator Poisson brackets $\{Q_i, Q_j\} = \{P_i, P_j\} = 0$, $\{Q_i, P_j\} = \delta_{ij}$ hold, and Φ is non-injective via (4).

Proof sketch. Standard cotangent lift for constant $\det J$; non-injectivity lifts from collisions using $BJ^T = I$. CAS regression tests in the software artifact [5].

5 Theorem C — Dual fields and Piola

Theorem 3 (C). *Let $X_j = \sum_k B_{jk} \partial_{q_k}$. Then $X_j(F_i) = \delta_{ij}$, $[X_i, X_j] = 0$, $\operatorname{div} X_j = 0$, and $H_j := -iX_j$ is symmetric on $C_c^\infty(\mathbb{R}^3)$ with $[F_i, H_j] = i\delta_{ij}$.*

Proof. $X_j(F_i) = (JB^T)_{ij} = \delta_{ij}$. Then $[X_i, X_j](F_k) = 0$ for all k ; since dF is a coframe, $[X_i, X_j] = 0$. $\operatorname{Cof}(DF) = (\det DF)B$. Piola gives $\operatorname{Div} \operatorname{Cof}(DF) = 0$; constant $\det DF = -2$ implies $\operatorname{Div} B = 0$. Symmetry of $-iX$ when $\operatorname{div} X = 0$ is standard. $\square$

6 Theorem D — Incompleteness of X_1

Theorem 4 (D). *The dual field X_1 is incomplete on $\mathbb{R}^3$.*

Proof. For $t \in [0, \frac{1}{2})$, with continuous extension $\gamma(0) = (1, 0, 0)$,

$$\begin{aligned} q_0(t) &= \frac{-2t - \sqrt{1 - 2t} + 1}{t(2t - 1)}, \\ q_1(t) &= t, \\ q_2(t) &= t^2(2t - 3\sqrt{1 - 2t} - 1). \end{aligned} \tag{6.1}$$

On $(0, \frac{1}{2})$, $F(\gamma(t)) = (0, t, 2)$ and $\gamma' = X_1 \circ \gamma$ (CAS-verified identities). As $t = \frac{1}{2} - \varepsilon$, $q_0 = \sqrt{2/\varepsilon} + O(1) \rightarrow +\infty$. Thus the forward maximal time from $(1, 0, 0)$ is at most $\frac{1}{2} < \infty$. $\square$

7 Flow-box coordinates

Write $H := -iX_1$ and $n_\pm := \dim \ker(H^* \mp i)$ in the usual Reed–Simon convention $n_+ = \dim \ker(H^* - i)$. On local inverse sheets of F , with $(a, s, c) = F(q)$,

$$dq = \frac{1}{2} da \, ds \, dc, \quad X_1 = \partial_s, \quad H = -i\partial_s. \tag{7.1}$$

Eliminating variables yields

$$A(s; a, c) q_0^3 + B(s; c) q_0 + C(c) = 0, \tag{7.2}$$

with $A = -cs^3 + s^2 + 18acs - 27a^2c^2 - 16a$, $B = 3cs - 4$, $C = 2c$. If $A \rightarrow 0$ and $B \neq 0$, then some $|q_0| \rightarrow \infty$.

8 Theorems E–F — Deficiency indices

Proposition 1 (Forward wall). *At $(0, \frac{1}{2}, 2)$, $A = 0$ and $\partial_s A = -\frac{1}{2} \neq 0$. IFT yields open $U_+ \ni (0, 2)$ and a finite upper wall $\beta : U_+ \rightarrow \mathbb{R}$.*

Proposition 2 (Backward wall). *At $a = \frac{1}{54}$, $c = 2$, $A = -\frac{(2s-1)(3s^2-1)}{3}$, so $A(\frac{1}{2}) = 0$, $\partial_s A(\frac{1}{2}) = \frac{1}{6} > 0$, $B(\frac{1}{2}) = -1$. For $s = \frac{1}{2} + h$, $h \downarrow 0^+$, large roots $q_0 = \pm\sqrt{6/h} + O(1)$ give $\|q\| \rightarrow \infty$. IFT yields open U_- of finite lower walls α .*

Lemma 1 (Positive measure). $\text{Leb}_3(\{T_+ < \infty\}) > 0$ and $\text{Leb}_3(\{T_- > -\infty\}) > 0$.

Proof. Open flow-box slabs below β (resp. above α), pushed by local inverses of F , have positive Lebesgue measure in q -space. $\square$

Theorem 5 (F). $(n_+, n_-) = (\infty, \infty)$.

Proof. For $\chi \in C_c^\infty(W_+)$, $W_+ \Subset U_+$, set

$$u_-(a, s, c) = \chi(a, c) e^{s-\beta(a,c)} \mathbf{1}_{(\beta-\delta, \beta)}(s).$$

Then $u_- \in L^2$ and $(H^* + i)u_- = 0$. Varying χ gives $n_- = \infty$. Similarly

$$u_+(a, s, c) = \chi(a, c) e^{-(s-\alpha(a,c))} \mathbf{1}_{(\alpha, \alpha+\delta)}(s)$$

gives $n_+ = \infty$. Separability yields countably infinite dimensions [2]. $\square$

Theorem 6 (E). H is not essentially self-adjoint on $C_c^\infty(\mathbb{R}^3)$.

Proof. Immediate from $n_\pm \geq 1$ in Theorem F. $\square$

Corollary 1 (Extensions). *Since $n_+ = n_- = \infty$, infinitely many self-adjoint extensions exist. Algebraic CCR/Poisson data do not select a preferred extension.*

Erratum. The v0.2.2 claim $(0, \infty)$ is withdrawn.

9 Non-claims

No unitary gate, channel, CP map, or computational advantage. No ESS claim for H_0, H_2 . No strong CCR after extensions. No discovery claim for F . No preferred physical extension.

10 Conclusion

Starting from the Alpöge–Fable Keller map, Poisson/Weyl lifts exist, X_1 is incomplete, and $H = -iX_1$ has deficiency indices (∞, ∞) : not ESS, infinitely many self-adjoint extensions, none preferred by algebraic data.

References

- [1] P. R. Chernoff, Essential self-adjointness of powers of generators of hyperbolic equations, *J. Funct. Anal.* **12** (1973), 401–414.
- [2] M. Reed and B. Simon, *Methods of Modern Mathematical Physics II*, Academic Press, 1975.
- [3] H. Bass, E. H. Connell, and D. Wright, The Jacobian conjecture: reduction of degree and formal expansion of the inverse, *Bull. Amer. Math. Soc. (N.S.)* **7** (1982), 287–330.
- [4] O.-H. Keller, Ganze Cremona-Transformationen, *Monatsh. Math. Phys.* **47** (1939), 299–306.
- [5] D. E. Fredriksen, EXOTIC-CCR A001 software artifact, Zenodo (2026), DOI 10.5281/zenodo.21477350.
- [6] D. E. Fredriksen, exotic-ccr-lean,
texttt<https://github.com/Quantyra/exotic-ccr-lean>.
- [7] L. Alpöge, announcement that the Jacobian conjecture is false in dimension 3, public post, 19–20 July 2026 (credits Fable); see also software provenance in [5].
- [8] D. Cureton, independent Lean 4 formalization,
texttt<https://github.com/deancureton/jacobian>.
- [9] D. Speyer, The new counterexample to the Jacobian conjecture, Secret Blogging Seminar, 20 July 2026.