

A NOTE ON ERDŐS PROBLEM #856

QUANYU TANG

1. INTRODUCTION

Let $k \geq 2$ and $N \geq 1$ be integers. Following Erdős [2], we are interested in the structure of subsets $A \subset \{1, \dots, N\}$ which avoid “LCM patterns” of size k . More precisely, we say that A contains an LCM- k -tuple if there exist distinct $a_1, \dots, a_k \in A$ such that all pairwise least common multiples coincide:

$$\text{lcm}(a_i, a_j) = \text{lcm}(a_1, a_2) \quad \text{for all } 1 \leq i < j \leq k.$$

In [2, p. 124], Erdős introduced the counting function $F(k, x)$, defined as the largest integer s for which one can find

$$1 \leq a_1 < \dots < a_s \leq x$$

with the property that no k of the a_i have pairwise equal least common multiple. He originally conjectured that $F(k, x) = o(x)$ for every $k \geq 3$, but later showed that this is false for $k \geq 4$, see [2, Theorem 1]. In particular, for every $k \geq 4$ there exists $\delta_k > 0$ such that

$$F(k, x) \geq \delta_k x$$

for all sufficiently large x .

In the same paper, Erdős also considered the *harmonic* analogue of this problem. He showed in [2, Eq. (19)] that there is a constant $c_k > 0$ such that, whenever x is sufficiently large and $a_1 < \dots < a_s \leq x$ satisfy

$$\sum_{i=1}^s \frac{1}{a_i} > c_k \log x, \tag{1.1}$$

then there must exist k of the a_i which have pairwise equal least common multiple.

Motivated by this, it is natural to isolate the harmonic problem

$$f_k(N) := \max \left\{ \sum_{n \in A} \frac{1}{n} : A \subset \{1, \dots, N\} \text{ contains no LCM-}k\text{-tuple} \right\}.$$

Erdős’ argument (see [2, p. 127]) shows that

$$f_k(N) \ll_k \frac{\log N}{\log \log N}.$$

However, very little is known about lower bounds for $f_k(N)$.

The following problem appears as Problem #856 on Bloom’s Erdős Problems website [1].

Problem 1.1. *Let $k \geq 3$ and $f_k(N)$ be the maximum value of $\sum_{n \in A} \frac{1}{n}$, where A ranges over all subsets of $\{1, \dots, N\}$ which contain no subset of size k with the same pairwise least common multiple.*

Estimate $f_k(N)$.

It is immediate from the definition that we have the monotonicity

$$k \geq 3 \implies f_k(N) \geq f_3(N).$$

In the comments part of [1], Bloom observed that $f_3(N) \gg \log \log N$. A closer inspection of the same construction yields the stronger bound

$$f_3(N) \gg_t (\log \log N)^t \quad \text{for every integer } t \geq 1.$$

Since the detailed proofs in the comment section are rather long, we have included complete versions of these arguments in the Appendix of this note.

In this note we go significantly further and prove that

$$f_3(N) \gg (\log N)^{1/e-o(1)}.$$

2. MAIN RESULT

We recall that $f_3(N)$ denotes the maximum of

$$\sum_{n \in A} \frac{1}{n}$$

over all subsets $A \subset \{1, \dots, N\}$ which contain no triple $\{a, b, c\}$ of distinct elements with

$$\text{lcm}(a, b) = \text{lcm}(a, c) = \text{lcm}(b, c).$$

Theorem 2.1. *For every constant $0 < c < 1/e$, we have*

$$f_3(N) \gg_c (\log N)^c$$

for all sufficiently large N .

Proof. Fix $c \in (0, 1/e)$ and define

$$g(\alpha) := \alpha \log \frac{1}{\alpha}, \quad 0 < \alpha < 1.$$

It is well known that g attains its maximum $1/e$ at $\alpha = 1/e$, and g is continuous on $(0, 1)$. Hence we may choose $\alpha \in (0, 1)$ such that

$$g(\alpha) > c.$$

Let $\varepsilon > 0$ be such that

$$g(\alpha) - 2\varepsilon > c.$$

For each sufficiently large N , set

$$L := \log \log N, \quad t := \lfloor \alpha L \rfloor,$$

so that $t \rightarrow \infty$ and $t = \alpha L + O(1)$, and define

$$x := N^{1/t}.$$

Then $x \rightarrow \infty$ as $N \rightarrow \infty$. We also define $P_0 := \lceil t^2 \rceil$, and work with primes in the interval $[P_0, x]$.

For $y \geq 2$, let

$$H(y) := \sum_{p \leq y} \frac{1}{p}$$

denote the harmonic sum over primes $\leq y$. By Mertens' theorem,

$$H(y) = \log \log y + M + O\left(\frac{1}{\log y}\right) \quad (y \rightarrow \infty),$$

for some absolute constant M .

We first estimate $H(x)$. Note that

$$\log \log x = \log \left(\frac{\log N}{t} \right) = \log \log N - \log t = L - \log t.$$

Since $t = \alpha L + O(1)$, we have $\log t = \log L + O(1)$ and hence

$$\log \log x = L - \log L + O(1),$$

so in particular

$$H(x) = \log \log x + O(1) = L + O(\log L).$$

Let

$$H_{>P_0}(x) := \sum_{P_0 \leq p \leq x} \frac{1}{p} = H(x) - H(P_0^-),$$

where $H(P_0^-) = \sum_{p < P_0} 1/p$. By Mertens' theorem applied at $y = P_0$,

$$H(P_0^-) = \log \log P_0 + O(1) = \log \log(t^2) + O(1) = \log \log t + O(1) = O(\log \log L),$$

so

$$H_{>P_0}(x) = L + O(\log L).$$

Consequently,

$$\frac{H_{>P_0}(x)}{t} = \frac{L + O(\log L)}{\alpha L + O(1)} = \frac{1}{\alpha} + O\left(\frac{\log L}{L}\right) = \frac{1}{\alpha} + o(1) \quad (N \rightarrow \infty). \quad (2.1)$$

List the primes in $[P_0, x]$ in increasing order as

$$q_1 < q_2 < \cdots < q_M \leq x.$$

We now partition $\{q_1, \dots, q_M\}$ into t disjoint subsets P_1, \dots, P_t by the following greedy algorithm.

Initialize $P_i := \emptyset$ and $S_i := 0$ for $1 \leq i \leq t$. For $j = 1, 2, \dots, M$, having already assigned q_1, \dots, q_{j-1} to some of the P_i , choose an index $i_j \in \{1, \dots, t\}$ such that S_{i_j} is minimal, and put $q_j \in P_{i_j}$, updating $S_{i_j} \leftarrow S_{i_j} + 1/q_j$.

For each i we write

$$S_i(x) := \sum_{p \in P_i} \frac{1}{p}.$$

We claim that for every N ,

$$\max_{1 \leq i \leq t} S_i(x) - \min_{1 \leq i \leq t} S_i(x) \leq \frac{1}{P_0}. \quad (2.2)$$

This is proved by induction on j . Let $S_i^{(j)}$ denote the sums after the first j primes have been assigned, and let $M_j := \max_i S_i^{(j)}$, $m_j := \min_i S_i^{(j)}$. Initially $M_0 = m_0 = 0$. Suppose $M_{j-1} - m_{j-1} \leq 1/P_0$. When we assign q_j of weight $w_j = 1/q_j \leq 1/P_0$ to a bucket i_j with $S_{i_j}^{(j-1)} = m_{j-1}$, we obtain $S_{i_j}^{(j)} = m_{j-1} + w_j$ and all other $S_i^{(j)}$ remain unchanged. Thus $m_j \geq m_{j-1}$ and

$$M_j = \max\{M_{j-1}, m_{j-1} + w_j\}.$$

If $M_{j-1} \geq m_{j-1} + w_j$, then $M_j = M_{j-1}$ and

$$M_j - m_j \leq M_{j-1} - m_{j-1} \leq \frac{1}{P_0}.$$

If $M_{j-1} < m_{j-1} + w_j$, then $M_j = m_{j-1} + w_j$ and

$$M_j - m_j \leq (m_{j-1} + w_j) - m_{j-1} = w_j \leq \frac{1}{P_0}.$$

In either case $M_j - m_j \leq 1/P_0$, completing the induction and proving (2.2).

Let $m(x) := \min_i S_i(x)$. Since $H_{>P_0}(x) = \sum_{i=1}^t S_i(x) \geq tm(x)$, we have

$$m(x) \leq \frac{H_{>P_0}(x)}{t}.$$

Combining this with (2.2) yields, for every i ,

$$S_i(x) \geq m(x) \geq \frac{H_{>P_0}(x)}{t} - \frac{1}{P_0}. \quad (2.3)$$

Using (2.1) and the fact that $P_0 \geq t^2$, hence $1/P_0 \leq 1/t^2 \rightarrow 0$, we deduce that

$$S_i(x) \geq \frac{1}{\alpha} + o(1) \quad (N \rightarrow \infty), \quad (2.4)$$

uniformly in $1 \leq i \leq t$.

By continuity of the logarithm, there exists $\delta > 0$ such that

$$\log\left(\frac{1}{\alpha} - \delta\right) \geq \log \frac{1}{\alpha} - \varepsilon.$$

From (2.4) we have $S_i(x) \geq \frac{1}{\alpha} - \delta$ for all $1 \leq i \leq t$ and all sufficiently large N , and thus

$$\log S_i(x) \geq \log \frac{1}{\alpha} - \varepsilon \quad (1 \leq i \leq t) \quad (2.5)$$

for all sufficiently large N .

Define

$$A_N := \left\{ n = p_1 \cdots p_t : p_i \in P_i \text{ for } 1 \leq i \leq t \right\}.$$

Each p_i is a prime $\leq x$, so any $n \in A_N$ satisfies

$$n \leq x^t = N,$$

and hence $A_N \subset \{1, \dots, N\}$.

We claim that A_N contains no triple $\{a, b, c\}$ of distinct elements with all pairwise least common multiples equal, so A_N is admissible in the definition of $f_3(N)$. Indeed, write

$$a = \prod_{i=1}^t p_i^{(a)}, \quad b = \prod_{i=1}^t p_i^{(b)}, \quad c = \prod_{i=1}^t p_i^{(c)},$$

where $p_i^{(\cdot)} \in P_i$ for each i . For a fixed i , consider the primes from P_i appearing in the pairwise least common multiples. In $\text{lcm}(a, b)$ the primes from P_i form the set $\{p_i^{(a)}, p_i^{(b)}\}$ (with multiplicities ignored), and similarly for $\text{lcm}(a, c)$ and $\text{lcm}(b, c)$ we obtain the sets $\{p_i^{(a)}, p_i^{(c)}\}$ and $\{p_i^{(b)}, p_i^{(c)}\}$ respectively.

If

$$\text{lcm}(a, b) = \text{lcm}(a, c) = \text{lcm}(b, c),$$

then for each i these three sets of primes from P_i must be equal:

$$\{p_i^{(a)}, p_i^{(b)}\} = \{p_i^{(a)}, p_i^{(c)}\} = \{p_i^{(b)}, p_i^{(c)}\}.$$

A simple check shows that three two-element sets of the form $\{x, y\}, \{x, z\}, \{y, z\}$ can be equal only if $x = y = z$. Thus $p_i^{(a)} = p_i^{(b)} = p_i^{(c)}$ for each i , and hence $a = b = c$, contradicting distinctness. Therefore A_N is admissible, and

$$f_3(N) \geq \sum_{n \in A_N} \frac{1}{n}.$$

We have

$$\sum_{n \in A_N} \frac{1}{n} = \sum_{p_1 \in P_1} \cdots \sum_{p_t \in P_t} \frac{1}{p_1 \cdots p_t} = \prod_{i=1}^t \sum_{p \in P_i} \frac{1}{p} = \prod_{i=1}^t S_i(x).$$

Taking logarithms and using (2.5), we obtain, for all sufficiently large N ,

$$\log \left(\sum_{n \in A_N} \frac{1}{n} \right) = \sum_{i=1}^t \log S_i(x) \geq t \left(\log \frac{1}{\alpha} - \varepsilon \right).$$

Recalling that $t = \alpha L + O(1)$ with $L = \log \log N$, we deduce

$$\log \left(\sum_{n \in A_N} \frac{1}{n} \right) \geq (\alpha L + O(1)) \left(\log \frac{1}{\alpha} - \varepsilon \right) = (\alpha \log \frac{1}{\alpha} - \alpha \varepsilon + o(1)) L$$

as $N \rightarrow \infty$. Hence, for all sufficiently large N ,

$$\sum_{n \in A_N} \frac{1}{n} \geq \exp((g(\alpha) - 2\varepsilon)L) = (\log N)^{g(\alpha) - 2\varepsilon}.$$

Since $g(\alpha) - 2\varepsilon > c$, we have

$$(\log N)^{g(\alpha) - 2\varepsilon} = (\log N)^c (\log N)^{g(\alpha) - 2\varepsilon - c} \geq (\log N)^c$$

for all sufficiently large N (because $g(\alpha) - 2\varepsilon - c > 0$ is fixed). Thus there exists $N_0(c)$ such that for all $N \geq N_0(c)$,

$$f_3(N) \geq \sum_{n \in A_N} \frac{1}{n} \gg_c (\log N)^c,$$

where the implied constant may depend on c but not on N . This proves the theorem. \square

REFERENCES

- [1] T. F. Bloom, Erdős Problem #856, <https://www.erdosproblems.com/856>, accessed 2025-12-10.
- [2] Erdős, Paul, Some extremal problems in combinatorial number theory. Mathematical Essays Dedicated to A. J. Macintyre (1970), 123–133.

APPENDIX A. COMMENTS (15:18 ON 09 DEC 2025)

For $k = 3$ one can obtain a lower bound stronger than $\gg \log \log N$ from Bloom's comment (12:38 on 09 Dec 2025) by looking at products of two primes from different congruence classes. Fix, for instance,

$$P_1 := \{p \text{ prime} : p \equiv 1 \pmod{8}\}, \quad P_2 := \{q \text{ prime} : q \equiv 3 \pmod{8}\},$$

and for each N set

$$A_N := \{n = pq \leq N : p \in P_1, q \in P_2\}.$$

I claim that A_N contains no triple $\{a, b, c\}$ with all pairwise least common multiples equal. Indeed, write

$$a = p_1 q_1, \quad b = p_2 q_2, \quad c = p_3 q_3$$

with $p_i \in P_1$ and $q_i \in P_2$. In the prime factorization of $\text{lcm}(a, b)$, the contribution from primes in P_1 is exactly the set $\{p_1, p_2\}$, and similarly for the other two lcm's we obtain the sets $\{p_1, p_3\}$ and $\{p_2, p_3\}$. These three two-element sets can only be equal if $p_1 = p_2 = p_3$, and the same argument applied to the primes in P_2 forces $q_1 = q_2 = q_3$, hence $a = b = c$. Thus A_N is admissible for the definition of $f_3(N)$. Now

$$f_3(N) \geq \sum_{n \in A_N} \frac{1}{n} = \sum_{\substack{p \in P_1 \\ q \in P_2 \\ pq \leq N}} \frac{1}{pq} \geq \sum_{\substack{p \in P_1 \\ p \leq \sqrt{N}}} \frac{1}{p} \sum_{\substack{q \in P_2 \\ q \leq N/p}} \frac{1}{q}.$$

By Mertens' theorem in arithmetic progressions one has

$$\sum_{\substack{p \leq x \\ p \equiv a \pmod{8}}} \frac{1}{p} = \frac{1}{\varphi(8)} \log \log x + O(1) = \frac{1}{4} \log \log x + O(1)$$

for each residue class a coprime to 8. In particular, there exists a constant $c_0 > 0$ and $x_0 \geq 2$ such that, for all $x \geq x_0$ and all $a \in \{1, 3, 5, 7\}$,

$$\sum_{\substack{r \leq x \\ r \equiv a \pmod{8}}} \frac{1}{r} \geq c_0 \log \log x.$$

Assume N is large enough that $\sqrt{N} \geq x_0$. Then, for any prime $p \in P_1$ with $p \leq \sqrt{N}$, we have $N/p \geq \sqrt{N} \geq x_0$, so

$$\sum_{\substack{q \in P_2 \\ q \leq N/p}} \frac{1}{q} \geq c_0 \log \log \left(\frac{N}{p} \right).$$

Since $p \leq \sqrt{N}$, we have $N/p \in [\sqrt{N}, N]$, and hence

$$\log \log \left(\frac{N}{p} \right) \geq \log \log \sqrt{N} = \log \log N - \log 2.$$

Thus, enlarging N further if necessary, we may absorb the additive constant $\log 2$ into the implicit constant and obtain

$$\sum_{\substack{q \in P_2 \\ q \leq N/p}} \frac{1}{q} \gg \log \log N \quad \text{for all } p \in P_1 \text{ with } p \leq \sqrt{N}.$$

Similarly, applying the same Mertens-type estimate with $x = \sqrt{N}$ and $a = 1$ gives

$$\sum_{\substack{p \in P_1 \\ p \leq \sqrt{N}}} \frac{1}{p} \geq c_0 \log \log \sqrt{N} = c_0 (\log \log N - \log 2) \gg \log \log N$$

for all sufficiently large N . Combining these lower bounds, we deduce

$$f_3(N) \geq \sum_{n \in A_N} \frac{1}{n} \gg (\log \log N) \cdot (\log \log N) = (\log \log N)^2.$$

APPENDIX B. COMMENTS (15:30 ON 09 DEC 2025)

I think the above construction extends in a straightforward way to give higher powers of $\log \log N$. Indeed, fix an integer $t \geq 1$ and choose t distinct residue classes $a_1, \dots, a_t \pmod{q}$ with $(a_i, q) = 1$. For each $1 \leq i \leq t$ let

$$P_i := \{p \text{ prime} : p \equiv a_i \pmod{q}\},$$

and define

$$A_N^{(t)} := \{n = p_1 \cdots p_t \leq N : p_i \in P_i \ (1 \leq i \leq t)\}.$$

By the same argument as in the case $t = 2$, one checks that $A_N^{(t)}$ contains no triple $\{a, b, c\}$ with all pairwise least common multiples equal. Furthermore, using Mertens' theorem in arithmetic progressions in each of the t sets P_i and restricting to the box $p_i \leq N^{1/t}$, one obtains

$$\sum_{n \in A_N^{(t)}} \frac{1}{n} \geq \prod_{i=1}^t \sum_{\substack{p \in P_i \\ p \leq N^{1/t}}} \frac{1}{p} \gg_t (\log \log N)^t$$

for all sufficiently large N , where the implied constant depends only on t . Hence for every fixed t there exists a constant $c_t > 0$ such that

$$f_3(N) \geq c_t (\log \log N)^t$$

for all sufficiently large N .

APPENDIX C. COMMENTS (08:19 ON 10 DEC 2025)

Thanks to Bloom for the very helpful heuristic comment (17:24 on 09 Dec 2025). Motivated by his suggestion that choosing t to be a small multiple of $\log \log N$ might lead to a power of $\log N$, one can in fact turn this into a completely rigorous construction.

Theorem C.1. *There exist constants $c > 0$ such that*

$$f_3(N) \gg_c (\log N)^c$$

for all sufficiently large N .

Proof. Fix a real number $B > 1$, and choose a small parameter $\gamma \in (0, B/2)$. Set

$$B' := B + \gamma.$$

Next choose a real number $P_0 > 0$ so large that for every prime $p \geq P_0$ we have

$$\frac{1}{p} \leq \gamma.$$

(This is possible since $1/p \rightarrow 0$ as $p \rightarrow \infty$.)

We will work with primes $\geq P_0$, and we write

$$H(x) := \sum_{p \leq x} \frac{1}{p}$$

for the usual harmonic sum over primes. By Mertens' theorem we have

$$H(x) = \log \log x + O(1) \quad (x \rightarrow \infty).$$

Let us now choose a small constant $\kappa := \frac{1}{16B'}$. For each large N we set $L := \log \log N$, $t := \lfloor \kappa L \rfloor$, and define $x := N^{1/t}$. For N sufficiently large we have $t \geq 1$ and $x \rightarrow \infty$.

First, we have

$$\log \log x = \log \left(\frac{\log N}{t} \right) = \log \log N - \log t = L - \log t.$$

Since $t = \kappa L + O(1)$, we have $\log t = \log L + O(1)$, so

$$\log \log x = L - (\log L + O(1)) \geq \frac{L}{2}$$

for all sufficiently large N . Hence Mertens' theorem gives

$$H(x) = \log \log x + O(1) \geq \frac{L}{2} - C_0$$

for some constant C_0 and all large N .

Let

$$S_{\text{tot}}(x) := \sum_{P_0 \leq p \leq x} \frac{1}{p} = H(x) - H(P_0^-),$$

where $H(P_0^-)$ is a fixed constant depending only on P_0 . Thus

$$S_{\text{tot}}(x) \geq \frac{L}{2} - C_1$$

for some constant C_1 and all large N . In particular, there exists N_1 such that for all $N \geq N_1$,

$$S_{\text{tot}}(x) \geq \frac{L}{4}. \quad (\text{eq:Stot-large})$$

On the other hand

$$tB' \leq (\kappa L)B' \leq \frac{L}{16}$$

for all large N , since $\kappa = 1/(16B')$ and $t \leq \kappa L$. Thus, by Eq. (eq:Stot-large), for all $N \geq N_1$ we have

$$S_{\text{tot}}(x) \geq tB'. \quad (\text{eq:Stot-vs-tBprime})$$

List the primes in $[P_0, x]$ in increasing order as

$$q_1 < q_2 < \cdots < q_M \leq x.$$

We will construct, by a greedy algorithm, disjoint subsets $P_1, \dots, P_k \subset \{q_1, \dots, q_M\}$, where $k \geq t$, such that each P_i satisfies

$$B \leq \sum_{p \in P_i} \frac{1}{p} < B'. \quad (\text{eq:Pi-bounds})$$

We proceed as follows. Set $P_1 = \emptyset$ and $i = 1$. Process the primes q_1, q_2, \dots, q_M in order, adding q_j to the current set P_i as long as

$$\sum_{p \in P_i} \frac{1}{p} < B.$$

As soon as this sum first becomes $\geq B$, we stop filling P_i and, if $i < t$, we start a new set P_{i+1} from the next unused prime in the list. We continue in this way until either we have formed t such sets or we have exhausted all primes in $[P_0, x]$.

By construction, whenever a set P_i is completed, its harmonic sum satisfies $\sum_{p \in P_i} 1/p \geq B$, and since each added prime p satisfies $p \geq P_0$ and hence $1/p \leq \gamma$, we have

$$\sum_{p \in P_i} \frac{1}{p} < B + \gamma = B'.$$

Thus Eq. (eq:Pi-bounds) holds for every completed set P_i .

We now show that for $N \geq N_1$ the algorithm must produce at least t such sets. Suppose instead that the process terminates after forming only $k < t$ completed sets P_1, \dots, P_k . There are two possibilities: Case 1: There is no unfinished set at termination. Then all primes in $[P_0, x]$ have been used to form the sets P_1, \dots, P_k , and

$$S_{\text{tot}}(x) = \sum_{i=1}^k \sum_{p \in P_i} \frac{1}{p}.$$

Using Eq. (eq:Pi-bounds), we obtain

$$S_{\text{tot}}(x) < \sum_{i=1}^k B' = kB' \leq (t-1)B' < tB'.$$

Case 2: There is an unfinished $(k+1)$ -st set at termination. In this case, after forming P_1, \dots, P_k , there remain some primes in $[P_0, x]$ which have been partially used to form a final set P_{k+1} with

$$\sum_{p \in P_{k+1}} \frac{1}{p} < B,$$

since otherwise P_{k+1} would also be a completed set. Let P_{k+1} denote this (possibly empty) set of remaining primes. Then

$$S_{\text{tot}}(x) = \sum_{i=1}^k \sum_{p \in P_i} \frac{1}{p} + \sum_{p \in P_{k+1}} \frac{1}{p}.$$

Using Eq. (eq:Pi-bounds) again, we obtain

$$S_{\text{tot}}(x) < kB' + B \leq kB' + B' = (k+1)B' \leq tB'.$$

In both cases we conclude that $S_{\text{tot}}(x) < tB'$, which contradicts Eq. (eq:Stot-vs-tBprime). Therefore, for every $N \geq N_1$ the greedy algorithm produces at least t disjoint subsets P_1, \dots, P_t of primes in $[P_0, x]$, each satisfying Eq. (eq:Pi-bounds).

Now define

$$A_N := \left\{ n = p_1 \cdots p_t : p_i \in P_i \text{ for } 1 \leq i \leq t \right\}.$$

Since every prime in P_i is $\leq x$, any $n \in A_N$ satisfies

$$n \leq x^t = N,$$

so $A_N \subset \{1, \dots, N\}$. Clearly, A_N contains no triple $\{a, b, c\}$ of distinct elements with all pairwise least common multiples equal.

Finally, we have

$$\sum_{n \in A_N} \frac{1}{n} = \sum_{p_1 \in P_1} \cdots \sum_{p_t \in P_t} \frac{1}{p_1 \cdots p_t} = \prod_{i=1}^t \sum_{p \in P_i} \frac{1}{p}.$$

By Eq. (eq:Pi-bounds), each factor satisfies

$$\sum_{p \in P_i} \frac{1}{p} \geq B,$$

so

$$\sum_{n \in A_N} \frac{1}{n} \geq B^t.$$

Since A_N is admissible, this implies

$$f_3(N) \geq B^t.$$

Recalling that $t = \lfloor \kappa L \rfloor$ with $L = \log \log N$, we obtain

$$B^t \geq B^{\kappa L - 1} = \exp((\kappa L - 1) \log B) = (\log N)^{\kappa \log B} B^{-1}.$$

Thus there exists $c := \kappa \log B > 0$ and a constant $C > 0$ (depending only on B, γ, P_0) such that

$$f_3(N) \geq C (\log N)^c$$

for all sufficiently large N . Q.E.D. □

For example, if we let $B = 2$ and $\gamma = \frac{1}{2}$ in the proof of Theorem 1, then $B' = B + \gamma = \frac{5}{2}$ and $\kappa = \frac{1}{16B'} = \frac{1}{40}$. Thus

$$f_3(N) \gg (\log N)^c, \quad \text{where } c = \frac{1}{40} \log 2 > 0.017.$$

Of course, with a more refined construction one can improve this value of c , but it seems that this method can at best yield exponents in the range $0 < c < 1/e$.

SCHOOL OF MATHEMATICS AND STATISTICS, XI'AN JIAOTONG UNIVERSITY, XI'AN 710049, P. R. CHINA
Email address: `tang_quanyu@163.com`