

Part I

A First Course in General Relativity¹

Chapter 1

Special Relativity

On “Principle of relativity (Galileo)”

Galilean invariance

Newton’s laws of motion hold in all frames related to one another by a Galilean transformation. In other words, all frames related to one another by such a transformation are inertial (meaning, Newton’s equation of motion is valid in these frames).² The proof has been given by the book on page 2.

1.5 - Construction of the coordinates used by another observer

Why would the tangent of the angle is the speed in Fig. 1.2?


Suppose \mathcal{O} and $\bar{\mathcal{O}}$ both start out at the same position where $\bar{\mathcal{O}}$ moves along the x at some speed. After t_1 , observer \mathcal{O} sees $\bar{\mathcal{O}}$ at position x_1 :

$$\bar{\mathcal{O}}_1 = (x_1, t_1)$$

Observer $\bar{\mathcal{O}}$, however, still sees themselves at $x = 0$:

$$\bar{\mathcal{O}}_1 = (0, t_1)$$

By definition where “ \bar{t} is the locus of events at constant $\bar{x} = 0$ ”, \bar{t} is the straight line that passes the origin and the (x_1, t_1) :



1.6 - Why $(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 - (\Delta t)^2 = 0$ for two events in the same light beam?

Let's say, in a simplified 1D case, event $\mathcal{E} = (x_0, t_0)$ and $\mathcal{P} = (x_1, t_1)$.

$$(\Delta x)^2 - (\Delta t)^2 = (x_1 - x_0)^2 - (t_1 - t_0)^2$$

Since the speed of light is 1,

$$(x_1 - x_0)^2 - (t_1 - t_0)^2 = (x_1 - x_0)^2 - (t_1 \times 1 - t_0 \times 1)^2 = (x_1 - x_0)^2 - (x_1 - x_0)^2 = 0$$

Why do we have a 2nd term in equation 1.3 on p.10?

$$\Delta \bar{s}^2 = \sum_{\alpha=0}^3 \sum_{\beta=0}^3 M_{\alpha\beta} (\Delta x^\alpha) (\Delta x^\beta)$$

Bibliography

¹ Bernard Schutz. *[A First Course on General Relativity](#)*. 2009.

² Wikipedia. [Galilean invariance](#).