

1     **Dispersal capacities of anadromous Allis shad population inferred from a**  
2                                   **coupled genetic and otolith approach**

3

4     **Jean Martin<sup>1</sup>, Quentin Rougemont<sup>2</sup>, Hilaire Drouineau<sup>1</sup>, Sophie Launey<sup>2</sup>, Philippe**  
5     **Jatteau<sup>1</sup>, Gilles Bareille<sup>3</sup>, Sylvain Berail<sup>3</sup>, Christophe Pécheyran<sup>3</sup>, Eric Feunteun<sup>4</sup>,**  
6     **Séverine Roques<sup>1</sup>, David Clavé<sup>5</sup>, David Jose Nachón<sup>6</sup>, Carlos Antunes<sup>7</sup>, Micaela Mota<sup>7</sup>,**  
7                                   **Elodie Réveillac<sup>4,8</sup>, Françoise Daverat<sup>1</sup>**

8

9     <sup>1</sup> **IRSTEA, Estuarine Ecosystems and Diadromous Fish Research Unit, 50 avenue de**  
10                                   **Verdun, 33612 Cestas Cedex, FRANCE, jean.martin@irstea.fr ,**  
11     **francoise.daverat@irstea.fr, severine.roques@irstea.fr, hilaire.drouineau@irstea.fr,**  
12                                   **philippe.jatteau@irstea.fr**

13     <sup>2</sup> **Institut National de la Recherche Agronomique, Unité Mixte de Recherche, 985**  
14     **Ecologie et Sante des Ecosystemes, 65 rue de Saint Briec, F-35042 Rennes Cedex,**  
15                                   **France, Sophie.Launey@rennes.inra.fr, Quentin.Rougemont@rennes.inra.fr**

16     <sup>3</sup> **Laboratoire de Chimie Analytique Bio-Inorganique et Environnement (LCABIE),**  
17                                   **UMR CNRS-UPPA 5254 IPREM, Hélioparc Pau Pyrénées, 64053 Pau, France,**  
18     **gilles.bareille@univ-pau.fr, christophe.pecheyran@univ-pau.fr, sylvain.berail@univ-**  
19                                   **pau.fr**

20     <sup>4</sup> **Muséum National d'Histoire Naturelle, UMR BOREA 7208, Station Marine de Dinard,**  
21                                   **Dinard, France, Eric.Feunteun@mnhn.fr**

22 <sup>5</sup> Association MIGADO, 18 Ter rue de la Garonne, BP 95, 47520 Le Passage, France,  
23 [clave.migado@orange.fr](mailto:clave.migado@orange.fr)

24 <sup>6</sup> Hydrobiology Station “Encoro do Con”, Castroagudín s/n, 36617 Vilagarcía de Arousa,  
25 Pontevedra, Spain, [davidjose.nachon@usc.es](mailto:davidjose.nachon@usc.es)

26 <sup>7</sup> Centre of Marine and Environmental Research (CIIMAR/CIMAR), University of  
27 Porto, Rua dos Bragas 289, P 4050-123 Porto, Portugal, [cantunes@ciimar.up.pt](mailto:cantunes@ciimar.up.pt),  
28 [mmota@ciimar.up.pt](mailto:mmota@ciimar.up.pt)

29 <sup>8</sup> Agrocampus Ouest, UMR985 ESE Ecologie et Santé des Ecosystèmes, 65 rue de Saint-  
30 Brieuc, CS 84215, 35042 Rennes cedex, France, [elodie.reveillac@agrocampus-ouest.fr](mailto:elodie.reveillac@agrocampus-ouest.fr)

31  
32 \* Corresponding author. Email: [francoise.daverat@irstea.fr](mailto:francoise.daverat@irstea.fr)

33

34

## Abstract

Understanding dispersal capacities for migratory species is crucial for their management. By coupling otolith microchemistry and microsatellite genetic analyses, we provided information on snapshot and long-term dispersal capacity of Allis shad, an anadromous clupeid in decline throughout its distribution range. The allocation of natal origin was obtained from water chemistry, signatures in otoliths of juveniles and spawners within a Bayesian model. The majority of adults were assigned to a source river with high degrees of confidence, only 4% were undetermined. Otolith natal origins were used to define a population baseline by grouping individuals from the same natal river and not from the same sampling location as usually done. While *Alosa alosa* exhibited a high level of natal site fidelity, this species showed weak genetic structure which supported the evidence of a significant flow of strayers between river basins in vicinity or at longer distances. However, long distance straying was probable but not frequent. In a context of global change, straying would be a key mechanism to drive dispersal and allow resilience of Allis shad populations.

**Keywords:** Natal origins, Straying pattern, Allis shad, Otolith chemistry, Microsatellites.

## Introduction

Understanding population structure and connectivity for migratory species across broad geographic areas (e.g., species' distribution) is of major interest for their management and conservation. Anadromous fish are defined by McDowall (2008) as fish which spend most of their life at sea and migrate to freshwater to reproduce. Philopatry to natal sites and straying behavior, defined as adult migration to non-natal rivers, are two fundamental life-history traits of most anadromous fishes that jointly convey adaptive advantages to their persistence. Selection in spatially heterogeneous environment leads to local adaptation which can be enhanced by natal site fidelity hence providing fitness benefits (Hendry et al. 2004). On the other hand, straying allows for colonization of new habitats, the maintenance of genetic diversity, and can mitigate spatial and temporal variation in habitat quality (Keefer and Caudill 2014). Regardless whether fish are straying by habitat choice or by mischance of returning home through inadequate imprinting, accumulating evidence suggests that the degree to which straying is undertaken probably varies within and among populations of many fish species (McDowall 2001). Thus, understanding this straying behavioral diversity at the individual scale may be necessary to ensure anadromous fishes resilience in the face of anthropogenic threats and large scale climatic and environmental changes (Hamann and Kennedy 2012).

The combination of otolith microchemistry and genetic markers has shown promise in identifying patterns of connectivity among populations at different scales (Bradbury et al. 2008; Selkoe et al. 2008; Barnett-Johnson et al. 2010; Hughes et al. 2014). Genetic studies provide insight on long-term connectivity patterns on evolutionary time scales whereas chemical tracers can reveal dispersal patterns within the time scale of an individual lifetime (Feyrer et al. 2007). However, magnitudes of dispersal or straying rate present particular challenges to genetic investigation of connectivity for anadromous species (Selkoe et al. 2008;

77 Humston et al. 2010). The use of otolith based techniques and genetic markers on the same  
78 individuals may provide valuable information about the dispersal pattern of a given fish  
79 during its lifetime, but also about longer term gene flow between populations (Fromentin et  
80 al. 2009).

81 Otoliths are paired aragonitic structures in the inner ear of bony fish that grow by  
82 continuous accretion of metabolically inert material (Campana 1999). Otolith chemical  
83 composition reflects, to some degree, the chemistry of ambient waters at the time of material  
84 deposition (Bath et al. 2000; Walther and Thorrold 2006). Otoliths from fish hatched in  
85 chemically distinct water masses will record unique signatures, reflecting their habitats. The  
86 use of trace element ratios recorded in otoliths has provided separation of anadromous fish  
87 according to their natal river (Barnett-Johnson et al. 2008; Walther et al. 2008). Strontium and  
88 Barium have been amongst the most powerful elements because their concentrations in  
89 otoliths are largely influenced by their respective metal-to-Ca ratios in the ambient water  
90 (Bath et al. 2000; Dorval et al. 2007; Elsdon et al. 2008; Martin et al. 2013b). In addition,  
91 natural Sr isotopic signatures have proved to be useful as fish markers, particularly for species  
92 that inhabit freshwater during their juvenile stages (Ingram and Weber 1999; Kennedy et al.  
93 2000; Martin et al. 2013a). Otolith  $^{87}\text{Sr}:^{86}\text{Sr}$  ratios directly reflect dissolved ambient ratios  
94 (Blum et al. 2000), and have been found to remain stable over the years, making them “ideal”  
95 spatial markers for characterizing natal sources (Kennedy et al. 2000). Fine-scale geographic  
96 discrimination of rivers could be achieved by the combined use of elemental and Sr isotope  
97 ratios.

98 Microsatellites are neutral genetic markers not affected by selective pressures. They  
99 consist of short tandem repeat sequences of nucleotides (2-5 base pairs repeated units)  
100 (O'Connell and Wright 1997) that undergo fast mutation rates and thus lead to high levels of  
101 polymorphism. They are among the most commonly used markers to test the level of among-

102 population connectivity (Bentzen et al. 1996; Selkoe and Toonen 2006; Bradbury et al. 2014)  
 103 and have been useful for fine-scale population discrimination in anadromous fishes (Banks et  
 104 al. 2000; Spidle et al. 2003). However, the success of genetics-based approaches to quantify  
 105 natal homing is dependent on the levels of interbreeding among populations since even very  
 106 low exchange rates can prevent populations' differentiation (Collins et al. 2013).

107 In the present study, we integrated multiple approaches (i.e. otolith microchemistry,  
 108 and genetics) to assess the dispersal capacity of Allis shad (*Alosa alosa*), an anadromous  
 109 clupeid in decline throughout its distribution range (Baglinière et al. 2003). This species has a  
 110 pelagic marine existence but upon maturation (from 4 to 6 years) migrates to spawn in the  
 111 higher middle watercourse of rivers. Eggs hatch at spawning site and seaward emigration  
 112 takes place in the summer and fall of their first year of life (Lochet et al. 2009). Allis shad was  
 113 distributed historically in latitude from the North of Morocco to Norway (Baglinière et al.  
 114 2003). Since the middle of the 20th century, the impact of human activities such as  
 115 overfishing, damming of rivers and pollution, has led to a drastic restriction of its distribution  
 116 area (De Groot 2002; Limburg and Waldman 2009). Climate change is also expected to shift  
 117 the species range towards higher latitudes and southern populations are expected to be  
 118 particularly affected by global warming (Lassalle et al. 2008a). The species was classified as  
 119 'vulnerable' and has been included in Appendix III of the Bern Convention as well as in  
 120 Annexes II and V of the European Community Habitats Directive. Currently, the most  
 121 successful breeding populations, excluding relict or residual ones, are distributed across the  
 122 north-east Atlantic coastline in a few rivers in western France and Portugal (Baglinière et al.  
 123 2003).

124 Very little is known about Allis shad population structure in Europe. Divergence in  
 125 life history characteristics among populations was observed with southern populations  
 126 exhibiting larger length at maturity compared to basins further north (Lassalle et al. 2008b).

Studies reporting discrimination of natal origins of European Allis shad populations using otolith microchemistry are scarce and were restricted to the Gironde population, where Tomas et al. (2005) discriminated fish from Garonne and Dordogne Rivers. Genetic studies generally indicated low levels of genetic structure among populations but some authors (Alexandrino et al. 2006; Jolly et al. 2012) described a slight divergence between French and Portuguese populations and a process of isolation-by-distance over 2,000-4,000 km, suggesting that gene flow primarily takes place between neighbouring populations. Although there is a pressing need to understand the migratory dynamics of Allis shad throughout its range for management and conservation purposes, all previous studies have been restricted to a limited geographic area or have compared only a handful of populations.

The main goal of the present study was to couple otolith chemistry with genetic analysis on the same individuals to study population connectivity and dispersal of *A. alosa* over ecological and evolutionary time scales. This integrative approach was used to identify the Allis shad's natal rivers and assess the level of gene flow between populations, and thus determine the temporal and spatial scale over which straying occurs.

## Materials and methods

### Fish and water sampling

Mature Allis shad ( $n = 425$ ) were collected from upstream spawning areas to tidal freshwater portions of 15 rivers along the Atlantic coast, from northern France to Portugal, between April and June 2009 to 2013 (supplementary Figs. S1, S2, Table S1). Spawners were sampled either by sport fishing, by commercial fishermen using a trammel net, or found dead after spawning during summer. Muscle tissues were removed and stored in 95% ethanol. The lengths of shad were measured to the nearest millimeter. Fifty decaying fish carcasses, sampled after spawning, were not available for length and weight measurements. All fish were frozen upon collection.

Juvenile shad ( $n = 26$ ) were sampled in four French rivers (Blavet, Dordogne, Loire and Vilaine) between June and October 2013 (supplementary Table S1). Juveniles from the Minho River in Portugal ( $n = 20$ ) were sampled between 2009 and 2012 from September to January. The Blavet, Dordogne, Loire, Vilaine and Minho Rivers have been sampled for both adults and juveniles. Fish were collected using bongo nets and seines in upper-estuarine regions of each river prior to their downstream migration to the sea.

Spatial variability in water chemistry was investigated across 17 major spawning rivers from throughout the native range of Allis shad. These rivers corresponded to the rivers where fish were sampled. The Charente, Oloron and Nive rivers were sampled despite no fish collection. At each river, water samples were collected, once a month, from late May to September 2013, close to well-known spawning area of Allis shad. At each location, 100 mL of river water was collected for Sr:Ca, Ba:Ca and  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  analyses. Water samples were passed through 0.45  $\mu\text{m}$  Nalgene polytetrafluoroethylene filters (<http://www.vgdusa.com/Nalgene-Syringe-Filters-PTFE-25mm.htm>) with syringes into acid-washed low-density polyethylene (LDPE) bottles and acidified (2%) using concentrated, ultrapure  $\text{HNO}_3$  (JT Baker, Ultrex II; [www.jtbaker.nl](http://www.jtbaker.nl)). Samples were kept on ice in the field and refrigerated upon return to the laboratory.

### Microsatellite DNA genotyping

Genetic analyses were performed on all collected specimens except for Portuguese rivers where no fish tissue was available. Genomic DNA was extracted from ethanol-preserved muscle tissue using a modified Chelex protocol from Estoup et al. (1996). Genotyping was performed using 13 microsatellite markers: Alo1, Alo6, Alo7, Alo9, Alo15, Alo16, Alo17, Alo26, Alo29, Alo32, Alo33, Alo43 and Alo45 (Rougemont et al. 2014) specifically developed for *Alosa alosa* (and its closely related species *A. fallax*) by following the protocol recommended by the authors. We found no evidence of null alleles and scoring



errors due to large allele dropout or stutter peaks by using the software Micro-Checker (Van Oosterhout et al. 2004).

### Otolith preparation and analysis

Frozen fish were thawed and dissected to remove pairs of sagittal otoliths. Otoliths were rinsed in distilled water, air-dried, and one sagitta per fish was embedded in epoxy resin (Araldite 2020, Escil) with the primordial surface downwards. Resin blocks were ground with ultrapure water and sandpaper (1200 – 4000 grit) until the primordium was reached. Finally, otoliths were rinsed with ultrapure water, and then air-dried before being stored in individually labeled plastic vials.

A C-shaped ablation trajectory centered on the primordium (40  $\mu\text{m}$  away from the primordium, ablation width 60  $\mu\text{m}$ ; Fig. 1) was applied on otoliths of all individuals collected (adult and juvenile Allis shad), which avoided the maternal influence on the chemical signature (Kalish 1990; Rieman et al. 1994). We assume that the first feeding mark was always before 40  $\mu\text{m}$  from the primordium for each sample of Allis shad (Lochet et al. 2008). Semi coronas of 60 $\mu\text{m}$  thick (difference between the inner 40  $\mu\text{m}$  and outer radius 100  $\mu\text{m}$ ) were constructed by fast rotation of an 8  $\mu\text{m}$  spot (using 8 concentric semicircles). A UV high-repetition-rate femtosecond laser ablation (fs-LA) system (Nexeya SA, Canejan, France) was employed (Pulse duration: 360fs; wavelength: 257 nm).

Otoliths were analyzed for Sr:Ca and Ba:Ca ratios on a High Resolution (Thermo Scientific, USA) inductively coupled plasma quadrupole mass spectrometer (HR-ICP-MS). A He gas stream carried ablated material to the HR-ICP-MS (carrier gas flow rate 0.68 L $\cdot\text{min}^{-1}$ ). Elemental ratios were quantified by monitoring  $^{43}\text{Ca}$ ,  $^{86}\text{Sr}$  and  $^{138}\text{Ba}$ . Calcium was used as an internal standard to improve the reliability of the concentration measurement (Campana 1999). Elements were standardized to calcium based on the stoichiometry of calcium carbonate (389 000  $\mu\text{g Ca}\cdot\text{g}^{-1}$  otolith) (Brown and Severin 2009): Sr:Ca ( $\text{mg}\cdot\text{g}^{-1}$ ), Ba:Ca

( $\mu\text{g}\cdot\text{g}^{-1}$ ). Quantification of trace elements in otoliths was achieved by external calibration using both carbonate pellets (Barats et al. 2007) and 3 NIST glass standards (610, 612, 616) to ensure the best accuracy. An otolith Certified Reference Material (NIES 22, National Institute for Environmental Studies) was also pelletized and used in the quality control of the analysis of selected trace elements in the fish otolith. The limits of detection ( $\mu\text{g}\cdot\text{g}^{-1}$  in otoliths) achieved in this study were as follows:  $^{86}\text{Sr}$ , 0.05 and  $^{138}\text{Ba}$ , 0.01. They were based on a  $3\sigma$  criterion, where  $\sigma$  is the standard deviation of the mean blank count for each isotope. All the elemental concentrations in the otolith were above the detection limits.

After elemental analysis, the same otoliths were used for  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  ratio and the laser traced out a semi corona of  $60\mu\text{m}$  thick opposed to the one ablated for elemental ratio measurements (Fig. 1). We used a Nu-Plasma multicollector ICP-MS (Nu Instruments, UK) coupled to the same fs-LA system described above. The otoliths were analysed for  $^{87}\text{Sr}$ : $^{86}\text{Sr}$ , following the method used by Martin et al. (2013a). The accuracy of this approach was checked through the analyses of NIES 22 obtained from otoliths of a marine fish (*Lutjanus sebae*). We analyzed NIES 22 pellets ( $n = 36$ ) with the same ablation strategy applied to otoliths and we obtained a mean value of  $0.70927 \pm 0.00011$  2 SD, for  $^{87}\text{Sr}$ : $^{86}\text{Sr}$ . The  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  ratios in pellets fell within the expected range for nearly constant modern sea water (0.70917) (Allègre et al. 2010). Finally, the  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  ratio of NIES 22 pellets was used as an in-house marine carbonate standard and was measured every 6 samples as an external check of reproducibility.

### Water sample preparation and analysis

Water samples were analyzed using solution-based ultra-sensitive inductively coupled plasma mass spectrometer (ICP-MS, Bruker Aurora Elite; www.bruker.com) to measure Ca, Sr and Ba concentrations following protocol described by Martin et al. (2013a). The general performance of the procedure was checked every 10 samples using the certified reference

freshwater SLRS-5 (NRCC; [www.nrc-cnrc.gc.ca](http://www.nrc-cnrc.gc.ca)). External precision (% relative standard deviation, R.S.D.) for the laboratory standard ( $n = 10$ ) were 3% for Ca and Ba, 4% for Sr.

Sr isotope analysis was performed using the Nu-Plasma MC-ICP-MS, and following protocol described by Martin et al. (2013a). Accuracy and precision were monitored with a Standard Reference Material (SRM 987; [www.nist.gov/srm](http://www.nist.gov/srm)). The mean $\pm$ s.d. value of  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  in SRM 987 ( $n = 54$ ) run throughout the analyses was  $0.71034 \pm 0.00003$ , which compares favourably with the accepted value of  $0.71034 \pm 0.00026$ .

## Data analysis

### *Water chemistry analyses*

Measurements of elemental and Sr isotopic ratios in water were statistically analyzed using non-parametric multivariate tests as data were not normally distributed. Water chemistry differences among sites were tested using the non-parametric Mann–Whitney  $U$ -Wilcoxon tests with a Bonferroni adjustment. Geographic differences in multivariate signatures among locations were visualized using canonical discriminant analysis. Canonical variate coefficients provide a useful way to measure the relative importance of each variable to the observed separation among rivers (Walther et al. 2008). Finally, a quadratic discriminant function analysis (QDFA) was employed to discriminate among the different rivers. A QDFA was used because this procedure does not assume homogeneity of covariance matrices and tolerates modest deviations from normality (McGarigal et al. 2000). The discriminant function analysis used a jackknife cross-validation procedure to determine classification accuracy. Data analyses were performed using R software (R. Development Core Team 2013).

### *Bayesian hierarchical mixture model: model specification*

Flash ablation is a laser strategy that allows ablating semi-coronas and sends the corresponding otolith powdered material at once to the ICP-MS (for elemental ratio

measurements) or the multicollector ICP-MS (for Sr isotope ratio measurements). Since we ablated the same period of life on the otolith for both semi-coronas, we used Sr:Ca, Ba:Ca and  $^{87}\text{Sr}:^{86}\text{Sr}$  as a three dimensional value system to characterize the natal origin chemical signature of a fish.

In the following section, braces were used to denote vectors while brackets were used to denote matrices. An approach similar to Pflugeisen and Calder (2013) was used to infer shad natal rivers based on their otolith microchemistry and habitat information. This approach was based on a Bayesian clustering method using otolith elemental compositions and Sr isotopic ratios as fish descriptors and similar measurements made on possible natal rivers as habitat information.

Each adult shad  $a$ , was characterised by a river of capture  $C_{ad}(a)$ , its otolith elemental composition  $\{E_{ad}(a)\}$ , and a Sr isotope ratio  $IR_{ad}(a)$ . The objective was to infer adult natal rivers  $N_{ad}(a)$  based on  $\{E_{ad}(a)\}$  and  $IR_{ad}(a)$ . Similarly, a juvenile  $j$  was characterised by its otolith elemental composition  $\{E_{ju}(j)\}$ , a Sr isotope ratio  $IR_{ju}(j)$  and a natal river  $N_{ju}(j)$  (which was known contrary to adult fish). Each river  $r$ , was also characterised by its elemental signature  $\{E_{ri}(r)\}$  and Sr isotope ratio  $IR_{ri}(r)$ .

We assumed that  $E_{ad}(a)$  and  $IR_{ad}(a)$  given the fish was born in river  $r$  followed respectively a multinormal ( $MN$ ) and a normal ( $N$ ) distribution:

$$(\{E_{ad}(a)\}|N_{ad}(a) = r) \sim MN(\{\alpha\} \cdot \{E_{ri}(r)\} + \{\beta\}, [\Sigma])$$

$$(IR_{ad}(a)|N_{ad}(a) = r) \sim N(IR_{ri}(r), \sigma_i)$$

with  $\{\alpha\}$  and  $\{\beta\}$  the partition coefficients of element assimilation (Farrell and Campana 1996; Bath et al. 2000). Uninformative priors were used for  $[\Sigma]$  and  $\sigma_i$  ( $i$  for isotope):

$$\sigma_i \sim InvGamma(0.01, 0.01)$$

$$[\Sigma] \sim InvWishart([I_n], n)$$

with  $[I_n]$  the identity matrix and  $n$  the number of element considered in the study. Vague

uniform priors were used for  $\{\alpha\}$  and  $\{\beta\}$ , respectively between  $[0, 2]$  and between  $[-3, 3]$ . No partition coefficients were required for isotopic ratio since otolith isotopic ratio is equal to the water ratio (Blum et al. 2000).

For juveniles, we assumed that:

$$\{E_{ju}\}(j) \sim MN\left(\{\alpha\} \cdot \{E_{ri}\}(N_{ju}(j)) + \{\beta\}, [\Sigma]\right)$$

$$IR_{ju}(j) \sim N\left(IR_{ri}(N_{ju}(j)), \sigma_i\right)$$

The natal origin of an adult caught in river  $C_{ad}(r)$  was given by:

$$N_{ad}(a) \sim Categorical(\{\theta_{C_{ad}(a)}\})$$

$\{\theta_r\}$  was a vector that contains the a priori probabilities  $\{\theta_r(1), \dots, \theta_r(n_r)\}$  that an adult caught in river  $r$  was born in each of the  $n_r$  rivers.

Similarly to Pflugeisen and Calder (2013), an uninformative prior was used for  $\{\theta_r\}$ :

$$\{\theta_r\} \sim Dirichlet(\{\gamma_k\})$$

with  $\gamma_1 = \dots = \gamma_{n_r} = \frac{1}{n_r}$

#### *Model fitting and convergence diagnostics*

Three independent parallel chains were run for 20 000 iterations after a burn-in period of 10 000 iterations. The model was implemented using the software JAGS (Plummer 2003) and was run using runjags (Denwood 2013), a package that facilitated the communication between JAGS and R statistical software. Convergence diagnostics were made out using the coda library (Plummer et al. 2006). Gelman and Rubin (1992) tests and visual inspection of a posteriori distribution were carried out. In addition, visual inspections of rivers reallocation were carried out to check consistency between the three chains.  $\{E_{ri}(r)\}$ ,  $\{E_{ju}(j)\}$  and  $\{E_{ad}(a)\}$  were previously scaled and centered for each element. Centering covariates is useful in Bayesian inference to decrease the correlation between regression parameters. Scaling was required to have a common scale of variations among elements.  $\{\alpha\}$ ,  $\{\beta\}$ ,  $[\Sigma]$ ,  $\sigma_i$  and  $N_{ad}(a)$

were monitored.  $\{\alpha\}$ ,  $\{\beta\}$  provided information on the partition function.  $[\Sigma]$ ,  $\sigma_i$  provided information on the relative influence of each element and Sr isotopes. Finally,  $N_{aa}(a)$  corresponded to the adult shad reallocations.

### *Genetic data analysis*

We first tested deviation from Hardy-Weinberg Equilibrium (HWE) and genotypic disequilibrium between pairs of loci for each population using Genepop (Rousset 2008). Basic indices of population genetic diversity, namely observed heterozygosity and unbiased expected heterozygosity ( $H_o$  and  $H_{nb}$ ) were computed using Genetix (Belkhir et al. 2004). Allelic richness ( $A_r$ ) averaged over loci was computed using Fstat (Goudet 2001) and this software was also used to compute population level genetic differentiation ( $F_{ST}$ ) estimated by  $\theta$  (Weir and Cockerham 1984). Significance was assessed using 1000 permutations in Fstat. We used Bonferonni correction to adjust significance level for multiple tests (Rice 1989). As  $F_{ST}$  estimation is dependent on within and between population allele frequency (Meirmans and Hedrick 2011), we also computed  $D_{ST}$  (Jost 2008) using SMOGD (Crawford 2010) and 999 bootstrap replicates.  $D_{ST}$  is a standardized measure of population differentiation, not dependent on allele frequencies within and among populations. Population genetic structure was further investigated using the DAPC approach (Jombart et al. 2010) using the Adegenet package (Jombart 2008) implemented in R software. We used the Bayesian method implemented in GeneClass (Piry et al. 2004) to assign individuals to their potential river of origin. In a first approach, the population baseline was defined based on the river the fish were sampled in. The likelihood that sampled fish came from one of the river was tested with the Paetkau et al. (2004) algorithm using 100 000 simulated individuals. We finally quantified the relationship between genetic differentiation ( $F_{ST}$ ) and geographic distance by testing for a signal of isolation by distance (IBD) using a simple Mantel test in R with the package Vegan

(Oksanen et al. 2009). Population genetic distance was transformed following Rousset (1997) formulae ( $F_{ST} / (1-F_{ST})$ ).

### *Combining population genetic and otolith natal origins*

Otolith fingerprints were used to define a new population baseline by grouping individuals from the same natal river (and not from the same sampling location, as above). Measures of intra and inter-population differentiation ( $A_r$ ,  $H_o$ ,  $H_{nb}$ , pairwise  $F_{ST}$ ) were computed as above for these reconstructed populations. Differences in allelic richness and observed heterozygosity were tested using Wilcoxon paired signed-rank test (using values per locus) in R software. We then used this new baseline to assign all fish for which genetic data was available to potential groups using GeneClass (Piry et al. 2004). These results were then compared to those obtained previously.

## **Results**

### **Spatial differences in water signatures**

Mean water Sr:Ca, Ba:Ca and  $^{87}\text{Sr}:^{86}\text{Sr}$  ratios were significantly different among sites (Bonferroni adjustment,  $P < 0.05$ ) (Fig. 2). Water  $^{87}\text{Sr}:^{86}\text{Sr}$  ratios separated three groups of rivers: Portuguese rivers (Lima, Mondego and Minho rivers) with elevated  $^{87}\text{Sr}:^{86}\text{Sr}$  ratios (range between  $0.72261 \pm 1.03\text{E-}04$  and  $0.71571 \pm 1.81\text{E-}04$ ), Normandy and Brittany's rivers (Vire, Blavet, Scorff, Aulne, Loire and Vilaine rivers) with lower ratios (range between  $0.71160 \pm 1.00\text{E-}04$ ,  $0.71429 \pm 1.01\text{E-}04$ ), and southern French rivers from Charente River to Nivelle River (range between  $0.70823 \pm 5.70\text{E-}05$  and  $0.71037 \pm 1.22\text{E-}04$ ). Ba:Ca and Sr:Ca ratios separated most of the rivers within the three groups with the exception of the Oloron Saison and Nive Rivers which had overlapping elemental and Sr isotopic ratios.

Canonical Discriminant Analysis (CDA) showed strong geographical separation of water samples based on the geochemical signatures in rivers (Fig. 3). The 95% confidence

ellipse around the mean value for each river suggested that the water chemistry did not change significantly over the course of the 2013 sampling season (from late May to September 2013). The first two canonical variates explained more than 80% of the variation within the data (Wilks' lambda = 0.419,  $P < 0.01$ ). Rivers were generally separated along the first canonical variate, with the exception of the Saison, the Nive, the Oloron Rivers, and also, the Vire and the Vilaine Rivers. Canonical structure coefficients indicated the relative importance of the three variables to the separation in geochemical signatures amongst the rivers (Table 1). The first canonical variate was primarily driven by Sr isotopes ratios. Sr:Ca ratios also contributed to separation among rivers on the first canonical variate. Loadings on the second canonical variate were dominated by variation in Sr:Ca ratios with smaller contributions from  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  ratios. Finally Ba:Ca ratios appeared to contribute moderately to signature separation along the first and second variates. Owing to differences in Sr:Ca, Ba:Ca and  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  ratios, the cross-validation classification accuracies of rivers based on their geochemical signatures ranged from 80% to 100% (not presented). Among the seventeen rivers, misclassifications occurred in three neighboring rivers (the Oloron, the Saison and the Nive Rivers) and also between the Vilaine and the Vire Rivers. Those results likely arose because of similarities in chemical signatures between rivers as described in the Figure 2.

### Determination of adult natal streams

For all samples tested, apparent convergence under the Gelman-Rubin diagnostic was achieved. The Bayesian hierarchical mixture model was used to analyze otolith microchemistry signatures and provided reliable estimates of the proportion of individuals that displayed natal site fidelity or reproduced outside of their watershed of origin, i.e., migrants (supplementary Table S2). Posterior conditional assignment probabilities (i.e. the probability of assignment to each natal river) were higher than 0.80 for 85% of fish, indicating that the majority of adults were assigned to a source river with high degrees of confidence. In



the present results, while most individuals seemed to return to their natal watersheds, their fidelity to the natal river within the watershed of origin appeared less precise. The majority of spawners captured in the Scorff River and the Garonne River originated from adjacent spawning tributaries (i.e., the Blavet River and the Dordogne River respectively). Seventeen adults were classified as “undetermined” in supplementary Table S2 indicating that those individuals represent heterogeneous signatures that did not match those from the water and juvenile reference data set. Some individuals captured in the Vilaine River ( $n = 3$ ), the Loire River ( $n = 3$ ), the Garonne River ( $n = 14$ ), the Saison River ( $n = 3$ ), the Lima River ( $n = 2$ ) and the Mondego River ( $n = 11$ ) strayed into non-natal rivers to spawn but originated from neighbouring watersheds (distance between natal and spawning rivers were  $< 300$  km) (Table 2). Some nonresident spawning adults captured in the Vire River ( $n = 34$ ), the Adour watershed ( $n = 2$ ), the Minho River ( $n = 1$ ) and the Mondego River ( $n = 4$ ) travelled long (300-700 km) and ultra-long distances ( $> 700$  km) between natal and spawning river. The Vire River, the Garonne River and the Saison River displayed immigrants originating from the South (Table 2). The Vilaine River, the Loire River, the Adour watershed and all Portuguese Rivers displayed immigrants originating from the North.

### Genetic analysis

A total of 287 individuals were genotyped at 13 loci (supplementary Table S3). Significant deviations from HW equilibrium (heterozygote deficiency) were found in the Garonne population both in 2012 and 2013 (supplementary Table S3). Further investigation showed no significant linkage disequilibrium in these populations. Allelic richness ranged from 3.96 to 5.56 and averaged  $H_{nb}$  and  $H_o$  values were 0.61 and 0.60 respectively. Global  $F_{ST}$  was 0.026 (95%Ci 0.019-0.033,  $P = 0.0001$ ). 10 out of the 45 pairwise  $F_{ST}$  comparison were non-significant with values ranging from -0.0004 to 0.0745 (supplementary Table S4). The Nivelle showed the highest genetic differentiation with other rivers. Genetic

differentiation was similar using  $D_{ST}$  (0.032) with pairwise values ranging from -0.0132 to 0.745. The DAPC analysis indicated the existence of 5 genetically distinct groups based on the Bayesian Information Criterion. However, these groups were widely admixed (results not shown). Further investigations showed a significant pattern of isolation by distance ( $r = 0.6359$ ,  $P = 0.001$ ). GeneClass analysis assigned 81 out of 287 individuals at a threshold greater than 90% (Table 3). Individual assignment to their river of sampling was low and also supported a weak population structure and thus high connectivity. In five cases (Aulne, Scorff, Nivelle, Blavet and Vire), the majority of individuals were assigned to their respective sampling river, whereas for the remaining rivers, only few individuals were assigned to their river of sampling.

#### **Coupling otolith microchemistry and genetic markers**

The new baseline as defined by otolith natal origins was composed of eight groups representing the major French drainages. These new groups displayed similar level of genetic diversity ( $A_r$ ,  $H_o$ ,  $H_{nb}$ ) when compared to populations defined according to their sampling site (supplementary Table S5). Levels of  $F_{IS}$  were also reduced (supplementary Table S5). Level of genetic differentiation was also similar (Table 4). When using the new baseline 17 other individuals were assigned to a potential group at a 90% probability (Table 5). According to the new baseline, a greater number of individuals were miss-assigned to other rivers than their sampling river except for the Aulne, Nivelle and Vire Rivers where high rate of self-assignment were already observed using the genetic baseline.

### **Discussion**

#### **Variations in water and otolith chemistry**

Our analyses of water chemistry of Allis shad natal rivers revealed significant spatial variation from northern France to Portugal. This observed variation among rivers likely arose

415 from contrasting bedrock geology among respective drainage basins ([www.brgm.fr](http://www.brgm.fr)).  
416 However, similarities in water chemical signatures were observed in the case of three  
417 neighboring rivers from southern France (the Oloron, the Saison and the Nive rivers) and two  
418 northern French Rivers (the Vilaine and the Vire rivers), making the discrimination of these  
419 rivers more difficult.

420 It is likely that the otolith signatures measured in the present study had enough  
421 stability through time to enable the correct inference from 2013 signatures of the natal origin  
422 of fish hatched five or six years before (2007-2008). While previous works have shown that  
423 otolith  $^{87}\text{Sr}$ : $^{86}\text{Sr}$  ratios remained very stable from year to year as reported for American shad  
424 (Walther and Thorrold 2009), interannual stability was not well established in the case of  
425 trace elements. Both intra-and interannual variability in otolith elemental fingerprints may  
426 potentially confound spatial variation (Gillanders 2002). Thus, pooling juvenile otoliths  
427 and/or water samples collected from several years was recommended to account for the range  
428 of element values likely to be found in the adult cohorts (Walther and Thorrold 2009). Owing  
429 to a lack of water chemistry data and no juveniles' collection over the past few years in each  
430 study river, we did not know if ambient concentrations of Sr and Ba have fluctuated over  
431 years. However, comparison between recent water chemistry (data presented in this paper)  
432 and previous sampling years in 2001 (Tomas et al. 2005), in 2010 to 2013 in the Garonne  
433 River, the Dordogne River (F. Daverat, unpublished data) and tributaries of the Adour  
434 watershed (Martin et al. 2013a) indicated minor variation in Sr and Ba concentrations. This  
435 variability was neglectable compared to strong spatial variation in water chemistry among  
436 rivers. Moreover, significant interannual variability in otolith element signature was less  
437 frequently reported in freshwater species than estuarine or marine species (Hamer et al. 2003;  
438 Rooker et al. 2003; Patterson et al. 2004; Bergenius et al. 2005; Patterson et al. 2005; Elsdon  
439 and Gillanders 2006; Martin et al. 2013a). It appeared that strontium isotopes in otoliths

mainly drove the allocation of natal origins of Allis shad spawners. To illustrate its importance we re-ran the Bayesian hierarchical mixture model without including  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratios. We found that posterior conditional assignment probabilities (i.e. the probability of assignment to each natal river) dropped from 0.80 for 85% of fish to 0.6 for 90% of fish. It indicated that the majority of adults were assigned to a natal river with low degrees of confidence when Sr isotopes ratios were excluded from the model. First, the  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratio varies among rivers according to underlying geology (Kennedy et al. 1997; Ingram and Weber 1999; Hobbs et al. 2005) and is temporally stable (Kennedy et al. 2000; Walther and Thorrold 2009; Walther and Limburg 2012). Second, Sr isotopes are not trophically fractionated and the  $^{87}\text{Sr}:$  $^{86}\text{Sr}$  ratios in otoliths closely match that of stream water (Kennedy et al. 2000; Martin et al. 2013a). These properties highlighted the utility of Sr isotope ratios to distinguish natal origins of Allis shad at a local level. Further, given its strong spatial variability, Sr:Ca was also a key ratio for discriminating among natal rivers. Otolith Sr:Ca reflect ambient water composition (Bath et al. 2000), particularly in freshwater systems (Wells et al. 2003). Sr provided another temporally stable signature (Martin et al. 2013a) as its concentration in water is primarily controlled by the weathering of different lithologies and related soils (Rondeau et al. 2005). Therefore, we assumed that the assignation of natal origin using the water and otolith signatures cannot be impaired by minor temporal variability of chemical signatures.

### **The value of a Bayesian framework**

Bayesian tools were especially useful to determine geographical origins of individuals as they can incorporate several sources of uncertainty in a single analysis (Pella and Masuda 2001; Munch and Clarke 2008; Smith and Campana 2010; Pflugeisen and Calder 2013). In our study, we used water composition and otolith juvenile composition as additional sources of information to build the Bayesian hierarchical mixture model. Based on the linear

relationship between water and otolith signatures, the model allowed us to predict otolith chemistry for rivers where juveniles were not available (Pflugeisen and Calder 2013). Our ability to classify natal origins of Allis shad (85% overall accuracy) compared reasonably well with other studies on freshwater natal habitats of anadromous species (Thorrold et al. 1998; Milton and Chenery 2001; Tomas et al. 2005). A key assumption for inferring natal origins by the use of classification-based methods is that the signatures of all potential spawning rivers are known (i.e., exhaustive sampling) (Neubauer et al. 2013). Although we did not sample all potential natal rivers across the distribution range of Allis shad, our sampling was representative of the main spawning sites and likely exhaustive. Unsourced rivers were limited and also exhibited low natural reproductive success for Allis shad (Baglinière et al. 2003). Thus, we did not use the Bayesian models described by Neubauer et al. (2013) which inferred dispersal from incomplete geochemical baselines, taking into account potential unsampled sources. Our modelling approach assumed that most potential natal rivers have been characterized and that the assignment of individuals to a finite set of sources was non erroneous. Of the 410 individuals analyzed, 96% were assigned to rivers with high probability and only 4% were undetermined indicating that those individuals represented heterogeneous signatures not well established in the database. Those adults of unknown origin were probably from rivers outside the database. Thus, immigration of individuals from those rivers was likely low. Nevertheless, the unidentified origins of those individuals remain to be determined by expanding this work in geographical coverage. Some secondary rivers could deserve to be included in the signature references set to strengthen the power of origin detection such as the Vienne, the Creuse and the Mayenne rivers in the Loire River Basin, the Trieux and the Elorn River in Brittany, and in a southerly direction the Sèvre Niortaise River.

#### **Patterns of straying and philopatry**

Our genetic and otolith-based analysis suggested different patterns of philopatry and straying behavior between natal and spawning rivers for Allis shad. We found a significant proportion of individuals hatched and grown in a different watershed than the one they were collected from. Even if, the true probability of straying can only be estimated based on the population sizes in each location (Munch and Clarke 2008), that are not available for all the rivers we sampled, our study provided a valuable qualitative view on the dispersal and homing patterns of Allis shad. While *A. alosa* exhibited a high fidelity to the natal site on ecological timescales, as inferred from otolith microchemistry, this species on the other hand showed modest genetic differentiation between collection sites, and followed an isolation by distance pattern, which imply moderate but nevertheless sufficient straying to uniformize the Allis shad population genetic structure. Extensive gene flow between populations of Allis shad at this spatial scale were already suggested in French rivers using allozymes markers (Alexandrino et al. 2006) and mtDNA (Faria et al. 2012). Supporting this high rate of gene flow, genetic assignment success to natal sources remained relatively modest, even when using the otolith natal origin to define new baseline genetic groups. To date, the extent of intraspecific population structure is still unclear over the entire geographic range of Allis shad (Jolly et al. 2012).

In contrast, genetic analyses for *A. sapidissima* (Waters et al. 2000; Hasselman et al. 2013) and geochemical signatures from otolith (Walther et al. 2008) established a higher degree of homing behavior which induced significant but subtle population structuring. Conventional tagging (Melvin et al. 1986; Hendricks et al. 2002) also revealed that more than 90% of American shad spawners return to natal rivers. In our study, some rivers displayed more than 10% of spawners; up to 23% of spawners originated from other watersheds. Such a proportion of immigrants from neighbouring sites may explain the weaker genetic differences between discrete spawning populations in *Alosa alosa* (Wright 1950) as compared to *Alosa*

514 *sapidissima*. In comparison, anadromous salmonids tend to show a greater proportion of  
515 homing behavior than *Alosa alosa* (Jonsson and Jonsson 2011).

516 Allis shad site fidelity was not spatially precise, as adults returned to spawn within the  
517 river basins of origin rather within the river of origin in the Garonne-Dordogne Basin and  
518 Blavet-Scorff Basin. This suggests again, that the imprinting and homing behavior was not as  
519 strong in *Alosa alosa* as in other species of anadromous fish such as sockeye salmon (Stewart  
520 et al. 2003; Quinn et al. 2006; Quinn et al. 2012). Nevertheless, several authors showed that  
521 straying is not spatially random and fish are more likely to stray in adjacent watersheds (with  
522 similar environmental settings) than to enter more distant river drainages (Candy and  
523 Beacham 2000; Correa and Gross 2008; Hamann and Kennedy 2012).

524 While within our sample, straying occurred more frequently between neighbouring  
525 river basins, the lack of genetic structure but significant pattern of isolation by distance along  
526 the whole Atlantic coast (at the exception of Nivelle river) also supported the evidence of a  
527 significant flow of strayers between river basins in vicinity or at longer distances (two  
528 individuals captured in the Adour watershed seemed to originate from the Aulne River in  
529 Brittany located 600 km northward). However, if long distance straying is probable it seems  
530 not frequent, as we found that five individuals moved from France (natal area) to Portugal  
531 (spawning destination), and three individuals from Blavet toward the Vire River. The low  
532 probability of long distance straying between Portugal and northern populations was  
533 supported by the significant genetic differentiation of Portuguese and French populations  
534 (Alexandrino et al. 2006) and the lack of observations of abnormally large bodied shad  
535 spawners in the northern part of our study. As a matter of fact, Portuguese shad spawners  
536 compared to other populations, achieve a larger size at reproduction, which was attributed to  
537 the location of their marine nursery in an upwelling zone (Lassalle et al. 2008b). Furthermore,  
538 the IBD patterns suggest that long distance dispersal is limited and is probably stronger

between adjacent rivers. However, even a small number of migrants per generation may be enough to homogenize allele frequencies between populations (Allendorf 1983) and the impact of these few migrating individuals should not be neglected.

As the Vilaine and the Vire water signatures were very close to each other, the assignation of the Vire spawners as Vilaine River origin ( $n = 31$ ) can be doubtful. Hence, results were interpreted with caution concerning the reallocation of individuals sampled in the Vire River which originated from the Vilaine River. The confusion between the Vilaine and Vire otolith chemical signatures did not allow drawing definite conclusions on the intensity of flow of Allis shad straying from southern Brittany out to the Manche area, even though few spawners originating from Blavet were found in the Vire River.

Whether straying occurs in the early stage of sea nursery phase or when the growth is completed at sea remains unclear. Allis shad were found to inhabit shallow waters ( $< 150$  m) and aggregate at low-salinity areas of coastal regions (river mouths of most important watersheds) (Taverny and Elie 2001). A later study, Trancart et al. (2014), demonstrated that Allis shad populations remained in the same geographic area throughout the year, but the study lacked to be extended to the southern part of the Atlantic coastal area below the Gironde estuary mouth. Lassalle et al. (2008b) also concluded that the marine growing areas are probably located near the natal river. Straying would then occur between the coastal nursery grounds corresponding to different river basins origins. The genetic differences observed between the French and Portuguese Allis shad populations may be due to the lower ability of shad to cross over large coastal distances and deep sea bathymetric barriers such as Capbreton canyon (Bay of Biscay, SW France) (Carlsson et al. 2011). *Salmo salar* disperse as a spawner from a single marine nursery located offshore and in consequence, strayers may originate from rivers far away (Perrier et al. 2011). It seems that a precise homing behavior especially in salmonids that undertake long distance migrations, made the successful evolution of



anadromy possible, by maximizing returns of individuals back to suitable habitats where they were spawned (McDowall 2001). As marine growing areas of Allis shad seem to be located near their natal rivers, the ability to home could be an evolutionary feature of clupeid biology that is not as critical as compared to salmonid species. Moreover, straying may be more frequent for Allis shad compared to other anadromous fish through inadequate juvenile imprinting due to the shorter freshwater growing period (only two months). Twaité shad also has a short freshwater growing period but genetic studies provided strong evidences for a greater segregation of populations compared to Allis shad (Jolly et al. 2012). The marine distribution of *Alosa fallax* is more coastal (water depths < 50 m) than for *Alosa alosa* (Taverny and Elie 2001) and the species is clumped in aggregations around the major catchments for reproduction, which could be consistent with a higher fidelity to spawning grounds and/or limited migration during the adult phase.

In a context of global change, straying would be a key mechanism to drive dispersal and allow resilience of Allis shad populations. Such behavioral diversity may be critical to long-term persistence of the species by linking local populations that independently would be vulnerable to extinction or by refunding those that do go extinct (Schlosser and Angermeier 1995; Rieman and Dunham 2000; Rieman and Allendorf 2001). The recovery of a depleted population relies on the size and proximity of neighboring populations, as well as on the proportion of immigrants that contribute to the depleted population (Hamann and Kennedy 2012; Keefer and Caudill 2014). Our result suggested that a river may act as a source of spawners when the other rivers had experienced a lower recruitment success. The disproportionate contribution of the Dordogne River and the Blavet River to the adult population entering both river basins may be due to fish in these habitats having more successful recruitment to the adult population. Similarly, Minho River provided a source of migrants into neighbouring rivers (i.e., the Lima and Mondego Rivers). Whether the provision

589 of spawners may be sufficient to restore a population remains doubtful when habitats may not  
 590 be suitable for juveniles. The Minho River is the only Portuguese River which has free access  
 591 to suitable spawning grounds for Allis shad since the first downstream barrier is located far  
 592 from river mouth (Mota and Antunes 2012). In the Lima and Mondego Rivers, dams blocked  
 593 access to historical spawning reaches, causing severe fish population declines. In the specific  
 594 case of Dordogne-Garonne river basin, within our entire sample, no spawner with a Garonne  
 595 otolith signature was identified for two consecutive years (2012 and 2013) suggesting either  
 596 that the survival of juveniles or the spawning activity in the Garonne river five years earlier  
 597 might have been critical. Despite the immigration of Dordogne and other river spawners into  
 598 the Garonne River, no sign of population recovery could be observed. A significant  $F_{IS}$  value  
 599 was observed for the Garonne population in both 2012 and 2013, suggesting a Wahlund  
 600 effect. This effect suggests that the depleted populations may not be sufficiently supported by  
 601 straying itself. The depletion of Allis shad populations along the Atlantic coast may not allow  
 602 the recovery of specific stocks if an Allee effect is confirmed for this species (Rougier et al.  
 603 2012).

604       Lassalle et al. (2009), showed that under global warming the environment would be  
 605 more favorable for shad towards the north of its distribution area. Our results did not conclude  
 606 on the predominance of a northward dispersal of Allis shad with more strayers going  
 607 northward. However, the recent increase of spawners abundance in the Vire River at the  
 608 northern part of our sampling, may be the combined result of a larger abundance of strayers  
 609 and a more successful recruitment. Thus the analysis of straying data together with data on the  
 610 relative abundance of river populations would bring insight on a northward predominance of  
 611 dispersal. This suggests a possibility for the stock to regain its historical distribution area by  
 612 recolonizing rivers of the North of Europe, such as the Seine or the Rhine, providing that their  
 613 longitudinal connectivity and spawning habitats are accessible, suitable and/or restored.

## Acknowledgements

Research was funded by the IRSTEA (Institut national de Recherche en Sciences et Technologies pour l'Environnement et l'Agriculture). Femtosecond laser ablation system and HR-ICP-MS were funded by Region Aquitaine and Europe. We are grateful to various workers who assisted with sample collection (Fédération de Pêche, Syndicat Mixte Aménagement Tourisme Aulne et Hyères: SMATAH, Centre Régional Expérimentation et Application Aquacole: CREAA, Association Bretagne Grands Migrateurs, Association Migrateur Garonne Dordogne: MIGADO). Part of the samples were collected thanks to a research project funded by the French Ministry of Environment and led by the French Museum of Natural History (MNHN) on the efficiency of Marine Protected Areas for Shads and Lampreys. We thank fishermen P. Boisneau and D. Macé for collection of adult fish in the Loire and Vilaine Rivers respectively. We would like to thank P. Lambert from the IRSTEA institute for its help in implementing the results and discussion of the present work. Many thanks to Anne-Laure Besnard for her help in molecular biology analyses. We finally thank H. Tabouret and A. Holub for their help in water sample preparation and analysis.

## References

- Alexandrino, P., Faria, R., Linhares, D., Castro, F., Le Corre, M., Sabatié, R., Baglinière, J.L., and Weiss, S. 2006. Interspecific differentiation and intraspecific substructure in two closely related clupeids with extensive hybridization, *Alosa alosa* and *Alosa fallax*. *Journal of Fish Biology* **69**(SUPPL. B): 242-259.
- Allègre, C.J., Louvat, P., Gaillardet, J., Meynadier, L., Rad, S., and Capmas, F. 2010. The fundamental role of island arc weathering in the oceanic Sr isotope budget. *Earth and Planetary Science Letters* **292**(1-2): 51-56.
- Allendorf, F.W. 1983. Isolation, gene flow, and genetic differentiation among populations. *Genetics and conservation* **18**(3): 51-65.
- Baglinière, J.L., Sabatié, R., Rochard, E., Alexandrino, P., and Aprahamian, M.W. 2003. The allis shad *Alosa alosa*: biology, ecology, range, and status of populations. *American Fisheries Society Symposium* **35**: 85-102.
- Banks, M.A., Rashbrook, V.K., Calavetta, M.J., Dean, C.A., and Hedgecock, D. 2000. Analysis of microsatellite DNA resolves genetic structure and diversity of chinook salmon (*Oncorhynchus tshawytscha*) in California's Central Valley. *Canadian Journal of Fisheries and Aquatic Sciences* **57**(5): 915-927.
- Barats, A., Amouroux, D., Pecheyran, C., Chauvaud, L., and Donard, O.F.X. 2007. High-Frequency Archives of Manganese Inputs To Coastal Waters (Bay of Seine, France) Resolved by the LA ICPMS Analysis of Calcitic Growth Layers along Scallop Shells (*Pecten maximus*). *Environmental Science & Technology* **42**(1): 86-92.
- Barnett-Johnson, R., Pearson, T.E., Ramos, F.C., Grimes, C.B., and Bruce MacFarlane, R. 2008. Tracking natal origins of salmon using isotopes, otoliths, and landscape geology. *Limnology and Oceanography* **53**(4): 1633-1642.

- 654 Barnett-Johnson, R., Teel, D.J., and Casillas, E. 2010. Genetic and otolith isotopic markers  
655 identify salmon populations in the Columbia River at broad and fine geographic scales.  
656 Environmental Biology of Fishes **89**(3): 533-546.
- 657 Bath, G.E., Thorrold, S.R., Jones, C.M., Campana, S.E., McLaren, J.W., and Lam, J.W.H.  
658 2000. Strontium and barium uptake in aragonitic otoliths of marine fish. Geochimica et  
659 Cosmochimica Acta **64**(10): 1705-1714.
- 660 Belkhir, K., Borsa, P., Chikhi, L., Raufaste, N., and Bonhomme, F. 2004. GENETIX 4.05,  
661 logiciel sous Windows TM pour la génétique des populations. version 4.05. Laboratoire  
662 Génome, Populations, Interactions, CNRS UMR 5171, Université de Montpellier II,  
663 Montpellier. Available from <http://www.genetix.univ-montp2.fr/genetix/genetix.htm>.
- 664 Bentzen, P., Taggart, C.T., Ruzzante, D.E., and Cook, D. 1996. Microsatellite polymorphism  
665 and the population structure of Atlantic cod (*Gadus morhua*) in the northwest Atlantic.  
666 Canadian Journal of Fisheries and Aquatic Sciences **53**(12): 2706-2721.
- 667 Bergenius, M.A.J., Mapstone, B.D., Begg, G.A., and Murchie, C.D. 2005. The use of otolith  
668 chemistry to determine stock structure of three epinepheline serranid coral reef fishes on  
669 the Great Barrier Reef, Australia. Fisheries Research **72**(2-3): 253-270.
- 670 Blum, J.D., Taliaferro, E.H., Weisse, M.T., and Holmes, R.T. 2000. Changes in Sr/Ca, Ba/Ca  
671 and <sup>87</sup>Sr/<sup>86</sup>Sr ratios between trophic levels in two forest ecosystems in the northeastern  
672 U.S.A. Biogeochemistry **49**(1): 87-101.
- 673 Bradbury, I.R., Campana, S.E., and Bentzen, P. 2008. Estimating contemporary early life-  
674 history dispersal in an estuarine fish: Integrating molecular and otolith elemental  
675 approaches. Molecular Ecology **17**(6): 1438-1450.
- 676 Bradbury, I.R., Hamilton, L.C., Robertson, M.J., Bourgeois, C.E., Mansour, A., and  
677 Dempson, J.B. 2014. Landscape structure and climatic variation determine Atlantic salmon

- 678 genetic connectivity in the Northwest Atlantic. Canadian Journal of Fisheries and Aquatic  
679 Sciences **71**(2): 246-258.
- 680 Brown, R.J., and Severin, K.P. 2009. Otolith chemistry analyses indicate that water Sr:Ca is  
681 the primary factor influencing otolith Sr:Ca for freshwater and diadromous fish but not for  
682 marine fish. Canadian Journal of Fisheries and Aquatic Sciences **66**(10): 1790-1808.
- 683 Campana, S.E. 1999. Chemistry and composition of fish otoliths: pathways, mechanisms and  
684 applications. Marine Ecology-Progress Series **188**: 263-297.
- 685 Candy, J.R., and Beacham, T.D. 2000. Patterns of homing and straying in southern British  
686 Columbia coded-wire tagged chinook salmon (*Oncorhynchus tshawytscha*) populations.  
687 Fisheries Research **47**(1): 41-56.
- 688 Carlsson, J., Shephard, S., Coughlan, J., N. Trueman, C., Rogan, E., and Cross, T.F. 2011.  
689 Fine-scale population structure in a deep-sea teleost (orange roughy, *Hoplostethus*  
690 *atlanticus*). Deep-Sea Research Part I: Oceanographic Research Papers **58**(6): 627-636.
- 691 Collins, S.M., Bickford, N., McIntyre, P.B., Coulon, A., Ulseth, A.J., Taphorn, D.C., and  
692 Flecker, A.S. 2013. Population structure of a neotropical migratory fish: Contrasting  
693 perspectives from genetics and Otolith microchemistry. Transactions of the American  
694 Fisheries Society **142**(5): 1192-1201.
- 695 Correa, C., and Gross, M.R. 2008. Chinook salmon invade southern South America.  
696 Biological Invasions **10**(5): 615-639.
- 697 Crawford, N.G. 2010. SMOGD: software for the measurement of genetic diversity. Molecular  
698 Ecology Resources **10**(3): 556-557.
- 699 De Groot, S.J. 2002. A review of the past and present status of anadromous fish species in the  
700 Netherlands: Is restocking the Rhine feasible? Hydrobiologia **478**: 205-218.

- 701 Denwood, M. 2013. runjags: An R Package Providing Interface Utilities, Distributed  
702 Computing Methods and Additional Distributions For MCMC Models in JAGS. Journal of  
703 Statistical Software.
- 704 Dorval, E., Jones, C.M., Hannigan, R., and Van Montfrans, J. 2007. Relating otolith  
705 chemistry to surface water chemistry in a coastal plain estuary. Canadian Journal of  
706 Fisheries and Aquatic Sciences **64**(3): 411-424.
- 707 Elsdon, T.S., and Gillanders, B.M. 2006. Temporal variability in strontium, calcium, barium,  
708 and manganese in estuaries: Implications for reconstructing environmental histories of fish  
709 from chemicals in calcified structures. Estuarine, Coastal and Shelf Science **66**(1-2): 147-  
710 156.
- 711 Elsdon, T.S., Wells, B.K., Campana, S.E., Gillanders, B.M., Jones, C.M., Limburg, K.E.,  
712 Secor, D.E., Thorrold, S.R., and Walther, B.D. 2008. Otolith chemistry to describe  
713 movements and life-history measurements of fishes: hypotheses, assumptions, limitations,  
714 and inferences using five methods. Oceanography and Marine Biology: an Annual Review  
715 **46**: 297-330.
- 716 Estoup, A., Largiader, C.R., Perrot, E., and Chourrout, D. 1996. Rapid one-tube DNA  
717 extraction for reliable PCR detection of fish polymorphic markers and transgenes.  
718 Molecular Marine Biology and Biotechnology **5**(4): 295-298.
- 719 Faria, R., Weiss, S., and Alexandrino, P. 2012. Comparative phylogeography and  
720 demographic history of European shads (*Alosa alosa* and *A. fallax*) inferred from  
721 mitochondrial DNA. BMC Evolutionary Biology **12**(1).
- 722 Farrell, J., and Campana, S.E. 1996. Regulation of calcium and strontium deposition on the  
723 otoliths of juvenile tilapia, *Oreochromis niloticus*. Comparative Biochemistry and  
724 Physiology - A Physiology **115**(2): 103-109.

- 725 Feyrer, F., Hobbs, J., Baerwald, M., Sommer, T., Yin, Q.Z., Clark, K., May, B., and Bennett,  
 726 W. 2007. Otolith microchemistry provides information complementary to microsatellite  
 727 DNA for a migratory fish. Transactions of the American Fisheries Society **136**(2): 469-  
 728 476.
- 729 Fromentin, J.M., Ernande, B., Fablet, R., and De Pontual, H. 2009. Importance and future of  
 730 individual markers for the ecosystem approach to fisheries. Aquatic Living Resources  
 731 **22**(4): 395-408.
- 732 Gelman, A., and Rubin, D.B. 1992. Inference from iterative simulation using multiple  
 733 sequences. Statistical science: 457-472.
- 734 Gillanders, B.M. 2002. Temporal and spatial variability in elemental composition of otoliths:  
 735 Implications for determining stock identity and connectivity of populations. Canadian  
 736 Journal of Fisheries and Aquatic Sciences **59**(4): 669-679.
- 737 Goudet, J. 2001. Fstat, a program to estimate and test gene diversities and fixation indices.  
 738 Version 2.9.3. Available from <http://www2.unil.ch/popgen/softwares/fstat.htm> .
- 739 Hamann, E.J., and Kennedy, B.P. 2012. Juvenile dispersal affects straying behaviors of adults  
 740 in a migratory population. Ecology **93**(4): 733-740.
- 741 Hamer, P.A., Jenkins, G.P., and Gillanders, B.M. 2003. Otolith chemistry of juvenile snapper  
 742 *Pagrus auratus* in Victorian waters: Natural chemical tags and their temporal variation.  
 743 Marine Ecology Progress Series **263**: 261-273.
- 744 Hasselman, D.J., Ricard, D., and Bentzen, P. 2013. Genetic diversity and differentiation in a  
 745 wide ranging anadromous fish, American shad (*Alosa sapidissima*), is correlated with  
 746 latitude. Molecular Ecology **22**(6): 1558-1573.
- 747 Hendricks, M.L., Hoopes, R.L., Arnold, D.A., and Kaufmann, M.L. 2002. Homing of  
 748 hatchery-reared American shad to the Lehigh River, a tributary to the Delaware River.  
 749 North American Journal of Fisheries Management **22**(1): 243-248.



- 750 Hendry, A.P., Castric, V., Kinnison, M.T., and Quinn, T.P. 2004. The evolution of philopatry  
751 and dispersal: homing versus straying in salmonids. *In* Evolution illuminated: salmon and  
752 their relatives. *Edited by* A.P. Hendry and S.C. Stearns. Oxford, New York. pp. 52-91.
- 753 Hobbs, J.A., Yin, Q.Z., Burton, J., and Bennett, W.A. 2005. Retrospective determination of  
754 natal habitats for an estuarine fish with otolith strontium isotope ratios. *Marine and*  
755 *Freshwater Research* **56**(5): 655-660.
- 756 Hughes, J.M., Schmidt, D.J., Macdonald, J.I., Huey, J.A., and Crook, D.A. 2014. Low  
757 interbasin connectivity in a facultatively diadromous fish: Evidence from genetics and  
758 otolith chemistry. *Molecular Ecology* **23**(5): 1000-1013.
- 759 Humston, R., Priest, B.M., Hamilton, W.C., and Bugas, P.E. 2010. Dispersal between  
760 tributary and main-stem rivers by juvenile smallmouth bass evaluated using otolith  
761 microchemistry. *Transactions of the American Fisheries Society* **139**(1): 171-184.
- 762 Ingram, B.L., and Weber, P.K. 1999. Salmon origin in California's Sacramento-San Joaquin  
763 river system as determined by otolith strontium isotopic composition. *Geology* **27**(9): 851-  
764 854.
- 765 Jolly, M.T., Aprahamian, M.W., Hawkins, S.J., Henderson, P.A., Hillman, R., O'Maoiléidigh,  
766 N., Maitland, P.S., Piper, R., and Genner, M.J. 2012. Population genetic structure of  
767 protected allis shad (*Alosa alosa*) and twaite shad (*Alosa fallax*). *Marine Biology* **159**(3):  
768 675-687.
- 769 Jombart, T. 2008. Adegnet: A R package for the multivariate analysis of genetic markers.  
770 *Bioinformatics* **24**(11): 1403-1405.
- 771 Jombart, T., Devillard, S., and Balloux, F. 2010. Discriminant analysis of principal  
772 components: A new method for the analysis of genetically structured populations. *BMC*  
773 *Genetics* **11**.

- 774 Jonsson, B., and Jonsson, N. 2011. Ecology of Atlantic salmon and brown trout: habitat as a  
775 template for life histories. Springer.
- 776 Jost, L. 2008. GST and its relatives do not measure differentiation. *Molecular Ecology* **17**(18):  
777 4015-4026.
- 778 Kalish, J.M. 1990. Use of otolith microchemistry to distinguish the progeny of sympatric  
779 anadromous and non-anadromous salmonids. *Fishery Bulletin* **88**(4): 657-666.
- 780 Keefer, M.L., and Caudill, C.C. 2014. Homing and straying by anadromous salmonids: A  
781 review of mechanisms and rates. *Reviews in Fish Biology and Fisheries* **24**(1): 333-368.
- 782 Kennedy, B.P., Folt, C.L., Blum, J.D., and Chamberlain, C.P. 1997. Natural isotope markers  
783 in salmon. *Nature* **387**(6635): 766-767.
- 784 Kennedy, B.P., Blum, J.D., Folt, C.L., and Nislow, K.H. 2000. Using natural strontium  
785 isotopic signatures as fish markers: Methodology and application. *Canadian Journal of*  
786 *Fisheries and Aquatic Sciences* **57**(11): 2280-2292.
- 787 Lassalle, G., Béguer, M., Beaulaton, L., and Rochard, E. 2008a. Diadromous fish  
788 conservation plans need to consider global warming issues: an approach using  
789 biogeographical models. *Biological Conservation* **141**(4): 1105-1118.
- 790 Lassalle, G., Trancart, T., Lambert, P., and Rochard, E. 2008b. Latitudinal variations in age  
791 and size at maturity among allis shad *Alosa alosa* populations. *Journal of Fish Biology*  
792 **73**(7): 1799-1809.
- 793 Lassalle, G., Crouzet, P., and Rochard, E. 2009. Modelling the current distribution of  
794 European diadromous fishes: An approach integrating regional anthropogenic pressures.  
795 *Freshwater Biology* **54**(3): 587-606.
- 796 Limburg, K.E., and Waldman, J.R. 2009. Dramatic declines in north Atlantic diadromous  
797 fishes. *BioScience* **59**(11): 955-965.

- 798 Lochet, A., Jatteau, P., Tomàs, J., and Rochard, E. 2008. Retrospective approach to  
799 investigating the early life history of a diadromous fish: Allis shad *Alosa alosa* (L.) in the  
800 Gironde-Garonne-Dordogne watershed. *Journal of Fish Biology* **72**(4): 946-960.
- 801 Lochet, A., Boutry, S., and Rochard, E. 2009. Estuarine phase during seaward migration for  
802 allis shad *Alosa alosa* and twaite shad *Alosa fallax* future spawners. *Ecology of Freshwater*  
803 *Fish*.
- 804 Martin, J., Bareille, G., Berail, S., Pecheyran, C., Daverat, F., Bru, N., Tabouret, H., and  
805 Donard, O. 2013a. Spatial and temporal variations in otolith chemistry and relationships  
806 with water chemistry: A useful tool to distinguish Atlantic salmon *Salmo salar* parr from  
807 different natal streams. *Journal of Fish Biology* **82**(5): 1556-1581.
- 808 Martin, J., Bareille, G., Berail, S., Pecheyran, C., Gueraud, F., Lange, F., Daverat, F., Bru, N.,  
809 Beall, E., Barracou, D., and Donard, O. 2013b. Persistence of a southern Atlantic salmon  
810 population: Diversity of natal origins from otolith elemental and Sr isotopic signatures.  
811 *Canadian Journal of Fisheries and Aquatic Sciences* **70**(2): 182-197.
- 812 McDowall, R.M. 2001. Anadromy and homing: two life-history traits with adaptive synergies  
813 in salmonid fishes? *Fish and Fisheries* **2**: 78-85.
- 814 McDowall, R.M. 2008. Diadromy, history and ecology: A question of scale. *Hydrobiologia*  
815 **602**(1): 5-14.
- 816 McGarigal, K., Cushman, S., and Stafford, S.G. 2000. Multivariate statistics for wildlife and  
817 ecology research.
- 818 Meirmans, P.G., and Hedrick, P.W. 2011. Assessing population structure: FST and related  
819 measures. *Molecular Ecology Resources* **11**(1): 5-18.
- 820 Melvin, G.D., Dadswell, M.J., and Martin, J.D. 1986. Fidelity of American shad, *Alosa*  
821 *sapidissima* (Clupeidae), to its river of previous spawning. *Canadian Journal of Fisheries*  
822 *and Aquatic Sciences* **43**(3): 640-646.

- 823 Milton, D.A., and Chenery, S.R. 2001. Sources and uptake of trace metals in otoliths of  
 824 juvenile barramundi (*Lates calcarifer*). Journal of Experimental Marine Biology and  
 825 Ecology **264**(1): 47-65.
- 826 Mota, M., and Antunes, C. 2012. A preliminary characterisation of the habitat use and feeding  
 827 of Allis shad (*Alosa alosa*) juveniles in the Minho River tidal freshwater wetlands.  
 828 Limnetica **31**(1): 165-172.
- 829 Munch, S.B., and Clarke, L.M. 2008. A Bayesian approach to identifying mixtures from  
 830 otolith chemistry data. Canadian Journal of Fisheries and Aquatic Sciences **65**(12): 2742-  
 831 2751.
- 832 Neubauer, P., Shima, J.S., and Swearer, S.E. 2013. Inferring dispersal and migrations from  
 833 incomplete geochemical baselines: Analysis of population structure using Bayesian infinite  
 834 mixture models. Methods in Ecology and Evolution **4**(9): 836-845.
- 835 O'Connell, M., and Wright, J.M. 1997. Microsatellite DNA in fishes. Reviews in Fish Biology  
 836 and Fisheries **7**(3): 331-363.
- 837 Oksanen, J., Kindt, R., Legendre, P., O'Hara, B., Simpson, G.L., Solymos, P., Stevens,  
 838 M.H.H., and Wagner, H. 2009. Vegan: community ecology package. R package version  
 839 1.15-4.
- 840 Paetkau, D., Slade, R., Burden, M., and Estoup, A. 2004. Genetic assignment methods for the  
 841 direct, real-time estimation of migration rate: A simulation-based exploration of accuracy  
 842 and power. Molecular Ecology **13**(1): 55-65.
- 843 Patterson, H.M., Kingsford, M.J., and McCulloch, M.T. 2004. The influence of oceanic and  
 844 lagoonal plume waters on otolith chemistry. Canadian Journal of Fisheries and Aquatic  
 845 Sciences **61**(6): 898-904.
- 846 Patterson, H.M., Kingsford, M.J., and McCulloch, M.T. 2005. Resolution of the early life  
 847 history of a reef fish using otolith chemistry. Coral Reefs **24**(2): 222-229.

- 848 Pella, J., and Masuda, M. 2001. Bayesian methods for analysis of stock mixtures from genetic  
849 characters. *Fishery Bulletin* **99**(1): 151-167.
- 850 Perrier, C., Daverat, F., Evanno, G., Pécheyran, C., Bagliniere, J.L., and Roussel, J.M. 2011.  
851 Coupling genetic and otolith trace element analyses to identify river-born fish with  
852 hatchery pedigrees in stocked Atlantic salmon (*Salmo salar*) populations. *Canadian Journal*  
853 *of Fisheries and Aquatic Sciences* **68**(6): 977-987.
- 854 Pflugeisen, B.M., and Calder, C.A. 2013. Bayesian hierarchical mixture models for otolith  
855 microchemistry analysis. *Environmental and Ecological Statistics* **20**(2): 179-190.
- 856 Piry, S., Alapetite, A., Cornuet, J.M., Paetkau, D., Baudouin, L., and Estoup, A. 2004.  
857 GENECLASS2: A software for genetic assignment and first-generation migrant detection.  
858 *Journal of Heredity* **95**(6): 536-539.
- 859 Plummer, M. 2003. JAGS: a program for analysis of Bayesian graphical models using Gibbs  
860 sampling. *Proceedings of the 3rd International workshop on distributed statistical*  
861 *computing*.
- 862 Plummer, M., Best, N., Cowles, K., and Vines, K. 2006. CODA: Convergence diagnosis and  
863 output analysis for MCMC. *R news* **6**(1): 7-11.
- 864 Quinn, T.P., Stewart, I.J., and Boatright, C.P. 2006. Experimental evidence of homing to site  
865 of incubation by mature sockeye salmon, *Oncorhynchus nerka*. *Animal Behaviour* **72**(4):  
866 941-949.
- 867 Quinn, T.P., Rich Jr, H.B., Gosse, D., and Schtickzelle, N. 2012. Population dynamics and  
868 asynchrony at fine spatial scales: A case history of sockeye salmon (*Oncorhynchus nerka*)  
869 population structure in Alaska, USA. *Canadian Journal of Fisheries and Aquatic Sciences*  
870 **69**(2): 297-306.

- 871 R. Development Core Team. 2013. R: A language and environment for statistical computing.  
 872 R Foundation for Statistical Computing, Vienna, Austria. ISBN 3-900051-07-0, URL  
 873 <http://www.R-project.org>. R Foundation for Statistical Computing, Vienna, Austria.
- 874 Rice, W.R. 1989. Analyzing tables of statistical tests. *Evolution*: 223-225.
- 875 Rieman, B.E., Myers, D.L., and Nielsen, R.L. 1994. Use of Otolith Microchemistry to  
 876 Discriminate *Oncorhynchus-Nerka* of Resident and Anadromous Origin. *Canadian Journal*  
 877 *of Fisheries and Aquatic Sciences* **51**(1): 68-77.
- 878 Rieman, B.E., and Dunham, J.B. 2000. Metapopulations and salmonids: A synthesis of life  
 879 history patterns and empirical observations. *Ecology of Freshwater Fish* **9**(1-2): 51-64.
- 880 Rieman, B.E., and Allendorf, F.W. 2001. Effective population size and genetic conservation  
 881 criteria for bull trout. *North American Journal of Fisheries Management* **21**(4): 756-764.
- 882 Rondeau, B., Cossa, D., Gagnon, P., Pham, T.T., and Surette, C. 2005. Hydrological and  
 883 biogeochemical dynamics of the minor and trace elements in the St. Lawrence River.  
 884 *Applied Geochemistry* **20**(7): 1391-1408.
- 885 Rooker, J.R., Secor, D.H., Zdanowicz, V.S., De Metrio, G., and Relini, L.O. 2003.  
 886 Identification of Atlantic bluefin tuna (*Thunnus thynnus*) stocks from putative nurseries  
 887 using otolith chemistry. *Fisheries Oceanography* **12**(2): 75-84.
- 888 Rougemont, Q., Besnard, A.-L., Baglinière, J.-L., and Launey, S. 2014. Characterization of  
 889 thirteen new microsatellite markers for allis shad (*Alosa alosa*) and twaite shad (*Alosa*  
 890 *fallax*). *Conservation Genetics Resources*: 1-3.
- 891 Rougier, T., Lambert, P., Drouineau, H., Girardin, M., Castelnaud, G., Carry, L., Aprahamian,  
 892 M., Rivot, E., and Rochard, E. 2012. Collapse of allis shad, *Alosa alosa*, in the Gironde  
 893 system (southwest France): Environmental change, fishing mortality, or Allee effect? *ICES*  
 894 *Journal of Marine Science* **69**(10): 1802-1811.

- 895 Rousset, F. 1997. Genetic differentiation and estimation of gene flow from F-statistics under  
896 isolation by distance. *Genetics* **145**(4): 1219-1228.
- 897 Rousset, F. 2008. genepop'007: a complete re-implementation of the genepop software for  
898 Windows and Linux. *Molecular Ecology Resources* **8**(1): 103-106.
- 899 Schlosser, I.J., and Angermeier, P. 1995. Spatial variation in demographic processes of lotic  
900 fishes: conceptual models, empirical evidence, and implications for conservation.  
901 American Fisheries Society symposium.
- 902 Selkoe, K.A., and Toonen, R.J. 2006. Microsatellites for ecologists: A practical guide to using  
903 and evaluating microsatellite markers. *Ecology Letters* **9**(5): 615-629.
- 904 Selkoe, K.A., Henzler, C.M., and Gaines, S.D. 2008. Seascape genetics and the spatial  
905 ecology of marine populations. *Fish and Fisheries* **9**(4): 363-377.
- 906 Smith, S.J., and Campana, S.E. 2010. Integrated stock mixture analysis for continuous and  
907 categorical data, with application to genetic-otolith combinations. *Canadian Journal of*  
908 *Fisheries and Aquatic Sciences* **67**(10): 1533-1548.
- 909 Spidle, A.P., Kalinowski, S.T., Lubinski, B.A., Perkins, D.L., Beland, K.F., Kocik, J.F., and  
910 King, T.L. 2003. Population structure of Atlantic salmon in Maine with reference to  
911 populations from Atlantic Canada. *Transactions of the American Fisheries Society* **132**(2):  
912 196-209.
- 913 Stewart, I.J., Quinn, T.P., and Bentzen, P. 2003. Evidence for fine-scale natal homing among  
914 island beach spawning sockeye salmon, *Oncorhynchus nerka*. *Environmental Biology of*  
915 *Fishes* **67**(1): 77-85.
- 916 Taverny, C., and Elie, P. 2001. Répartition spatio-temporelle de la grande alose *Alosa alosa*  
917 (Linné, 1766) et de l'alse feinte *Alosa fallax* (Lacépède, 1803) dans le golfe de Gascogne.  
918 *Bulletin Français de la Pêche et de la Pisciculture* **362-363**: 803-821.

- 919 Thorrold, S.R., Jones, C.M., Campana, S.E., McLaren, J.W., and Lam, J.W.H. 1998. Trace  
 920 element signatures in otoliths record natal river of juvenile American shad (*Alosa*  
 921 *sapidissima*). Limnology and Oceanography **43**(8): 1826-1835.
- 922 Tomas, J., Augagneur, S., and Rochard, E. 2005. Discrimination of the natal origin of young-  
 923 of-the-year Allis shad (*Alosa alosa*) in the Garonne-Dordogne basin (south-west France)  
 924 using otolith chemistry. Ecology of Freshwater Fish **14**(2): 185-190.
- 925 Trancart, T., Rochette, S., Acou, A., Lasne, E., and Feunteun, E. 2014. Modeling marine shad  
 926 distribution using data from French bycatch fishery surveys. Marine Ecology Progress  
 927 Series **511**: 181-192.
- 928 Van Oosterhout, C., Hutchinson, W.F., Wills, D.P.M., and Shipley, P. 2004. MICRO-  
 929 CHECKER: Software for identifying and correcting genotyping errors in microsatellite  
 930 data. Molecular Ecology Notes **4**(3): 535-538.
- 931 Walther, B.D., and Thorrold, S.R. 2006. Water, not food, contributes the majority of  
 932 strontium and barium deposited in the otoliths of a marine fish. Marine Ecology Progress  
 933 Series **311**: 125-130.
- 934 Walther, B.D., Thorrold, S.R., and Olney, J.E. 2008. Geochemical signatures in otoliths  
 935 record natal origins of American shad. Transactions of the American Fisheries Society  
 936 **137**(1): 57-69.
- 937 Walther, B.D., and Thorrold, S.R. 2009. Inter-annual variability in isotope and elemental  
 938 ratios recorded in otoliths of an anadromous fish. Journal of Geochemical Exploration  
 939 **102**(3): 181-186.
- 940 Walther, B.D., and Limburg, K.E. 2012. The use of otolith chemistry to characterize  
 941 diadromous migrations. Journal of Fish Biology **81**(2): 796-825.



942 Waters, J.M., Epifanio, J.M., Gunter, T., and Brown, B.L. 2000. Homing behaviour facilitates  
943 subtle genetic differentiation among river populations of *Alosa sapidissima*: Microsatellites  
944 and mtDNA. Journal of Fish Biology **56**(3): 622-636.

945 Weir, B.S., and Cockerham, C.C. 1984. Estimating F-statistics for the analysis of population  
946 structure. evolution: 1358-1370.

947 Wells, B.K., Rieman, B.E., Clayton, J.L., Horan, D.L., and Jones, C.M. 2003. Relationships  
948 between water, otolith, and scale chemistries of westslope cutthroat trout from the Coeur  
949 d'Alene River, Idaho: The potential application of hard-part chemistry to describe  
950 movements in freshwater. Transactions of the American Fisheries Society **132**(3): 409-424.

951 Wright, S. 1950. Genetic structure of populations. British medical journal **2**(4669): 36.

952

953

954

## Table captions

Table 1. Canonical coefficients for the first two canonical variates (CV) performed on elemental and Sr isotopic ratios in water samples.

Table 2. Distance travelled by Allis shad strayers between natal and spawning river measured by a straight-line between the two river mouths. Distance classes were: short (20-100 km), Middle (100-300 km), Long (300-700 km) and Ultra long (> 700 km).

Table 3: Number of individuals from each river assigned to a sampling river. All individuals ( $n = 287$ ) were used in the baseline and in assignment tests. Only individuals with  $P > 90\%$  were reported here.

Table 4: Pairwise  $F_{ST}$  values between Allis shad populations (populations defined by otolith natal origin).

Table 5: Number of individuals from each river assigned to the otolith baseline. Only known origin individuals inferred from otolith chemistry ( $n = 275$ ) were used in the baseline. All individuals ( $n = 287$ ) were used in assignment tests. Only individuals with  $P > 90\%$  were reported here.

# Figure captions

Fig. 1. Picture of adult Allis shad otolith showing the two areas ablated by the laser prior to ICP-MS (right semi corona) and MC-ICP-MS analysis (left semi corona). Semi coronas of 60  $\mu\text{m}$  thick (difference between the inner 40  $\mu\text{m}$  and outer radius 100 $\mu\text{m}$ ) are centered on the primordium and correspond to the juvenile freshwater period of growth only.

Fig. 2. Box plot of mean (a) Sr:Ca, (b) Ba:Ca, and (c)  $^{87}\text{Sr}:^{86}\text{Sr}$  ratios of water samples from the sampling locations. Abbreviations are used for three French rivers: Adour river (Adour R.), Garonne river (Gar.), and Dordogne river (Dord.). Interquartile ranges (25th and 75th percentile) are shown by extent of boxes, and horizontal lines represent medians (50th percentile). Whiskers range from 10th to 90th percentiles, and values outside this range are plotted with circles.

Fig. 3. Canonical discriminant plot of isotope ( $^{87}\text{Sr}:^{86}\text{Sr}$ ) and elemental (Sr:Ca and Ba:Ca) signatures from water samples collected from 17 rivers in 2013. Symbols represent water samples, and ellipses are 95% confidence intervals around each group. Grid scale is given (d = 1).

Ratios	CV1	CV2
$^{87}\text{Sr}:^{86}\text{Sr}$	-0.880	0.362
Sr:Ca	0.151	-0.518
Ba:Ca	0.034	-0.205

Collection site	Individuals displaying northward movements	Individuals displaying southward movements	Distance between natal and spawning river			
			Short	Middle	Long	Ultra Long
Vire (34)	34 (100%)				34	
Aulne (12)						
Scorff (10)						
Blavet (7)						
Vilaine (19)		3 (16%)	3			
Loire (28)		3 (11%)	3			
Dordogne (71)						
Garonne (64)	14 (22%)			14		
Adour R. (6)		1 (17%)			1	
Adour E. (31)		1 (3%)			1	
Saison (6)	3 (50%)		3			
Nivelle (16)						
Minho (87)		1 (1%)				1
Lima (4)		2 (50%)	2			
Mondego (15)		15 (100%)		11		4

River	Reference populations based on geographic origins									
	Adour2013	Aulne2013	Dordogne2013	Blavet2013	Loire2013	Nivelle2009	Scorff2013	Vire2013	Garonne2012	Garonne2013
Adour (46)	9 (20%)					1				
Aulne (14)		9 (64%)								
Dordogne (69)			1 (1.4%)							
Blavet (17)				8 (47%)						
Loire (24)					8 (33%)					
Nivelle 2009 (17)						16 (94%)				
Scorff (10)							6 (60%)			
Vire (29)								17 (58%)	1	
Garonne 2012 (25)									4 (16%)	
Garonne 2013 (36)										1 (3%)

	Aulne	Blavet	Dordogne	Loire	Vilaine	Nivelle	Oloron
Adour	0.0354	0.0151	<b><i>0.0053</i></b>	<b><i>0.0131</i></b>	0.0305	0.036	<b><i>0.0037</i></b>
Aulne		<b><i>0.006</i></b>	0.0414	0.0415	0.0562	0.0447	0.035
Blavet			0.0278	0.0244	0.026	0.0409	0.0305
Dordogne				0.0109	0.027	0.0476	<b><i>0.017</i></b>
Loire					0.0372	0.0443	<b><i>0.0192</i></b>
Vilaine						0.0473	0.0588
Nivelle							0.0399


In bold-italic: non-significant pairwise  $F_{ST}$  value after Bonferroni correction.

River	Reference populations based on otolith natal origin							
	Adour R.	Oloron	Aulne	Dordogne	Blavet	Loire	Nivelle	Vilaine
Adour (46)	6 (13%)	9 (20%)	1				1	
Aulne (14)		1	9 (64%)		1			
Dordogne (69)	1	4 (6%)		3 (4%)		2		1
Blavet (17)					7 (41%)			
Loire (24)						11 (46%)		
Nivelle 2009 (17)							16 (64%)	
Scorff (10)					2			1
Vire (29)								18 (62%)
Garonne 2012 (25)								
Garonne 2013 (36)		1		5 (14%)				






Picture of adult Allis shad otolith showing the two areas ablated by the laser prior to ICP-MS (right semi corona) and MC-ICP-MS analysis (left semi corona). Semi coronas of 60  $\mu\text{m}$  thick (difference between the inner 40  $\mu\text{m}$  and outer radius 100 $\mu\text{m}$ ) are centered on the primordium and correspond to the juvenile freshwater period of growth only.  
484x361mm (72 x 72 DPI)



Box plot of mean (a) Sr:Ca, (b) Ba:Ca, and (c) <sup>87</sup>Sr:<sup>86</sup>Sr ratios of water samples from the sampling locations. Abbreviations are used for three French rivers: Adour river (Adour R.), Garonne river (Gar.), and Dordogne river (Dord.). Interquartile ranges (25th and 75th percentile) are shown by extent of boxes, and horizontal lines represent medians (50th percentile). Whiskers range from 10th to 90th percentiles, and values outside this range are plotted with circles.

192x194mm (300 x 300 DPI)



Canonical discriminant plot of isotope ( $^{87}\text{Sr}:^{86}\text{Sr}$ ) and elemental ( $\text{Sr}:\text{Ca}$  and  $\text{Ba}:\text{Ca}$ ) signatures from water samples collected from 17 rivers in 2013. Symbols represent water samples, and ellipses are 95% confidence intervals around each group. Grid scale is given ( $d = 1$ ).  
239x267mm (300 x 300 DPI)