Variation and spatial Simpson integration for reflected-clock iterated Brownian motion

Project Stardust

Reviewed editorial draft, October 30, 2026

Abstract

Let $Z_t = W(|B_t|)$, where W and B are independent standard Brownian motions. We prove convergence in probability of oriented spatial Simpson sums for every global C^3 one-form. The limit is its ordinary spatial primitive evaluated along Z. The proof uses a compact fourth-order Peano remainder and a direct quartic tightness estimate. Separately, an exact scenery-overlap bound and Brownian clock localization prove the uniform-grid quartic limit $3T$, quadratic-variation divergence and the need for compensation between raw Taylor channels. Continuous spatial weights yield a combined three-channel C^4 Taylor limit with correction $1/8$ times the fourth-derivative time integral, without global derivative bounds. A separate spatial-kernel concentration argument proves that a specific source-compensated grid has diverging L^2 norm even after projection onto the full actual IBM observable space. Nevertheless, fixed quadratic and bounded C^2 terminal tests admit strong clock-source contraction limits, with explicit capped-occupation corrections. A faithful quotient Banach domain also supports clock-L^1-valued limits of the original source contractions. We then prove a deterministic nonseparable C^1-time/C^4-space chain formula, uniform control over prefixes of one common grid, bounded random-horizon convergence for spatial one-forms, and an explicit Bernstein realization converging in continuous-path supremum norm. The time and space hypotheses, measurability, and limiting topologies are stated separately. These prescribed symmetric quadratures are not identified with a causal physical-time divergence. The principal results have exact declarations in a frozen Lean development; the exposition supplies full human proofs. Novelty of this package is unresolved.

1 The process and the quadrature prescription

Work on a probability space carrying independent real standard Brownian motions $B = (B_t)_{t\geq 0}$ and $W = (W_s)_{s\geq 0}$, starting at zero, with continuous versions. Define

$$Z_t = W(|B_t|), \qquad t \geq 0. \tag{1}$$

Thus the clock is reflected and the scenery is indexed by nonnegative spatial time. This convention is fixed throughout. The product Wiener realization is equivalent for these statements to the constructed independent Gaussian scenery–clock realization used in the formal sources. The process is jointly measurable and has continuous paths on a common full-measure set. All samplewise arguments below are on that set.

For a real function g, define the oriented spatial Simpson increment

$$S_g(x,y) = \frac{y-x}{6}\left(g(x) + 4g\left(\frac{x+y}{2}\right) + g(y)\right). \tag{2}$$

The sign of $y - x$ is retained. The midpoint is the spatial midpoint of the two endpoints; it is not evaluation of Z at the midpoint of a physical time interval.

Fix $T > 0$, use all uniform grids $t_i = iT/N$, $0 \le i \le N$, and abbreviate $Z_i = Z_{t_i}$. Define

$$P_{g,N}(k) = \sum_{i=0}^{k-1} S_g(Z_i, Z_{i+1}), \qquad 0 \le k \le N.$$

The finite sums are defined directly from the integrand and the process values. The limiting primitive will be proved to agree with them, rather than being substituted as their definition.

Theorem 1.1. *Let $g \in C^3(\mathbb{R})$ and set*

$$J_g(t) = \int_{Z_0}^{Z_t} g(x)\, dx,$$

where the integral is oriented. Then J_g has continuous paths, is jointly measurable, and

$$\max_{0 \le k \le N} |P_{g,N}(k) - J_g(t_k)| \xrightarrow{\mathrm{P}} 0. \tag{3}$$

No global growth bound on g or its derivatives is assumed.

The uniformity in (3) is over prefixes of one common grid. It is not an assertion that arbitrary partitions or arbitrary adapted time-dependent integrands are covered. The deterministic time–space formula and the continuous-path realization will have their own definitions below.

Stochastic integration for iterated Brownian motion has classical antecedents. Khoshnevisan and Lewis [2] use a two-sided scenery and a random partition induced by a spatial clock grid. Our reflected-clock convention and deterministic physical-time grids are different. Simpson-type stochastic sums have also been studied for fractional Brownian motion [3]; covariance or scaling similarity does not identify those processes with (1). The paper gives a verified, domain-explicit exposition, not a claim of first construction of symmetric integration for iterated Brownian motion.

2 A direct quartic tightness estimate

For an interval $[a, b]$ let $\Delta Z = Z_b - Z_a$. Conditional on the complete clock path, the two scenery times are fixed and ΔZ is a centered Gaussian variable of variance

$$L = \big||B_b| - |B_a|\big|.$$

Independence of W and B is essential here. Reversal of the order of the scenery times only changes the sign of a Brownian increment and not its Gaussian fourth moment. Therefore

$$\mathbb{E}[(\Delta Z)^4 \mid B] = 3L^2, \qquad \mathbb{E}[(\Delta Z)^4] \le 3(b - a), \tag{4}$$

using $L \le |B_b - B_a|$ and the Brownian increment variance. These expectations can equally be computed by Fubini on the independent scenery–clock product space, so no unspecified conditional version is required.

Put

$$Q_{4,N} = \sum_{i=0}^{N-1} |Z_{i+1} - Z_i|^4, \qquad M_N = \max_{i<N} |Z_{i+1} - Z_i|.$$

Equation (4) yields

$$\mathbb{E}Q_{4,N} \le 3T, \qquad \sup_N \mathbb{P}\{Q_{4,N} > A\} \le \frac{3T}{A} \quad (A > 0). \tag{5}$$

Thus the quartic sums are tight. No independence between different $\Delta_i Z$ is needed when summing their expectations. Continuity of each path on $[0, T]$ gives $M_N \to 0$ almost surely.

The sharper assertion $Q_{4,N} \xrightarrow{\mathrm{P}} 3T$ is proved separately in Section 8; the quadrature proofs need only tightness. In particular it would be incorrect to infer that assertion from (5) or the corresponding expectation calculation. The present argument uses only the bound it has actually proved.

Lemma 2.1. *If $U_N \to 0$ in probability and $V_N \ge 0$ is tight, then $U_N V_N \to 0$ in probability.*

Proof. For $A > 0$ and $\varepsilon > 0$,

$$\mathbb{P}\{|U_N V_N| > \varepsilon\} \le \mathbb{P}\{V_N > A\} + \mathbb{P}\{|U_N| > \varepsilon/A\}.$$

First choose A to make the first term uniformly small, then let $N \to \infty$ in the second. □

This elementary transfer principle is why a local remainder modulus is sufficient; neither an almost-sure variation bound nor global bounds on the integrand are needed.

3 Compact Simpson remainders

We now establish the deterministic estimate which controls every stochastic prefix simultaneously.

Lemma 3.1. *Let $g \in C^3(\mathbb{R})$ and $R > 0$. There is a nonnegative modulus $\omega_R(\delta) \to 0$ as $\delta \downarrow 0$ such that, for $|x|, |y| \le R$,*

$$\left| S_g(x, y) - \int_x^y g(z)\, dz \right| \le \frac{1}{3}\omega_R(|y - x|)|y - x|^4. \tag{6}$$

One may take ω_R to be the modulus of continuity of g''' on $[-R, R]$.

Proof. Let $d = y - x$. The case $d = 0$ is immediate. On the segment between x and y, Taylor's theorem in integral form gives

$$g(x + \theta d) = g(x) + g'(x)\theta d + \frac{g''(x)}{2}\theta^2 d^2 + \frac{g'''(x)}{6}\theta^3 d^3 + r(\theta)$$

for $0 \le \theta \le 1$, with $|r(\theta)| \le \omega_R(|d|)|d|^3/6$. Indeed, subtract the constant $g'''(x)$ inside the third-order integral remainder and use its uniform continuity. This proof works for either sign of d.

The Simpson functional on $[0, 1]$, with weights $1/6, 4/6, 1/6$, integrates polynomials of degree at most three exactly. Directly, its values on $1, \theta, \theta^2, \theta^3$ are $1, 1/2, 1/3, 1/4$. Thus all displayed polynomial terms cancel against the actual integral. The Simpson average of the remainder and its integral each have absolute value at most $\omega_R(|d|)|d|^3/6$. Multiplying their difference by $|d|$ proves (6). □

The estimate is fourth order with a vanishing compact coefficient. It requires g''' continuous, not a fourth derivative of g. Equivalently, when $g = f'$, it requires a C^4 primitive, not a fifth derivative. This is the regularity boundary used in the formal compact Peano result.

For a continuous path $z : [0, T] \to \mathbb{R}$ with range in $[-R, R]$, sum (6) over any grid prefix. Since the exact oriented integrals telescope through the ordinary primitive,

$$\max_{k \le N} \left| \sum_{i<k} S_g(z_i, z_{i+1}) - \int_{z_0}^{z_k} g \right| \le \frac{1}{3} \omega_R(M_N(z)) Q_{4,N}(z). \tag{7}$$

There is one modulus and one full nonnegative quartic sum for every prefix. No union bound with a factor growing in N is being applied.

4 The spatial formula and measurable random horizons

Proof of Theorem 1.1. The function $G(x) = \int_0^x g(z)\,dz$ is C^4 by the fundamental theorem of calculus, so $J_g(t) = G(Z_t) - G(Z_0)$ is continuous in t and jointly measurable. Each finite Simpson sum is measurable because it is a finite sum of continuous functions of measurable coordinates.

The range of a continuous sample path is bounded, so for any $\eta > 0$ choose deterministic R with $\mathbb{P}\{\sup_{t \le T} |Z_t| > R\} < \eta$. On its complementary event (7) applies. There $\omega_R(M_N) \to 0$ almost surely. Quartic tightness (5) and Lemma 2.1 show that the right side tends to zero in probability. The probability of the excluded event is less than η, which is arbitrary. This proves the uniform-prefix assertion. □

Corollary 4.1. *Let τ be any measurable random time with $0 \le \tau \le T$. Define*

$$k_N = \left\lfloor \frac{N\tau}{T} \right\rfloor, \qquad \tau_N = \frac{T}{N} k_N.$$

Then $P_{g,N}(k_N) \xrightarrow{\mathrm{P}} J_g(\tau)$ for $g \in C^3(\mathbb{R})$.

Proof. The integer index is measurable, takes values in $\{0, \dots, N\}$, and obeys $0 \le \tau - \tau_N < T/N$. Evaluating the finite collection of measurable prefix sums at that index is measurable. The error to $J_g(\tau_N)$ is bounded by the prefix maximum in (3). Pathwise continuity gives $J_g(\tau_N) \to J_g(\tau)$ almost surely, proving the result. □

No stopping-time hypothesis is used. For example τ can equal T or $T/2$ according to the sign of Z_T. This conclusion uses uniform approximation and continuity, not optional sampling of a martingale.

5 Deterministic time–space integrands

Let $F : \mathbb{R}^2 \to \mathbb{R}$ satisfy the following hypotheses: for every fixed t, $F(t, \cdot) \in C^4(\mathbb{R})$; for every fixed x, $F(\cdot, x) \in C^1(\mathbb{R})$; and the functions $\partial_t F$ and $\partial_x^4 F$ are jointly continuous. No mixed derivative or global derivative bound is assumed. For the continuous-path realization in Section 7, we additionally state joint continuity of F explicitly, as in its formal declaration.

Define the time-frozen spatial sums

$$P_{F,N}(k) = \sum_{i<k} S_{\partial_x F(t_i, \cdot)}(Z_i, Z_{i+1}), \tag{8}$$

and the continuous target

$$J_F(t) = F(t, Z_t) - F(0, Z_0) - \int_0^t \partial_t F(s, Z_s)\, ds. \tag{9}$$

The target is continuous. In fact the stated hypotheses already give joint continuity of F: near a fixed time t_0, write $F(t,x) = F(t_0,x) + \int_{t_0}^t \partial_t F(s,x)\, ds$ and use continuity and local boundedness of the joint time partial. We keep the explicit hypothesis for the later source correspondence.

Theorem 5.1. *Under these time–space hypotheses,*

$$\max_{k \le N} |P_{F,N}(k) - J_F(t_k)| \xrightarrow{\mathrm{P}} 0. \tag{10}$$

In particular the full sums converge in probability to (9) *at* $t = T$.

Proof. On a compact rectangle $[0,T] \times [-R,R]$, joint continuity of $\partial_x^4 F$ gives one horizontal modulus uniform in t. Apply Lemma 3.1 with $g = \partial_x F(t,\cdot)$. The proof uses only differences of $\partial_x^4 F$ at nearby spatial points, so its modulus works for every frozen time. Thus the maximum error in replacing (8) by

$$V_{F,N}(k) = \sum_{i<k} \big(F(t_i, Z_{i+1}) - F(t_i, Z_i)\big)$$

tends to zero in probability, by exactly the same localization and quartic-tightness argument as before.

It remains to identify $V_{F,N}$. The exact finite chain identity is

$$\begin{aligned} F(t_k, Z_k) - F(0, Z_0) = V_{F,N}(k) \\ + \sum_{i<k} \big(F(t_{i+1}, Z_{i+1}) - F(t_i, Z_{i+1})\big). \end{aligned} \tag{11}$$

The time fundamental theorem of calculus identifies the second sum as

$$\sum_{i<k} \int_{t_i}^{t_{i+1}} \partial_t F(s, Z_{i+1})\, ds.$$

For each fixed continuous sample path, uniform continuity of $\partial_t F$ on its compact time–space range shows that its difference from $\int_0^{t_k} \partial_t F(s, Z_s)\, ds$ is bounded, uniformly over k, by

$$T\, \nu_R\!\left(\sup_{|t-s| \le T/N} |Z_t - Z_s| \right) \longrightarrow 0.$$

Here ν_R is a horizontal modulus of the time partial on the compact rectangle. This is a pathwise limit. Combining it with the spatial error and (11) proves (10). □

The time correction is an ordinary finite Lebesgue integral on each sample path. It is not an assumed stochastic correction term. Joint measurability of $\partial_t F(s, Z_s)$ and integration over a bounded interval make it a measurable random variable. The same argument gives measurability of the finite sums, since each frozen spatial slice has a continuous derivative.

6 Weights and elementary checks on the prescription

Taking $F(t,x) = h(t)G(x)$, where $h \in C^1(\mathbb{R})$ and $G' = g \in C^3(\mathbb{R})$, gives

$$\sum_{i<N} h(t_i) S_g(Z_i, Z_{i+1}) \xrightarrow{\mathrm{P}} h(T) J_g(T) - \int_0^T h'(s) J_g(s)\, ds. \tag{12}$$

The equality of this target with the one in Theorem 5.1 follows from cancellation of the constant $G(Z_0)$ and $\int_0^T h' = h(T) - h(0)$. Thus the separable time-weighted rule is contained in the full nonseparable theorem. Random time coefficients are not included by replacing h with a random function without further work.

The spatial rule is exact for cubic one-forms. For example

$$S_{4z^3}(x,y) = y^4 - x^4.$$

It is not always an exact increment of a primitive: direct polynomial expansion gives

$$S_{5z^4}(x,y) - (y^5 - x^5) = \frac{(y-x)^5}{24}. \tag{13}$$

For a continuous path the sum of absolute fifth powers is at most $M_N Q_{4,N}$, which tends to zero in probability here. This example illustrates the finite prescription and its limiting primitive without conflating them.

The sign in (2) also matters. Replacing $y - x$ by $|y - x|$ would already destroy the oriented integral for a constant one-form. Similarly, replacing the spatial midpoint by $Z_{(t_i+t_{i+1})/2}$ changes the approximation scheme; the deterministic Simpson polynomial cancellations no longer apply to that expression.

7 An explicit continuous-path realization

Under the time–space hypotheses, including explicit joint continuity of F, define on $0 \le u \le 1$

$$A_{F,N}(u) = \sum_{k=0}^{N} \binom{N}{k} u^k (1-u)^{N-k} P_{F,N}(k). \tag{14}$$

This is an actual random polynomial assembled from the prefixes of the one common grid. Let $j_F(u) = J_F(Tu)$.

Theorem 7.1. *As random elements of $C([0,1],\mathbb{R})$ with supremum norm,*

$$\|A_{F,N} - j_F\|_\infty \xrightarrow{\mathrm{P}} 0. \tag{15}$$

The approximants and target are Borel measurable in this continuous-path space.

Proof. The Bernstein basis is nonnegative and sums to one, so

$$\|A_{F,N} - B_N j_F\|_\infty \le \max_{k \le N} |P_{F,N}(k) - j_F(k/N)|, \tag{16}$$

where B_N is the ordinary Bernstein approximation operator. The right side tends to zero in probability by Theorem 5.1.

For a fixed continuous f, Bernstein approximation gives $\|B_N f - f\|_\infty \to 0$. A direct proof writes $(B_N f)(u) = \mathbb{E}[f(K/N)]$ for a binomial K of parameters N, u. Split at $|K/N - u| \le \delta$ and use $\mathbb{E}(K/N - u)^2 = u(1-u)/N \le 1/(4N)$ to obtain

$$\|B_N f - f\|_\infty \le \omega_f(\delta) + \frac{2\|f\|_\infty}{4N\delta^2}.$$

First take $N \to \infty$ and then $\delta \downarrow 0$. Apply this pathwise to the continuous target j_F. The resulting almost-sure Bernstein error tends to zero in probability. Combine it with (16).

Each coefficient in (14) is measurable, hence the finite polynomial is a measurable $C([0,1])$-valued map. For the target, rational evaluation maps are measurable by the measurability arguments following Theorem 5.1. The Borel sigma algebra of the separable space $C([0,1])$ is generated by these evaluations: suprema of continuous-path distances equal suprema over a countable dense subset. Thus the target path map is measurable as well. □

The approximation has $A_{F,N}(0) = 0$ and $A_{F,N}(1) = P_{F,N}(N)$, since the Bernstein basis selects the first and last coefficients at the endpoints. At intermediate times, all grid prefixes can enter (14). Consequently it is a polynomial smoothing, not a claimed causal interpolation. No adaptedness or L^p convergence follows merely from (15).

8 Exact uniform-grid quartic variation

The tightness estimate used above is deliberately weaker than the library's full variation result. We now reconstruct that result without changing the proof or domain of the Simpson formulas. For $t_i = iT/N$, let

$$Q_N = \sum_{i=0}^{N-1} (Z_{i+1} - Z_i)^4, \qquad C_N = 3\sum_{i=0}^{N-1} (|B_{t_{i+1}}| - |B_{t_i}|)^2.$$

The second quantity is the exact scenery-conditional mean of the first, since each scenery increment has conditional variance $\ell_i = \big||B_{t_{i+1}}| - |B_{t_i}|\big|$.

Theorem 8.1. *For every $T > 0$, along all positive integer uniform-grid sizes,*

$$Q_N \xrightarrow{\mathrm{P}} 3T. \tag{17}$$

The scenery increments need not be independent after conditioning on the clock. A later traversal may revisit a previous scenery interval. We keep all such overlaps in the variance estimate.

The exact Gaussian overlap calculation

Write $a_i = |B_{t_i}|$, $b_i = |B_{t_{i+1}}|$ and $I_i = (\min(a_i, b_i), \max(a_i, b_i)]$. Let o_{ij} be the signed overlap of these oriented intervals; its absolute value is $|I_i \cap I_j|$. Brownian scenery increments $G_i = W_{b_i} - W_{a_i}$ satisfy, for this fixed clock sample,

$$\mathbb{E}_W G_i^2 = \ell_i, \qquad \mathbb{E}_W(G_i G_j) = o_{ij}.$$

The second equality follows by writing each increment as the oriented integral of an interval indicator against Brownian motion. It also follows directly by expanding the covariance

$\mathbb{E}(W_u W_v) = \min(u, v)$. For centered jointly Gaussian variables with variances l, m and covariance o, the pair-moment formula gives

$$\mathrm{Cov}(G^4, H^4) = 72lm\,o^2 + 24o^4. \tag{18}$$

To verify the constants, among pairings of four G's and four H's, zero cross-pairs contribute $9l^2m^2$, two cross-pairs contribute $\binom{4}{2}^2 2! = 72$ copies of $lm\,o^2$, and four cross-pairs contribute $4! = 24$ copies of o^4. Subtract $\mathbb{E}G^4\mathbb{E}H^4 = 9l^2m^2$. The pairing rule itself follows by expanding the Gaussian generating function $\exp((lx^2 + 2oxy + my^2)/2)$ at x^4y^4.

If all $\ell_i \le \delta$, then $|o_{ij}| \le \delta$ and

$$\mathbb{E}_W(Q_N - C_N)^2 \le 96\delta^3 \sum_{i,j} |o_{ij}|. \tag{19}$$

Indeed, each summand in (18) is bounded by its coefficient times $\delta^3|o_{ij}|$. Every finite eighth moment exists, so the double variance expansion is legitimate. Equivalently, if $n(x) = \sum_i \mathbf{1}_{I_i}(x)$ is the return count, then $\sum_{i,j} |o_{ij}| = \int n(x)^2 dx$. Repeated visits are counted quadratically, as required.

A Brownian endpoint envelope

Define $\tau_\delta(x) = (2\delta - |x|)_+$ and

$$E_{N,\delta} = \sum_{i,j=0}^{N-1} \tau_\delta(|B_{t_i}| - |B_{t_j}|), \qquad p(s) = (2\pi s)^{-1/2} \quad (s > 0).$$

For two intervals starting at a, c and of lengths at most δ, their overlap length is at most $(2\delta - |a - c|)_+$. This follows by enclosing them in $[a - \delta, a + \delta]$ and $[c - \delta, c + \delta]$ and intersecting these two intervals. Consequently, on the event

$$\mathcal{A}_{N,\delta} = \{\max_i |B_{t_{i+1}} - B_{t_i}| < \delta\},$$

the reverse triangle inequality gives $\ell_i \le \delta$ and $\sum_{i,j} |o_{ij}| \le E_{N,\delta}$.

For $s < t$, condition on $B_s = x$. The later value is $x + G$, where G is independent normal with variance $t - s$. The folded variable $|x + G|$ has density on $[0, \infty)$ bounded by $2p(t - s)$, uniformly in x. The tent has height at most 2δ, support length 4δ, and hence integral at most $8\delta^2$. Thus

$$\mathbb{E}\tau_\delta(|B_s| - |B_t|) \le 16\delta^2 p(t - s). \tag{20}$$

The diagonal is instead exactly 2δ; it is not evaluated through a singular zero-variance density formula. Set $h = T/N$. Since $p(kh) = p(h)/\sqrt{k}$ and

$$\sum_{k=1}^{N} \frac{1}{\sqrt{k}} \le 2\sqrt{N},$$

summing the ordered off-diagonal pairs gives

$$\mathbb{E}E_{N,\delta} \le 2N\delta + 64\delta^2 p(h) N\sqrt{N} = 2N\delta + 64p(T)N^2\delta^2. \tag{21}$$

The inverse-square-root sum follows by telescoping $1/\sqrt{k} \le 2(\sqrt{k} - \sqrt{k-1})$.

Let $m_{20} = \mathbb{E}|G_0|^{20}$ for a standard normal G_0; this is a finite fixed constant. The union bound and Markov's inequality give

$$\mathbb{P}(\mathcal{A}_{N,\delta}^c) \le N(T/N)^{10} m_{20} \delta^{-20}. \tag{22}$$

Choose $\delta = \delta_N = N^{-4/9}$. The right side is $T^{10} m_{20} N^{-1/9} \to 0$, whereas

$$\delta_N^3 \mathbb{E}E_{N,\delta_N} \le 2N^{-7/9} + 64p(T)N^{-2/9} \longrightarrow 0. \tag{23}$$

For any $\varepsilon > 0$, conditional Chebyshev on the good-clock event, followed by Fubini, now yields

$$\mathbb{P}(|Q_N - C_N| \ge \varepsilon) \le \mathbb{P}(\mathcal{A}_{N,\delta_N}^c) + \frac{96}{\varepsilon^2} \delta_N^3 \mathbb{E}E_{N,\delta_N} \longrightarrow 0. \tag{24}$$

On the bad event the conditional probability is bounded by one. No global joint L^2 error bound is inferred by removing that event.

The reflected-clock correction

For real a, b define $r(a,b) = 2(|ab| - ab) \ge 0$. Direct expansion gives

$$(|b| - |a|)^2 = (b-a)^2 - r(a,b). \tag{25}$$

Let $R_N = \sum_i r(B_{t_i}, B_{t_{i+1}})$. If $|b-a| \le \delta$, then

$$r(a,b) \le \delta \tau_\delta(a). \tag{26}$$

When $ab \ge 0$ the left side is zero. Otherwise the endpoints straddle zero, so $|a| \le \delta$ and $r(a,b) = 4|ab| \le (b-a)^2 \le \delta^2$; also $\tau_\delta(a) \ge \delta$. This proves the bound in both cases. Therefore, on $\mathcal{A}_{N,\delta}$,

$$0 \le R_N \le \delta \sum_i \tau_\delta(B_{t_i}).$$

The initial term is bounded by $2\delta^2$. At positive grid times, the ordinary Gaussian density and the same tent-mass bound give $\mathbb{E}\tau_\delta(B_{t_i}) \le 16\delta^2 p(t_i)$. Summing as before, with a harmless upper bound including one extra positive time, gives

$$\mathbb{E}\left[\delta \sum_i \tau_\delta(B_{t_i})\right] \le 2\delta^2 + 32p(T)N\delta^3.$$

For $\delta_N = N^{-4/9}$ this tends to zero. Markov's inequality, with the bad-clock probability kept separate, implies $R_N \xrightarrow{\mathrm{P}} 0$.

Finally $U_N = \sum_i (B_{t_{i+1}} - B_{t_i})^2$ has expectation T and variance $2T^2/N$, by independent Brownian increments. Thus $U_N \to T$ in L^2. Summing (25) gives $C_N = 3U_N - 3R_N \xrightarrow{\mathrm{P}} 3T$. Combining this with (24) proves Theorem 8.1.

Uniform prefixes on the same physical grid

The preceding estimates also control prefixes of the original grid, without remeshing a shorter time interval. Define

$$Q_N(m) = \sum_{i<m} (Z_{i+1} - Z_i)^4, \qquad M_N = \max_{0 \le m \le N} |Q_N(m) - 3Tm/N|.$$

We claim $M_N \xrightarrow{\mathrm{P}} 0$. First fix any deterministic sequence $0 \le m_N \le N$. Removing intervals only decreases $\sum_{i,j} |o_{ij}|$, so the same good-clock conditional-variance bound and bad-clock estimate prove that the scenery-centered prefix error tends to zero in probability. The Brownian-clock prefix quadratic sum has mean $m_N T/N$ and variance $2m_N(T/N)^2 \le 2T^2/N$. Its nonnegative reflection penalty is at most the full penalty R_N, already proved to vanish in probability. Therefore

$$Q_N(m_N) - 3Tm_N/N \xrightarrow{\mathrm{P}} 0.$$

All selections in this argument are deterministic.

To obtain the maximum, fix $K \ge 1$ and put $L = \lfloor N/K \rfloor + 1$, $r_j = \min(jL, N)$ for $0 \le j \le K$. These nodes cover every integer prefix, because $KL > N$. The increasing function $m \mapsto Q_N(m)$ brackets each prefix between its neighboring coarse values. Subtracting the linear target gives the deterministic bound

$$M_N \le \max_{0 \le j \le K} |Q_N(r_j) - 3Tr_j/N| + 3T/K + 3T/N.$$

For fixed K, the finite maximum tends to zero in probability by the prefix result and a union bound. Choose K large first and then let $N \to \infty$. This proves the claim. In particular this is a maximum of partial errors, rather than an unjustified union bound over an increasing number of prefixes.

Every continuous spatial weight

For every continuous $\varphi : \mathbb{R} \to \mathbb{R}$, including globally unbounded ones, we have

$$\sum_{i<N} \varphi(Z_i)(Z_{i+1} - Z_i)^4 \xrightarrow{\mathrm{P}} 3 \int_0^T \varphi(Z_t)\, dt. \tag{27}$$

Here the integral is a genuine finite pathwise time integral. No moment or global bound on φ is required.

We give the deterministic quadrature argument before applying it to random weights. Suppose nonnegative masses $q_{N,i}$ on the uniform grid obey

$$\max_{m \le N} \left| \sum_{i<m} q_{N,i} - 3Tm/N \right| = \eta_N \longrightarrow 0.$$

For a Lipschitz function p on $[0,T]$ with $|p| \le B$ and Lipschitz constant A, Abel summation bounds

$$\left| \sum_{i<N} p(t_i)(q_{N,i} - 3T/N) \right| \le \eta_N \left(|p(t_{N-1})| + \sum_{i<N-1} |p(t_{i+1}) - p(t_i)| \right) \le \eta_N (B + AT).$$

The left Riemann-sum error for $3\int_0^T p$ is at most $3AT^2/N$, by integrating $A|t-t_i| \le AT/N$ on each cell. For continuous g, choose a piecewise linear p with $\|g-p\|_\infty \le \varepsilon$. Such a choice follows by linear interpolation and uniform continuity on the compact interval. The additional error is bounded by

$$\varepsilon\left(\sum_{i<N} q_{N,i} + 3T\right).$$

The mass sum tends to $3T$. First pass to the limit in N with the approximation fixed, then let $\varepsilon \downarrow 0$. Thus the quadrature converges for every continuous g. The formal proof uses polynomial approximation in place of linear interpolation; the deterministic estimate and positive-mass mechanism are the same.

Now take any subsequence of grid sizes. Since $M_N \to 0$ in probability, it has a further subsequence on which $M_N \to 0$ almost surely. This standard extraction follows by choosing successive indices with probabilities $\mathbb{P}(M_N > 2^{-j}) < 2^{-j}$ and applying the summable-event lemma. On each continuous sample path in this full-measure set, apply the deterministic argument with $q_{N,i} = (Z_{i+1} - Z_i)^4$ and $g(t) = \varphi(Z_t)$. The resulting convergence is almost sure along that further subsequence. The subsequence criterion proves (27): if probability convergence failed, some subsequence would have a fixed positive tail probability, which its almost-sure further subsequence contradicts. Finite sums are measurable by continuity of φ and the coordinate maps; the time integral is measurable by joint measurability and integration over $[0,T]$. No measurability of a selected samplewise approximation or compact-path bound is needed.

The combined first three Taylor channels on the full C^4 domain

For $f \in C^4(\mathbb{R})$ define the actual finite sum

$$A_N(f) = \sum_{i<N}\left[f'(Z_i)\Delta Z_i + \tfrac{1}{2}f''(Z_i)(\Delta Z_i)^2 + \tfrac{1}{6}f'''(Z_i)(\Delta Z_i)^3\right], \qquad \Delta Z_i = Z_{i+1} - Z_i.$$

Then at each fixed $T > 0$,

$$A_N(f) \xrightarrow{\mathrm{P}} f(Z_T) - f(Z_0) - \tfrac{1}{8}\int_0^T f^{(4)}(Z_t)\,dt. \tag{28}$$

All derivatives may be globally unbounded. For a fixed continuous sample path, choose R containing its range. Taylor's theorem and uniform continuity of $f^{(4)}$ on $[-R,R]$ give

$$f(y) - f(x) = f'(x)(y-x) + \tfrac{1}{2}f''(x)(y-x)^2 + \tfrac{1}{6}f'''(x)(y-x)^3 + \tfrac{1}{24}f^{(4)}(x)(y-x)^4 + r(x,y),$$

where $|r(x,y)| \le \omega_R(|y-x|)|y-x|^4/24$ and $\omega_R(a) \to 0$ as $a \downarrow 0$. For example, the third-order Lagrange remainder is $f^{(4)}(z)(y-x)^4/24$ for a point z between x and y; subtracting the fourth coefficient proves this bound, also when $x = y$. Since the path mesh $m_N \to 0$ almost surely,

$$\sum_{i<N} |r(Z_i, Z_{i+1})| \le \omega_R(m_N)Q_N/24 \xrightarrow{\mathrm{P}} 0.$$

To justify the last passage without a random integrable dominator, extract from any subsequence a further one on which $Q_N \to 3T$ almost surely and use the pathwise bound. The same subsequence criterion proves the stated probability convergence.

Finally telescope the exact Taylor identities. The sum tends by (27) to $3\int_0^T f^{(4)}(Z_t)dt$; the total signed remainder is bounded by the vanishing total absolute remainder. Thus $3/24 = 1/8$

gives (28). The three lower ranks remain grouped in their genuine original grid sum. This is not an assertion of their separate convergence, an L^p limit, a simultaneous-in-time version, or a classical observed-filtration stochastic integral. It supplies the precise compensated change-of-variables statement consistent with the raw-channel obstruction below.
`IBMUniformSubfamilyQuartic`,
`IBMUniformPrefixClock`,
`IBMUniformPrefixQuartic`,
`IBMUniformPrefixMaximum`,
`IBMContinuousQuadrature`,
`IBMContinuousWeightedQuartic`,
`IBMFourthPeano`,
`IBMActualPeanoRemainder`,
`IBMActualTaylorChain`.

The classical quartic-variation antecedents cited above remain applicable context. Our contribution here is the complete human reconstruction of these frozen probability limits and their exact C^4 hypothesis, with no priority claim for a new IBM calculus.

9 Divergent raw channels and finite compensated sums

Let

$$V_N = \sum_i (Z_{i+1} - Z_i)^2, \qquad m_N = \max_i |Z_{i+1} - Z_i|.$$

Path continuity on $[0, T]$ implies $m_N \to 0$ almost surely, and the elementary inequality

$$Q_N \le m_N^2 V_N \tag{29}$$

keeps the quartic and quadratic sums distinct.

Corollary 9.1. *For $T > 0$ and every fixed real K,*

$$\mathbb{P}(V_N \le K) \longrightarrow 0.$$

Neither V_N nor the raw first Taylor channel $F_N = \sum_i 2Z_i(Z_{i+1} - Z_i)$ has a finite probability limit. Nevertheless their sum is exactly $Z_T^2 - Z_0^2$ for every grid.

Proof. For $K < 0$ the event is empty. For $K \ge 0$, on $\{V_N \le K\}$, (29) gives $Q_N \le K m_N^2$. Since $Q_N \xrightarrow{\mathrm{P}} 3T > 0$ and $K m_N^2 \xrightarrow{\mathrm{P}} 0$, the event has probability tending to zero. If $V_N \xrightarrow{\mathrm{P}} Y$ for a finite random variable, choose K with $\mathbb{P}(|Y| > K) < 1/4$. For all sufficiently large N, $\mathbb{P}(|V_N - Y| > 1) < 1/4$ would give $\mathbb{P}(V_N \le K + 1) > 1/2$, contradicting the lower-tail result.

Expanding $(Z_{i+1})^2 - Z_i^2$ and telescoping gives

$$F_N + V_N = Z_T^2 - Z_0^2.$$

A finite probability limit for F_N would therefore give one for V_N by subtraction from this fixed finite endpoint variable. This is impossible. The compensation is essential even for the polynomial z^2; it cannot be replaced by separately convergent raw first and second channels. □

Burdzy's classical *Variation of iterated Brownian motion* [1] already establishes deterministic quartic variation for two-sided-scenery IBM. Khoshnevisan–Lewis [2], Theorem 4.3, proves a stronger weighted quartic limit on its induced random partition. These are explicit antecedents. Our source-mapped proof retains the reflected-clock model, all uniform physical grids and convergence in probability; it claims no priority for the quartic phenomenon and no almost-sure or L^2 upgrade. The continuously weighted uniform-grid limit above is proved in probability.

10 Compensated grids still escape after observable projection

The raw-channel divergence does not by itself exclude a compensated L^2 limit. We now prove that a specific original compensated grid also has diverging norm, even after conditioning on the entire IBM observable sigma algebra. This assertion concerns a definite sequence; it does not exclude other stochastic integral domains or prescriptions.

Write $r_i = |B_{t_i}|$, $h_i = \mathbf{1}_{(0,r_i]} \in L^2(\mathbb{R}_+)$, $d_i = h_{i+1} - h_i$, and use the scenery isonormal notation $X(h) = \int h\, dW$, so $Z_i = X(h_i)$. All these formulas can be read conditionally on B, where h_i, d_i are deterministic. Define the actual finite random variables

$$R_N = \sum_{i<N} Z_i(Z_{i+1} - Z_i), \qquad C_N = \sum_{i<N} \langle d_i, h_i \rangle = \sum_{i<N} (\min(r_i, r_{i+1}) - r_i), \qquad S_N = R_N - C_N.$$

The conditional scenery mean of S_N is zero. Gaussian moments of the finitely many clock coordinates and conditional fourth moments show that $R_N, C_N, S_N \in L^2$ for every N; for example $\|d_i\|^2 = |r_{i+1} - r_i|$ and $\|h_i\|^2 = r_i$, and the conditional second moment of each product is bounded by $3\|d_i\|^2\|h_i\|^2$. Let

$$\mathcal{O} = \sigma(Z_t : t \geq 0), \qquad I_N = \mathbb{E}[S_N \mid \mathcal{O}].$$

This uses the full observable space, with no changing-past or predictable identification.

Theorem 10.1. *For every fixed $T > 0$ along all uniform physical grids,*

$$a_N := \mathbb{E}[S_N^2] \longrightarrow \infty, \qquad \mathrm{Var}(C_N) \leq \frac{9}{2}T, \qquad \|I_N\|_2 \longrightarrow \infty.$$

More precisely,

$$a_N^2 \leq \|I_N\|_2^2 (a_N + \mathrm{Var}(C_N)).$$

Thus this projected compensated sequence has no finite L^2 limit.

10.1 Folded clock mass and a concentrating spatial kernel

Put $h = T/N$ and $U_N = \sum_{i<N} (B_{t_{i+1}} - B_{t_i})^2$. Independent increments give $\mathbb{E}U_N = T$ and $\mathrm{Var}(U_N) = 2T^2/N$. We need the folded version

$$q_N = \sum_{i<N} (r_{i+1} - r_i)^2 \longrightarrow T \quad \text{in } L^1.$$

Here is a direct proof of that additional assertion. The loss $(y-x)^2 - (|y| - |x|)^2$ is nonnegative and at most $(y-x)^2$. It vanishes when x, y have the same sign. For any $\delta > 0$, a crossing with $|y - x| \leq \delta$ has $|x| \leq \delta$. Consequently, writing $\Delta_i B = B_{t_{i+1}} - B_{t_i}$,

$$0 \leq U_N - q_N \leq \sum_{i<N} (\Delta_i B)^2 \left(\mathbf{1}_{\{|B_{t_i}| \leq \delta\}} + \mathbf{1}_{\{|\Delta_i B| > \delta\}} \right).$$

The first summand at $i = 0$ has expectation h. For $i \geq 1$, independence of B_{t_i} and $\Delta_i B$ and the normal density bound give a contribution at most $h\sqrt{2/\pi}\,\delta/\sqrt{ih}$. Since $\sum_{i=1}^{N-1} i^{-1/2} \leq 2\sqrt{N}$, their sum is at most $2\sqrt{2/\pi}\,\delta\sqrt{T}$. The large-increment contribution is at most $\sum_i \mathbb{E}(\Delta_i B)^4/\delta^2 = 3Th/\delta^2$. For fixed δ, let $N \to \infty$, and then $\delta \downarrow 0$. This proves $\mathbb{E}(U_N - q_N) \to 0$ and hence the claimed L^1 limit. No pathwise quadratic-variation assertion has been substituted.

For a deterministic finite nonnegative radius sequence starting at zero, define the spatial interval

$$J_i = (\min(r_i, r_{i+1}), \max(r_i, r_{i+1})].$$

Form the nonnegative spatial kernel

$$K_N(x, y) = \sum_{i<N} \mathbf{1}_{J_i}(x)\mathbf{1}_{J_i}(y).$$

Its mass and energy are exactly

$$\int K_N = q_N, \qquad G_N := \int K_N^2 = \sum_{i,j<N} |J_i \cap J_j|^2 = \sum_{i,j<N} \langle d_i, d_j\rangle^2.$$

The last equality uses the signs of oriented indicator increments; their signs disappear upon squaring. If $0 \leq r_i \leq M$ and $|r_{i+1} - r_i| \leq \eta$, the kernel is supported inside $\{0 \leq x \leq M,\ |y-x| \leq \eta\}$, of area $2M\eta$. Ordinary integral Cauchy–Schwarz therefore gives

$$q_N^2 \leq 2M\eta G_N. \tag{30}$$

All integrals here are Lebesgue integrals on $\mathbb{R}^2$, and all kernels are finite sums of bounded finite-area indicators.

For fixed clock coordinates let $e_N(B) = \mathbb{E}_W S_N^2$. Write $\mathcal{W}(u, v) = X(u)X(v) - \langle u, v\rangle$. The Gaussian pairing identity gives

$$\mathbb{E}_W[\mathcal{W}(u, v)\mathcal{W}(a, b)] = \langle u, a\rangle\langle v, b\rangle + \langle u, b\rangle\langle v, a\rangle.$$

This follows by the three pairings of four centered Gaussian coordinates and cancellation of the product of means. Since $S_N = \sum_i \mathcal{W}(d_i, h_i)$, telescoping the identity

$$2\mathcal{W}(d_i, h_i) = \mathcal{W}(h_{i+1}, h_{i+1}) - \mathcal{W}(h_i, h_i) - \mathcal{W}(d_i, d_i)$$

and applying the pairing formula gives

$$e_N(B) = \frac{1}{2}\left(r_N^2 - 2\sum_i \langle h_N, d_i\rangle^2 + G_N\right).$$

Indeed the endpoint square has energy $2r_N^2$, the sum of increment squares has energy $2G_N$, and their cross term is $2\sum_i \langle h_N, d_i\rangle^2$. The map $r \mapsto \min(r_N, r)$ is 1-Lipschitz, so $\sum_i \langle h_N, d_i\rangle^2 \leq q_N \leq U_N$ for the actual clock. Hence $G_N \leq 2e_N(B) + 2U_N$. Combining with (30) yields the pathwise bound

$$q_N^2 \leq 4M\eta(e_N(B) + U_N). \tag{31}$$

We now prove $a_N = \mathbb{E}_B e_N(B) \to \infty$ without a four-time lower bound. Suppose there is a subsequence with $a_N \leq A$. The L^1 convergence of q_N supplies a further subsequence with $q_N \to T$ almost surely: select summable L^1 errors and apply Tonelli. On the common continuous clock

event, choose $M = 1 + \sup_{t\le T}|B_t| < \infty$ and $\eta_N = N^{-1} + \max_i |r_{i+1} - r_i| \to 0$. Bound (31) then forces $e_N(B) + U_N \to \infty$ on this further subsequence. Fatou applied to these nonnegative variables would give

$$\infty = \mathbb{E}\liminf(e_N(B) + U_N) \le \liminf \mathbb{E}(e_N(B) + U_N) \le A + T,$$

a contradiction. Thus no bounded-energy subsequence exists, which proves divergence to infinity along the entire sequence.

10.2 A bounded centered clock correction

The identity $2(\min(x, y) - x) = y - x - |y - x|$ gives

$$C_N = \frac{1}{2}\left(r_N - \sum_i |r_{i+1} - r_i|\right).$$

This uncentered quantity need not be bounded. Let instead

$$A_N = \sum_i (|\Delta_i B| - \mathbb{E}|\Delta_i B|), \quad L_N = \sum_i (|\Delta_i B| - |r_{i+1} - r_i|), \quad H_N = \sum_i \operatorname{sgn}(B_{t_i})\Delta_i B,$$

where $\operatorname{sgn}(0) = 0$, and put $d_N = \sum_i \mathbb{E}|\Delta_i B|$. Independent increments give $\mathbb{E}A_N^2 \le T$. The sign coefficients are measurable at the left endpoints and bounded by one; conditioning each cross term on its later left endpoint gives zero and hence $\mathbb{E}H_N^2 \le T$.

The elementary discrete Tanaka inequalities are

$$0 \le |y - x| - ||y| - |x|| \le |y| - |x| - \operatorname{sgn}(x)(y - x).$$

For equal signs the left difference vanishes and the upper expression is zero; for opposite signs both inequalities follow by inserting $x > 0 > y$ or $y > 0 > x$; at $x = 0$ the left side is zero and the right side is $|y|$. Summing yields $0 \le L_N \le r_N - H_N$. Therefore $\mathbb{E}L_N^2 \le 2\mathbb{E}r_N^2 + 2\mathbb{E}H_N^2 \le 4T$. The exact identity

$$2C_N + d_N = r_N - A_N + L_N$$

and $(u + v + w)^2 \le 3(u^2 + v^2 + w^2)$ give $\mathbb{E}(2C_N + d_N)^2 \le 18T$. Since deterministic shifts leave variance unchanged, $4\operatorname{Var}(C_N) \le 18T$, as claimed. This argument works for any deterministic increasing partition from zero, although only uniform grids are used in the escape theorem.

10.3 A detecting observable survives projection

The variable R_N is $\mathcal{O}$-measurable. Conditional scenery centering gives $\mathbb{E}S_N = 0$ and $\mathbb{E}[S_N(C_N - \mathbb{E}C_N)] = 0$, with all products integrable by Cauchy–Schwarz. Consequently

$$R_N - \mathbb{E}C_N = S_N + (C_N - \mathbb{E}C_N), \qquad \|R_N - \mathbb{E}C_N\|_2^2 = a_N + \operatorname{Var}(C_N).$$

By the defining orthogonal projection identity for conditional expectation,

$$\langle I_N, R_N - \mathbb{E}C_N\rangle = \langle S_N, R_N - \mathbb{E}C_N\rangle = a_N.$$

Cauchy–Schwarz proves the quantitative inequality in Theorem 10.1. Since $a_N \to \infty$ and the clock variance is bounded, it forces $\|I_N\|_2 \to \infty$. This proves the theorem. In particular, contraction

of the projection does not remove the divergence, and no commutation of that projection with a scenery-chaos projection has been assumed.

In the original source construction the finite field has scenery component $\sum_i X(h_i)d_i$ and zero clock component. Its ambient scenery divergence is S_N: Gaussian integration by parts gives $\delta_W(X(h_i)d_i) = \mathcal{W}(d_i, h_i)$ for fixed clock, and the finite L^2 bounds above allow fiberwise integration. Restricting the two-source derivative to its original IBM observable domain makes its adjoint value I_N. Explicitly, every test F in that original domain satisfies

$$\left\langle \sum_i X(h_i)d_i, D_W F \right\rangle_{L^2} = \langle S_N, F\rangle = \langle I_N, F\rangle.$$

This is the precise finite-graph identification recorded by the verified native-field declarations. The escape proof above was given directly for actual conditional expectations, independently of any enlargement of that test domain. It supplies neither a completed physical-time integral nor a theorem against all possible renormalizations or higher-channel constructions.

10.4 Failure of weak reconstruction on the full observable space

The norm escape has a stronger consequence than absence of a strong limit. Let $E = L^2(\mathcal{O})$, the real closed observable subspace of the actual scenery–clock probability space. For the same original projected native outputs $I_N \in E$ there exists a *single* $v \in E$ such that

$$\sup_{N\geq 1} |\langle I_N, v\rangle| = \infty.$$

Thus those pairings converge to no finite scalar. Moreover no subsequence I_{N_j} with strictly increasing indices converges weakly to any $w \in E$.

Here is the complete functional-analytic step. If every $v \in E$ had bounded pairings, the closed sets

$$A_k = \{v \in E : |\langle I_N, v\rangle| \leq k \text{ for all } N \geq 1\}, \qquad k = 1, 2, \ldots,$$

would cover E. It is a Hilbert space, hence complete. The Baire category theorem would give a ball $B(v_0, r) \subset A_k$ for some k and $r > 0$. For every $\|h\| \leq 1$, the two points $v_0 \pm (r/2)h$ belong to that ball. Subtract their pairings and use the two bounds by k to obtain

$$r|\langle I_N, h\rangle| \leq 2k \quad \text{for every } N.$$

Taking the supremum over the unit ball gives $\|I_N\| \leq 2k/r$, contrary to Theorem 10.1. Negating the assumed pointwise boundedness produces the stated fixed observable. This is the classical Banach–Steinhaus mechanism [5]; its use introduces no new probabilistic assumption.

For every prescribed strictly increasing sequence N_j, the same norm escape holds along j. Applying the argument to that subsequence gives a detecting observable $v_{(N_j)}$ with unbounded pairings. Weak convergence to any $w \in E$ would make every such pairing convergent, hence bounded, a contradiction. The detecting observable may depend on the subsequence; no one test detecting every possible subsequence is asserted.

This is a full-observable L^2 weak-topology obstruction for exactly the original finite-field adjoint outputs identified above. The argument does not put its detecting observable in the original closed derivative domain, nor make it smooth or bounded. Convergence against a smaller graph-domain test class, a distributional topology, a different integrand norm or a different compensated

prescription is not ruled out. In particular the spatial Simpson theorem and its actual pathwise primitive remain intact.

`IBMNativeWeakObstruction`:
`exists_original_unbounded_observable`,
`exists_original_nonconvergent_observable`,
`no_original_weak_limit`, `no_original_weak_subsequence`. The source uses $N+1$ cells for indices $N \geq 0$; the paper uses $N \geq 1$ cells, which is only an index shift.

11 Strong source contractions without a full observable limit

The preceding obstruction concerns the full outputs I_N. Certain fixed tests nevertheless have strong limits after averaging only over the scenery. We reconstruct these limits for actual quadratic observations and bounded C^2 terminal tests. Throughout this section the horizon is $T > 0$, there are $N \geq 1$ cells, $h = T/N$, and $t_i = ih$. Write $\mathbb{E}_W$ and $\mathbb{E}_B$ for the two source expectations. Observation times $R, S \geq 0$ are fixed and may lie inside or outside $[0, T]$.

11.1 One common observation may anticipate the clock grid

We first need a weighted Brownian bracket estimate that does not assume predictability of the weights. For bounded Borel functions $\phi_i : \mathbb{R}^2 \to \mathbb{R}$, $|\phi_i| \leq 1$, let

$$D_i = (B_{t_{i+1}} - B_{t_i})^2 - h, \qquad a_i = \phi_i(B_{t_i}, B_R), \qquad E_N = \sum_{i<N} D_i a_i.$$

Then the exact source bound is

$$\mathbb{E}_B E_N^2 \leq \frac{12T^2}{N} + \frac{16T^2}{\sqrt{N}}. \tag{32}$$

The functions may vary with the cell and with N, and the same bound even permits a different common observation time $R = R(N)$ for each grid. It is not a theorem for arbitrary functions of the entire future path.

Here is a direct proof. For $s < t \leq R$, jointly Gaussian regression onto $\mathcal{F}_s^B \vee \sigma(B_R)$ gives

$$\mathbb{E}_B[(B_t - B_s)^2 - (t-s) \mid \mathcal{F}_s^B, B_R] = \left(\frac{t-s}{R-s}\right)^2 ((B_R - B_s)^2 - (R-s)). \tag{33}$$

Indeed $B_t - B_s$ is the sum of $(t-s)(B_R - B_s)/(R-s)$ and a centered Gaussian residual independent of the conditioning variables, of variance $(t-s) - (t-s)^2/(R-s)$. Expanding the square proves the formula. Its L^2 norm is $\sqrt{2}(t-s)^2/(R-s)$, at most twice that ratio. If $R \leq s$, the corresponding conditional mean is zero by independent increments. These statements use the actual clock and its original Gaussian coupling.

`GaussianTerminalBracketPrediction.conditional_square_actual`.
`GaussianInteriorClockBracket.past_conditional_square`.

For each j set $A_j = \sum_{i<j} D_i a_i a_j$. This is measurable with respect to $\mathcal{F}_{t_j}^B \vee \sigma(B_R)$, lies in L^2, and has norm at most $2T$, because $\|D_i\|_2 = \sqrt{2}h \leq 2h$. The finite expansion is therefore legitimate:

$$\mathbb{E}_B E_N^2 = \sum_i \|D_i a_i\|_2^2 + 2\sum_j \mathbb{E}_B[D_j A_j].$$

The diagonal sum is at most $4T^2/N$. Put $m = \min(N, \lfloor R/h \rfloor)$. For $j < m$, the cell ends before R and $R - t_j \geq h(m-j)$. Conditioning as in (33) bounds the absolute row pairing by $4T^2/(N(m-j))$.

For $j > m$ it is zero; the possible row $j = m < N$ is bounded by $4T^2/N$ using Cauchy–Schwarz alone. Finally $\sum_{k=1}^{m} k^{-1} \le 2\sqrt{m} \le 2\sqrt{N}$. For example $k^{-1} \le 2(\sqrt{k} - \sqrt{k-1})$, so this bound telescopes. Summing the rows proves (32), including $R = 0$.
`GaussianAllObservationBracket.error_norm_sq_bound`.
`GaussianAllObservationBracket.varying_error_tendsto`.

11.2 Capped random levels and a genuine mixed bracket

For a nonnegative radius r put $c_r(x) = \min(|x|, r)$. Define

$$Q_N^{R,S} = \sum_{i<N} [c_{|B_R|}(B_{t_{i+1}}) - c_{|B_R|}(B_{t_i})][c_{|B_S|}(B_{t_{i+1}}) - c_{|B_S|}(B_{t_i})],$$
$$O^{R,S} = \int_0^T \mathbf{1}_{\{|B_t| < \min(|B_R|, |B_S|)\}}\, dt.$$

These are actual functions of the original clock. We will prove

$$Q_N^{R,S} \longrightarrow O^{R,S} \quad \text{in } L^2(\mathbb{P}_B). \tag{34}$$

Random radii here are not asserted to be stopping times.

First consider $R = S$. For each fixed $t \neq R$, $\mathbb{P}_B(|B_t| = |B_R|) = 0$: the event implies $B_t - B_R = 0$ or $B_t + B_R = 0$, each a nondegenerate Gaussian equality. The single time $t = R$ has zero Lebesgue measure. Fubini thus proves that the random boundary $|B_t| = |B_R|$ has zero time occupation almost surely. The origin $B_t = 0$ also has zero time occupation by the same argument. Continuity of each clock path now implies convergence of the bounded left-node Riemann samples of $\mathbf{1}_{\{|B_t| < |B_R|\}}$ to its time integral, and vanishing of samples of shrinking bands about either boundary. To justify this last statement, represent the sums by step functions on the cells. At every time outside the null boundary their left nodes tend to that time and the band indicators eventually vanish. Dominated convergence first in time and then in clock probability proves the assertions in L^2, since all sample sums lie between zero and T.
`IBMAllObservationBoundary.distance_positive_all_windows`.

Equation (32) allows replacement of h by $(\Delta B_i)^2$ in all those sums, including bands whose widths vary with N. Away from an origin band, a cap boundary band, and increments larger than a width δ, the elementary identity

$$[c_r(y) - c_r(x)]^2 = \mathbf{1}_{\{|x| < r\}}(y - x)^2$$

holds: x, y have the same sign and lie on the same side of the cap. Always both quantities lie between zero and $(y - x)^2$. Consequently their absolute difference is bounded by $(y - x)^2$ times

$$\mathbf{1}_{\{|x| \le \delta\}} + \mathbf{1}_{\{||x| - r| \le \delta\}} + \mathbf{1}_{\{|y - x| > \delta\}}.$$

The first two grid sums vanish in L^1 by the band argument. For the last one, the actual Gaussian fourth moment gives

$$\mathbb{E}_B \sum_i (\Delta B_i)^2 \mathbf{1}_{\{|\Delta B_i| > \delta\}} \le \delta^{-2} \sum_i \mathbb{E}_B (\Delta B_i)^4 = \frac{3T^2}{N\delta^2}.$$

Take $\delta = N^{-1/4}$. We have proved the capped square limit in L^1, for every observation time R.
`IBMClippedClockCrossing.crossing_bound`.

`IBMClippedClockTail.mean_bound`.
`IBMAllObservationLimits.actual_absolute_error_tendsto_all`.

For two radii, select the capped square sum corresponding to the smaller of $|B_R|, |B_S|$. Its error relative to $O^{R,S}$ is bounded by the sum of the two individual absolute errors; no independence of that selector is needed. On one cell the mixed product equals the selected square unless the cell crosses the smaller cap. In particular the difference vanishes whenever $|y - x| \le \delta$ and both $||x| - |B_R||, ||x| - |B_S||$ exceed δ. Its absolute value is always at most $2(y - x)^2$. The same two band bounds and large-increment bound therefore show $Q_N^{R,S} \to O^{R,S}$ in L^1. This comparison uses only one-observation estimates separately, not an unjustified estimate for a weight depending on two observations.
`IBMClippedClockCovariation.mixedError_crossing_bound`.
`IBMMixedClockSelection.mixed_absolute_error_tendsto`.

For the promised L^2 upgrade put $U_N = \sum_i (\Delta B_i)^2$. Both caps are 1-Lipschitz, so $|Q_N^{R,S}| \le U_N$, whereas $0 \le O^{R,S} \le T$. For real $|x| \le u$ and $|y| \le T$,

$$(x - y)^2 \le 4T|x - y| + (u - T)^2.$$

Indeed, with $d = |x - y| \le u + T$, use $2d^2 \le 2(u + T)d$ and $(d - (u - T))^2 \ge 0$ to obtain the inequality. Thus

$$\|Q_N^{R,S} - O^{R,S}\|_2^2 \le 4T\|Q_N^{R,S} - O^{R,S}\|_1 + \frac{2T^2}{N}, \tag{35}$$

where the last term is the exact variance of ordinary Brownian quadratic variation. This proves (34).
`IBMMixedClockStrongL2.mixed_error_bound`.
`IBMMixedClockStrongL2.mixedL2_tendsto`.

11.3 Actual quadratic observations select a finite clock limit

Retain the same compensated source grid S_N as before. Conditionally on the clock, Gaussian integration of a centered product gives

$$\mathbb{E}_W[(X(d_i)X(h_i) - \langle d_i, h_i\rangle)Z_R Z_S] = \langle d_i, h_R\rangle\langle h_i, h_S\rangle + \langle d_i, h_S\rangle\langle h_i, h_R\rangle.$$

This is the two nonconstant pairings in the fourth Gaussian moment; the constant pairing cancels the subtracted covariance. Since $\langle h_i, h_R\rangle = \min(|B_{t_i}|, |B_R|)$, the elementary finite product telescope yields the exact identity

$$\mathbb{E}_W[S_N Z_R Z_S] = M_N^{R,S} := c_{|B_R|}(B_T)c_{|B_S|}(B_T) - Q_N^{R,S}. \tag{36}$$

All finite products are integrable by Gaussian moments of the clock and scenery. The endpoint product is in clock L^2, since it is bounded in absolute value by $|B_T|^2$. It follows that

$$M_N^{R,S} \longrightarrow M^{R,S} := c_{|B_R|}(B_T)c_{|B_S|}(B_T) - O^{R,S} \quad \text{in clock } L^2. \tag{37}$$

For any original clock coefficient $a \in L^2(\mathbb{P}_B)$,

$$\mathbb{E}_B|a(M_N^{R,S} - M^{R,S})| \le \|a\|_2\|M_N^{R,S} - M^{R,S}\|_2 \longrightarrow 0.$$

This is an actual weighted L^1 conclusion, even for unbounded a.
`IBMRawGridMixedObservable.source_mixed_test`.

`IBMActualQuadraticL2Reconstruction.native_tendsto`.
`IBMActualQuadraticL2Reconstruction.weighted_absolute_tendsto`.

The test $Z_R Z_S$ is observable and in joint L^2. Hence the orthogonal projection defining I_N may be removed in this pairing, giving

$$\langle I_N, Z_R Z_S \rangle = \mathbb{E}_B M_N^{R,S} \longrightarrow \mathbb{E}_B M^{R,S}.$$

This removal is legitimate only inside the pairing with the observable test. It does not identify the random variable $M_N^{R,S}$ with I_N, nor assert that arbitrary clock weights are observable. Finite linear combinations of these quadratic tests follow by linearity and finite summation, without any completed-test-space membership assumption for the individual observation products.
`IBMMixedClockLimit.nativeMixedPairing_tendsto`.
`IBMActualQuadraticRandomReconstruction.finite_family_tendsto`.

11.4 Bounded C^2 tests and the actual second-jet coefficient

Let $\varphi \in C^2(\mathbb{R})$ with $\varphi, \varphi', \varphi''$ bounded. For the same fixed observation time R define the actual coefficient

$$b_\varphi(B) = \frac{1}{2}\mathbb{E}_W[\varphi''(Z_R) \mid B], \qquad |b_\varphi| \le \tfrac{1}{2}\|\varphi''\|_\infty.$$

Here the notation means integration over the independent scenery with the clock fixed; it is a genuine bounded clock L^2 class. There is no assigned intrinsic derivative of $\varphi(Z_R)$ in this definition.

For deterministic Gaussian directions d, h, k, two scalar Gaussian integrations by parts give

$$\mathbb{E}_W[(X(d)X(h) - \langle d, h\rangle)\varphi(X(k))] = \langle d, k\rangle\langle h, k\rangle \mathbb{E}_W \varphi''(X(k)). \tag{38}$$

For completeness, when $k \neq 0$ decompose d, h into their projections onto k and orthogonal residuals. The residuals are independent of $X(k)$; their covariance cancels the residual part of $\langle d, h\rangle$. The remaining term is

$$\frac{\langle d, k\rangle\langle h, k\rangle}{\|k\|^4}\mathbb{E}_W[(X(k)^2 - \|k\|^2)\varphi(X(k))].$$

Integrate twice against the scalar Gaussian density. Boundary terms vanish because the function and its first derivative are bounded, and the result is $\|k\|^4 \mathbb{E}_W \varphi''(X(k))$. All terms are integrable by the same bounds. If $k = 0$, both sides of (38) vanish directly. This also identifies the source second-chaos projection of $\varphi(X(k))$ as $\frac{1}{2}\mathbb{E}_W\varphi''(X(k))(X(k)^2 - \|k\|^2)$: testing against every centered square verifies the identity, and polarization makes their span dense in the second chaos.
`IBMFirstChaosMixedStein.second_centered_mixed`.
`IBMBoundedC2SecondProjection.projection_two`.

Apply (38) conditionally with $k = h_R$ and sum over the finite grid. The scalar square telescope has the factor two, so the exact source contraction is

$$\mathbb{E}_W[S_N \varphi(Z_R)] = b_\varphi M_N^{R,R}. \tag{39}$$

Consequently it converges in the original clock L^2 to $b_\varphi M^{R,R}$, with error at most $\frac{1}{2}\|\varphi''\|_\infty \|M_N^{R,R} - M^{R,R}\|_2$. This multiplication is justified by a bounded actual coefficient, not by an assumed product rule for a distribution. Since the terminal test is bounded and observable, it also proves

$$\langle I_N, \varphi(Z_R)\rangle \longrightarrow \mathbb{E}_B[b_\varphi M^{R,R}].$$

`IBMBoundedC2NativeReconstruction.native_actual`.
`IBMBoundedC2NativeReconstruction.native_tendsto`.
`IBMBoundedC2NativeReconstruction.intrinsic_pairing_tendsto`.

As human evaluations of the coefficient, for $\varphi(x) = \cos(\xi x)$ one has $b_\varphi = -\xi^2 e^{-\xi^2|B_R|/2}/2$; for $\varphi(x) = \sin(\xi x)$ it is zero. The latter also follows directly from scenery reflection: S_N is even and the sine test is odd. Thus good source-test convergence does not put the terminal test in the native clock derivative domain; the earlier sine-domain obstruction in the clock-domain companion remains in force.

The convergent families above are fixed tests or their explicitly controlled finite combinations. They do not include every observable L^2 test: the Baire argument above supplies tests with unbounded pairings for precisely this same sequence. No full-output weak limit, commutation of observable and chaos projections, observed-filtration integral, or physical-time Itô formula follows. The proof uses classical Gaussian regression, Wick pairing, Brownian bracket control and bounded-coefficient continuity; priority for this specific IBM source-contraction package has not been established.

12 A faithful Banach domain for clock-valued source limits

The fixed-test limits just proved do not give convergence against every observable L^2 test. A separate completed test domain does support continuous clock-L^1 limits of the original source contractions. We construct that domain explicitly, including the norm and the consistency needed to forget a test's spatial presentation.

12.1 Spatial square kernels and their precise completed domain

For $x, y \geq 0$ let $J(x,y) = (\min(x,y), \max(x,y)]$, and for a bounded continuous real function F on $\mathbb{R}^2$ put

$$C_F(x,y) = \int_{J(x,y)^2} F(u,v)\, du\, dv.$$

This rectangle is unoriented: its area is $(y-x)^2$ even when $y < x$. Thus $|C_F(x,y)| \leq \|F\|_\infty (y-x)^2$. For Lipschitz F with constant L in the maximum product metric,

$$|C_F(x,y) - F(x,x)(y-x)^2| \leq L|y-x|^3, \tag{40}$$

since each point of the square is within $|y-x|$ of (x,x). Let $\mathcal{C}$ be the supremum-norm completion of the bounded Lipschitz functions on $\mathbb{R}^2$. It embeds isometrically into $C_b(\mathbb{R}^2)$: a norm-Cauchy sequence converges uniformly to a bounded continuous function, with the same norm. This specifies the completed domain used below; we make no identification with all of C_b.
`IBMSpatialKernelTesting.cell_diagonal_error`.
`IBMSpatialCompletedTestLimit.realize`.

For the same original clock $r_i = |B_{t_i}|$, define

$$K_N F = \sum_{i<N} C_F(r_i, r_{i+1}), \qquad KF = \int_0^T F(|B_t|, |B_t|)\, dt.$$

For each deterministic $F \in \mathcal{C}$,

$$K_N F \longrightarrow KF \quad \text{in clock } L^1, \qquad \|K_N F\|_1,\ \|KF\|_1 \leq T\|F\|_\infty. \tag{41}$$

Here is a full proof on the Lipschitz core. Equation (40) bounds the expected error in replacing each cell by its diagonal value by

$$L\sum_i \mathbb{E}_B|r_{i+1}-r_i|^3 \le L\mathbb{E}|G|^3\,\frac{T^{3/2}}{\sqrt{N}}, \qquad G\sim N(0,1).$$

For the diagonal sum, replace $(\Delta r_i)^2$ by $(\Delta B_i)^2$. The resulting L^1 error is at most $\|F\|_\infty$ times the folded-clock quadratic-variation loss proved in Section 10, which tends to zero. The bounded coefficient $F(r_i, r_i)$ is measurable at t_i, so independent increments and centering give

$$\left\|\sum_i F(r_i,r_i)((\Delta B_i)^2-h)\right\|_2^2 \le \frac{2T^2\|F\|_\infty^2}{N}.$$

Cross-cell terms vanish by conditioning on the later left past; the diagonal conditional variance is $2h^2$. The remaining Riemann sum converges pathwise to KF by continuity and in L^1 by its deterministic bound $T\|F\|_\infty$. This proves convergence. The bounds in (41) follow from $|K_N F| \le \|F\|_\infty \sum(\Delta r_i)^2$, $\mathbb{E}\sum(\Delta r_i)^2 \le T$, and the time-integral bound. For $F \in \mathcal{C}$, choose core $F_j \to F$ uniformly. The two uniform operator bounds make the approximation error at most $2T\|F-F_j\|_\infty$, independently of N; first fix j, take $N\to\infty$, then $j\to\infty$. This proves the completed assertion and retains the displayed actual formulas.

`IBMSpatialLipschitzTestLimit.core_tendsto`.

`IBMSpatialCompletedTestLimit.completed_tendsto`.

12.2 Random completed tests need a separate extension argument

Set $\mathcal{R} = L^2(\mathbb{P}_B;\mathcal{C})$, the actual Bochner space of almost everywhere strongly measurable random tests. For $u \in \mathcal{R}$, define the pointwise expressions

$$K_N[u](B) = \sum_i C_{u(B)}(r_i, r_{i+1}), \qquad K[u](B) = \int_0^T u(B)(|B_t|,|B_t|)\,dt.$$

No independence between $u(B)$ and the clock is imposed. These evaluations are measurable. Indeed evaluation of either pathwise continuous linear functional at each fixed F is measurable in B, by the actual rectangle/time integrals. A strongly measurable version of u has simple-function approximations converging pointwise in $\mathcal{C}$; continuity of the functional then proves measurability of its evaluation at $u(B)$. Almost everywhere changes do not affect the resulting scalar classes. This argument requires no strong measurability of the random functional in the full norm-dual.

Let $U_N = \sum_i (\Delta B_i)^2$. Pathwise,

$$|K_N[u](B)| \le U_N(B)\|u(B)\|_\infty, \qquad |K[u](B)| \le T\|u(B)\|_\infty.$$

The original Gaussian moments give $\|U_N\|_2 = T\sqrt{1+2/N} \le 3T$. Cauchy–Schwarz yields actual L^1 classes and uniform bounds

$$\|K_N[u]\|_1,\ \|K[u]\|_1 \le 3T\|u\|_{\mathcal{R}}. \tag{42}$$

They also prove continuity and linearity in the actual equivalence classes.

`IBMWeakRandomEvaluation.evaluation_measurable`.

`IBMSpatialRandomTestBounds.finite_norm_bound`.

For a finite-valued random test $u = \sum_{j=1}^{m} \mathbf{1}_{A_j} F_j$ on disjoint measurable sets, its error is bounded by

$$\|K_N[u] - K[u]\|_1 \le \sum_{j=1}^{m} \|K_N F_j - K F_j\|_1 \to 0.$$

The events A_j may depend on the entire clock. Such simple functions are dense in the Bochner L^2 space. For arbitrary u, approximate it there by one simple function v. Equations (42) give total approximation error $6T\|u - v\|_{\mathcal{R}}$, uniformly in N. The fixed simple-function error tends to zero, proving

$$K_N[u] \to K[u] \quad \text{in clock } L^1 \quad (u \in \mathcal{R}). \tag{43}$$

Deterministic-test convergence alone would not justify this step; the common L^2 random bounds and the finite-valued approximation are essential to the argument.

`IBMSpatialRandomTestConvergence.random_tendsto`.

For later use put

$$H_N[u](B) = C_{u(B)}(0, |B_T|) - K_N[u](B), \qquad H[u](B) = C_{u(B)}(0, |B_T|) - K[u](B).$$

The endpoint term is bounded by $|B_T|^2 \|u(B)\|_\infty$, and $\||B_T|^2\|_2 = \sqrt{3}T \le 3T$. Hence

$$H_N[u] \to H[u] \text{ in } L^1, \qquad \|H_N[u]\|_1,\ \|H[u]\|_1 \le 6T\|u\|_{\mathcal{R}}. \tag{44}$$

The symbol H here denotes the difference of two real spatial actions; it changes no probability law. Its pathwise formula is a bounded functional on spatial tests, with bound $(|B_T|^2 + T)\|F\|_\infty$.

`IBMSpatialSignedRandomTests.signed_random_tendsto`.

`IBMSpatialPathReconstruction.signedPath_bound`.

12.3 The original joint tests and their spatial symbols

Use the physical scenery isometry X on $L^2(\mathbb{R})$ fixed by the original two-sided Brownian indicator coordinates; on $\mathbf{1}_{(0,r]}$, $r \ge 0$, it agrees with the scenery W_r already used. The clock-domain companion reconstructs this same-source isometry; no new independent noise is introduced. For real Schwartz functions g, k, write $X_g = X(g')$, $X_k = X(k')$. The exact overlap is

$$\langle \mathbf{1}_{(0,r]}, g' \rangle = g(r) - g(0), \qquad r \ge 0,$$

by the fundamental theorem of calculus. Put

$$q_{g,k,a}(W, B) = a(B) X_g(W) X_k(W), \qquad u_{g,k,a}(B)(x, y) = a(B) g'(x) k'(y), \quad a \in L^2(\mathbb{P}_B).$$

The joint test belongs to $\mathcal{E} = L^2(\mathbb{P}_W \otimes \mathbb{P}_B)$: the fixed Gaussian product $X_g X_k$ has finite second moment and is independent of a. The symbol belongs to $\mathcal{R}$, since $g'(x)k'(y)$ is bounded Lipschitz and its random norm is a constant times $|a(B)|$. These are original joint tests; intrinsic observable membership is an additional condition, not part of the definition.

`IBMPhysicalWickTestDomain.source_primitive`.

`IBMJointMixedSpatialTransport.jointTest_actual`.

For any $q \in \mathcal{E}$ define the finite source contraction $\mathsf{C}_N q = \mathbb{E}_W[S_N q \mid B]$, where this again means scenery integration with clock fixed. Fubini and Cauchy–Schwarz give

$$\|\mathsf{C}_N q\|_1 \le \|S_N\|_2 \|q\|_2.$$

It is therefore a continuous linear map $\mathcal{E} \to L^1(\mathbb{P}_B)$ for each fixed N; no uniform bound in this ambient norm is asserted. Gaussian Wick pairing and the same finite product telescope as before give the exact identity

$$\mathsf{C}_N q_{g,k,a} = H_N[u_{g,k,a}]. \tag{45}$$

To check the spatial side, the actual rectangle integral factors as

$$C_{g'\otimes k'}(x,y) = (g(y)-g(x))(k(y)-k(x)).$$

For $y < x$ both one-dimensional integrals reverse sign, leaving the same product. The endpoint term is $(g(|B_T|)-g(0))(k(|B_T|)-k(0))$, and subtracting the cell sum gives exactly the Gaussian overlap telescope. This also verifies that no extra factor $1/2$ belongs in (45).

`IBMJointRandomContraction.joint_bound`.

`IBMSpatialMixedWickTesting.cell_mixed`.

`IBMJointRandomGraphIdentity.generator_identity`.

12.4 Complete paired presentations, then quotient the zero joint tests

Give $\mathcal{E} \times \mathcal{R}$ the maximum product norm, and let $\mathcal{G}$ be the closed linear span of all the pairs $(q_{g,k,a}, u_{g,k,a})$. It is a Banach space. Denote its two continuous coordinate maps by j and s, so $\|jp\|_2, \|sp\|_{\mathcal{R}} \le \|p\|_{\mathcal{G}}$. For each fixed N, both sides of (45) are continuous on this product. They agree on the generating pairs, hence on their finite linear span and its closure. Consequently for every $p \in \mathcal{G}$,

$$\mathsf{C}_N(jp) = H_N[sp] \longrightarrow H[sp] \quad \text{in clock } L^1. \tag{46}$$

This extends the *random* identity, before taking expectations.

`IBMJointCompletedTestGraph.testGraph`.

`IBMJointRandomGraphIdentity.finite_identity`.

If $jp = 0$, every finite left side of (46) is zero. Uniqueness of its L^1 limit then forces $H[sp] = 0$ as an actual L^1 class. More generally two presentations with the same joint test have the same limiting clock class. No injectivity of the spatial symbol map is assumed. The closed kernel $\mathcal{K} = \ker j$ can therefore be quotiented out. Define

$$\mathcal{T} = \mathcal{G}/\mathcal{K}, \qquad \|[p]\|_{\mathcal{T}} = \inf_{k\in\mathcal{K}} \|p+k\|_{\mathcal{G}}.$$

It is Banach, and the induced realization $J : \mathcal{T} \to \mathcal{E}$, $J[p] = jp$, is injective and satisfies $\|Jx\|_2 \le \|x\|_{\mathcal{T}}$. Indeed $J[p] = 0$ means $p \in \mathcal{K}$, exactly the zero quotient class; the norm estimate follows by taking the infimum of $\|j(p+k)\|_2 \le \|p+k\|$. Thus these tests are faithful original joint random variables, with their specified quotient norm.

The limiting map

$$\mathsf{D}_T : \mathcal{T} \to L^1(\mathbb{P}_B), \qquad \mathsf{D}_T[p] = H[sp]$$

is well-defined by the kernel argument. For every $x \in \mathcal{T}$,

$$\mathsf{C}_N(Jx) \longrightarrow \mathsf{D}_T x \quad \text{in clock } L^1, \qquad \|\mathsf{C}_N(Jx)\|_1,\ \|\mathsf{D}_T x\|_1 \le 6T\|x\|_{\mathcal{T}}. \tag{47}$$

For the bounds, apply (44) to $s(p+k)$ for every $k \in \mathcal{K}$; the finite and limiting values are independent of k. Infimizing proves the inequalities. Continuity and linearity follow, and pointwise uniqueness follows from uniqueness of L^1 limits. On a generator its value is the actual formula

$$\mathsf{D}_T[q_{g,k,a}, u_{g,k,a}](B) = a(B)\left[(g(|B_T|)-g(0))(k(|B_T|)-k(0)) - \int_0^T g'(|B_t|)k'(|B_t|)\,dt\right].$$

The brackets here denote the corresponding quotient class of the pair, not a class of arbitrary original joint functions.

`IBMJointBanachTestSpace.realize_injective`.

`IBMJointRandomDistribution.native_tendsto`.

`IBMJointRandomDistribution.native_bound`.

`IBMJointRandomDistribution.distribution_bound`.

Taking clock expectation is a continuous map on L^1, so it recovers the scalar limiting functional. If Jx independently belongs to the intrinsic observable space, then $\langle I_N, Jx\rangle = \mathbb{E}_B \mathsf{C}_N(Jx)$, and hence these pairings converge to $\mathbb{E}_B \mathsf{D}_T x$. This does not assign observable membership to every joint generator.

`IBMJointRandomDistribution.expectation_eq`.

`IBMJointBanachTestSpace.intrinsic_native_tendsto`.

Finally the realized test range omits some intrinsic observables. Choose the fixed observable v whose pairings with I_N are unbounded, as constructed by the earlier Baire argument. If $v = Jx$, (47) and the last pairing identity would make those pairings converge to a finite scalar, a contradiction.

`IBMJointBanachTestSpace.exists_intrinsic_outside_range`. This is a genuine completed test-space result with a controlled stronger norm. It does not identify that range with all original observable L^2, prove that it is a closed subspace there, or put the fixed quadratic and bounded C^2 observations of Section 11 in $\mathcal{T}$. Those observations were handled separately without such membership assumptions. The construction uses classical completion, Bochner approximation and closed-kernel quotient arguments; specific priority for the actual IBM test-space package remains unresolved.

13 What this integration result means

The targets are genuine pathwise finite spatial or time integrals, and the approximations converge under the actual positive scenery–clock law. Here "oriented" refers to the signed spatial increment $y - x$, not to a signed probability measure for Z. This distinguishes the present setting from the higher-order signed heat pseudoprocess calculus.

The proof is local in the spatial range. It admits unbounded global C^3 one-forms and C^4 spatial slices because every continuous sample path has a compact range. It provides convergence in probability, not integrability of arbitrary unbounded targets. Neither expectation convergence nor strong L^2 convergence has been asserted without moment hypotheses.

It also does not construct a general physical-time adapted stochastic integral. Its spatial one-form depends only on the current spatial argument, and its time–space integrand comes from a deterministic primitive F. The rule has not been identified with the original intrinsic divergence or with an operator acting on arbitrary predictable integrands. Bounded random horizons are handled by approximation, not a martingale stopping theorem. Nonsmooth Itô–Tanaka formulas and arbitrary partitions have their own proof obligations.

The native compensated escape concerns a specific projected finite-field sequence, not the symmetric Simpson prescription or every admissible integrand.

The variation and raw-channel results above are independent of the Simpson proof, which needs only quartic tightness. Intrinsic graph-dual obstructions remain separately scoped; no stochastic-domain assertion is used to justify a deterministic quadrature estimate.

14 Formal correspondence and preliminary comparisons

The frozen mathematical baseline is

`bab688bb6140cd6316c80680095a914ea0d346db.`

The namespace prefix is

`OperatorDerivativeRewrite.Foundation`

The companion `theorem-map.json` contains exact declaration links, source line numbers and hashes.

Paper result	*Lean namespace and declaration*
Compact spatial estimate	`IBMSimpsonPeano.compact_error_bound`
Spatial convergence	`IBMActualSimpson.sums_tendstoInMeasure`
Uniform spatial prefixes	`IBMUniformSimpsonProcess.maximum_tendstoInMeasure`
Random horizons	`IBMSimpsonRandomHorizon.stopped_sums_tendstoInMeasure`
Time–space compact estimate	`IBMTimeSpacePeano.compact_error_bound`
Time quadrature	`IBMTimeSpaceQuadrature.increments_tendsto`
Time–space chain	`IBMTimeSpaceSimpson.sums_tendstoInMeasure`
Uniform time–space prefixes	`IBMUniformTimeSpace.maximum_tendstoInMeasure`
Time weights	`IBMTimeWeightedSimpson.weighted_tendstoInMeasure`
Quartic variation	`IBMUniformQuarticVariation.quartic_tendstoInMeasure`
Quadratic divergence	`IBMActualQuadraticDivergence.quadratic_lower_tail`
Raw first channel	`IBMActualQuadraticDivergence.first_no_finite_limit`

The continuous-path statements are `IBMContinuousSimpsonRealization.approximation_error_bound` and `approximation_tendstoInMeasure` in the same namespace.

The formal proof uses its verified quartic-variation machinery to control compact Peano errors. The human proof above reorganizes the same application using the weaker direct Gaussian moment bound (5). That bound is proved explicitly here; no separate named Lean declaration for this shortened argument is claimed. The endpoints, smoothness assumptions and probability topologies of the principal convergence statements are those in the formal declarations. The non-separable chain theorem in the source does not require joint continuity of F as a separate input; the continuous-path declaration retains it explicitly.

The baseline receipt records a complete successful Lean build and 54,943 declarations audited with only the standard three axioms. This editorial work changes no mathematical source and does not claim a new full build. Source hashes, baseline evidence, compiler diagnostics and visual-review receipts accompany the paper. The earlier signed-Simpson working note remains intact; this paper replaces its mechanism summaries with proofs for the focused quadrature and continuous-path scope.

The primary comparison [2] constructs symmetric integrals for a two-sided-scenery IBM using an induced random partition. It is not an identification with the reflected-clock convention or

deterministic grids used here. Harnett and Nualart [3] study the critical fractional Brownian Simpson problem and distributional limits. These are close antecedents for numerical prescriptions, but their process and limiting regime differ. The compact-remainder method itself is elementary; this paper does not turn its formal verification into a novelty claim. For the compensated-grid comparison, Liu's primary manuscript [4], introduction and Hypothesis 1.1, assumes a continuous Gaussian driving process. Our Z is not Gaussian: for $t > 0$, $\mathbb{E}Z_t^2 = \sqrt{2t/\pi}$ and $\mathbb{E}Z_t^4 = 3t$, whereas a centered Gaussian of that variance would have fourth moment $6t/\pi$. Conditional Gaussian scenery does not make these marginal laws Gaussian. Thus a general Gaussian Wick–Riemann convergence theorem cannot be imported solely from the covariance of Z. This is a bounded nonapplication check, not a novelty certification for the explicit projected escape theorem. The full weak/strong partition comparison remains separate.

Priority for the combined domain, prefix and path-space package requires further targeted comparison.

References

[1] K. Burdzy, Variation of iterated Brownian motion, *CRM Proceedings and Lecture Notes* **5** (1994), 35–53. University of Washington record.

[2] D. Khoshnevisan and T. M. Lewis, Stochastic calculus for Brownian motion on a Brownian fracture, *Ann. Appl. Probab.* **9** (1999), 629–667. `https://www.math.utah.edu/~davar/ps-pdf-files/ibm.pdf`.

[3] D. Harnett and D. Nualart, On Simpson's rule and fractional Brownian motion with $H = 1/10$, *J. Theor. Probab.* **28** (2015), 1651–1688. doi:10.1007/s10959-014-0552-1.

[4] Y. Liu, *Riemann–Skorohod and Stratonovich integrals for Gaussian processes*, primary author manuscript, introduction and Hypothesis 1.1. Author-hosted PDF.

[5] S. Banach and H. Steinhaus, *Sur le principe de la condensation de singularités*, Fundamenta Mathematicae **9** (1927), 50–61. `https://doi.org/10.4064/fm-9-1-50-61`. Publisher abstract and bibliographic record inspected; original full-text retrieval was unsuccessful. The needed Baire argument is proved above.