Gaussian clock regularity and domain obstructions for reflected-clock Brownian motion

Project Stardust

Reviewed editorial draft, October 29, 2026

Abstract

Let B, W be independent Brownian motions and $Z_t = B(|W_t|)$. At each positive time the actual point observable belongs to the scenery Gaussian Sobolev domain, with an explicit finite-energy gradient, but does not belong to the original clock Gaussian Sobolev domain. The bounded smooth observable $\sin Z_t$ has the same domain separation, so unbounded point growth is not the cause. The obstruction is proved in the scenery-valued Hilbert space: its clock section has squared increment norm proportional to distance, incompatible with a common locally finite-energy weak derivative. A conditional contraction of the full clock gradient would recover precisely such a derivative, so separate scalar identities cannot evade the obstruction. An exact correlated-clock noise formula further gives a sharp weighted integral-energy cutoff. The actual endpoint belongs to the closed clock Bessel domain of order s exactly when $s < 1/2$. An exact bridge identifies the first Bessel domain with the original clock derivative domain; smoothing by order 2β reaches that domain exactly when $\beta > 1/4$. At some positive observation time, all sufficiently large real smoothing orders nevertheless stay a uniform positive L^2 distance from every observed right-past variable. For a closed densely defined Hilbert operator, strong value approximation of a point outside its domain forces all derivative norms to escape to infinity, while the embedded derivatives converge in the dual of its adjoint graph with an explicit error bound. Applied to the ambient clock operator, this gives a nonrepresentable graph-dual derivative of Z_t. The conclusions concern the stated original domains and do not supply an observed-filtration stochastic integral or a physical-time Itô formula. The proofs reconstruct frozen Lean results using classical Gaussian and closed-operator methods; novelty remains unresolved.

1 The actual process and the two Gaussian domains

On a product of independent Wiener probability spaces $(\Omega_B, \mathbb{P}_B)$ and $(\Omega_W, \mathbb{P}_W)$, let B_s, W_t, $s, t \geq 0$, be standard Brownian motions starting at zero. We study

$$Z_t(\omega_B, \omega_W) = B_{|W_t(\omega_W)|}(\omega_B). \tag{1}$$

This is the simple Brownian-time Brownian motion of Allouba–Zheng [1]. The name "reflected-clock" specifies that the same scenery B is used at every visit to a given nonnegative clock level. It does not mean an excursion-by-excursion replacement of scenery or a two-sided scenery evaluated at the signed clock.

Set

$$E = L^2(\mathbb{P}_B \otimes \mathbb{P}_W), \qquad H_B = H_W = L^2(\mathbb{R}_+; ds).$$

The isonormal coordinates of B and W identify their Cameron–Martin derivative directions with H_B and H_W. Write

$$D_B : D(D_B) \subset E \longrightarrow L^2(\Omega_B \times \Omega_W; H_B), \qquad D_W : D(D_W) \subset E \longrightarrow L^2(\Omega_B \times \Omega_W; H_W)$$

for the original closed partial Gaussian derivatives. The other source is held as an L^2 parameter in each case; these are not derivatives in physical time t. Section 2 specifies their closure and fiberwise domains precisely.

Theorem 1.1. *For every $T > 0$, $Z_T \in D(D_B)$ and*

$$D_B Z_T = \mathbf{1}_{[0,|W_T|]} \quad \text{in } H_B, \qquad \|D_B Z_T\|_2^2 = \|Z_T\|_2^2 = \sqrt{2T/\pi}. \tag{2}$$

In contrast, $Z_T \notin D(D_W)$. Consequently the point is outside any joint derivative whose domain is contained in $D(D_B) \cap D(D_W)$, including the original joint derivative restricted to IBM-observable L^2 classes in the mapped development. At $T = 0$ both derivatives of $Z_0 = 0$ exist and are zero.

Theorem 1.2. *For every $T > 0$, $S_T = \sin Z_T$ is a bounded actual observable in $D(D_B)$, with*

$$D_B S_T = \cos Z_T \, \mathbf{1}_{[0,|W_T|]}, \qquad \|D_B S_T\|_2^2 \le \sqrt{2T/\pi}.$$

Nevertheless $S_T \notin D(D_W)$ and it is outside the original joint intrinsic domain. Thus that domain does not contain all bounded smooth functions of actual point observations.

We will also prove the following quantitative form of the distinction between the original graph and a weak test-dual extension.

Theorem 1.3. *Let $\mathcal{T} = D(D_W^*)$ with norm $\|v\|_{\mathcal{T}}^2 = \|v\|_2^2 + \|D_W^* v\|_2^2$. For either $f = Z_T$ or $f = S_T$ there is a bounded clock graph-dual derivative*

$$d_W f(v) = \mathbb{E}[f D_W^* v], \qquad \|d_W f\|_{\mathcal{T}^*} \le \|f\|_2.$$

For $T > 0$ no $V \in L^2(\Omega_B \times \Omega_W; H_W)$ represents it as $\mathbb{E}\langle V, v\rangle_{H_W}$ for all $v \in \mathcal{T}$. Whenever $u_n \in D(D_W)$ and $u_n \to f$ in E,

$$\|D_W u_n\|_2 \longrightarrow \infty, \tag{3}$$

$$\|j D_W u_n - d_W f\|_{\mathcal{T}^*} \le \|u_n - f\|_2 \longrightarrow 0, \tag{4}$$

where $jV(v) = \mathbb{E}\langle V, v\rangle_{H_W}$.

This theorem concerns the ambient original clock operator. Its tests can depend on both sources and are not identified with temporal predictable integrands or with the natural filtration of Z. The native point-domain obstruction in Theorem 1.1 is a separate consequence of domain inclusion.

2 Closed Gaussian derivatives and their fiberwise lift

Here are the Gaussian foundations used in the proof. For a real separable Hilbert space H, let $X(h)$ be an isonormal Gaussian family, with $\mathbb{E}X(h)X(k) = \langle h, k\rangle_H$. On finite polynomial cylinders define

$$D\, p(X(h_1), \dots, X(h_n)) = \sum_{i=1}^{n} (\partial_i p)(X(h_1), \dots, X(h_n)) h_i. \tag{5}$$

One may use orthonormal h_i after a finite linear change of coordinates. Polynomial cylinders are dense in Gaussian L^2. For completeness, in finite dimension if f is orthogonal to every polynomial, the entire transform $\int f(x)e^{z\cdot x}\,d\gamma_n(x)$ has all Taylor coefficients zero. Cauchy–Schwarz and Gaussian exponential moments justify its entireness and termwise differentiation. Its restriction to imaginary z is the Fourier transform of $f\gamma_n$, so Fourier uniqueness gives $f = 0$. In countably many coordinates, conditional expectations onto the first n coordinates converge in L^2, reducing density to the finite case.

Gaussian integration by parts gives, for polynomial cylinders p, a and a finite coordinate vector h,

$$\mathbb{E}[\langle Dp, h\rangle a] = \mathbb{E}[p\{X(h)a - D_h a\}]. \tag{6}$$

This proves closability of (5). Indeed if $p_n \to 0$ and $Dp_n \to V$ in their L^2 spaces, the right side tends to zero for each fixed a, h. Such scalar-polynomial vector tests are dense in $L^2(\Omega; H)$, so $V = 0$. The original Gaussian derivative is the closure of this graph. Its domain is dense by the preceding density argument. Smooth compactly supported coordinate tests also satisfy (6), by ordinary finite-dimensional integration by parts and passage through the closed graph. These standard source-space constructions are the setting of Gaussian Malliavin calculus; see [2] for the corresponding Gaussian-process conventions.

For an independent parameter probability space (S, ρ), the partial derivative on $L^2(S; L^2(\Omega))$ has the exact domain

$$\begin{aligned} D(\widetilde{D}) = \{f :&\text{there exists } V \in L^2(S; L^2(\Omega; H)) \text{ with} \\ &(f(s), V(s)) \in \operatorname{graph} D \text{ for } \rho\text{-almost every } s\}. \end{aligned} \tag{7}$$

The output V is unique, since the graph has a unique value in each fiber. This is the actual Bochner graph lift, not a definition using independently chosen scalar derivatives.

The lift is closed: from a convergent sequence of both graph coordinates, take a common subsequence converging almost everywhere in both fiber norms, and use closedness of the original graph in each fiber. It is densely defined because simple S-functions with values in $D(D)$ are dense in the Bochner L^2 space. It is also the closure of tensor cylinders. To see the nontrivial inclusion, a graph pair in (7) is a strongly measurable, square-integrable function into the closed Hilbert graph of D. Approximate it by simple graph-valued functions, and approximate each of their finitely many values by a polynomial graph pair of D. This yields tensor-cylinder approximation in both graph coordinates. No measurable selection of derivative values is required: the given pair already supplies the measurable output.

Taking $S = \Omega_W$ or Ω_B defines D_B and D_W as used above. Identifying product scalar L^2 with either Bochner order is an isometric Fubini identification; changing the order does not change the product law or the graph condition.

3 Finite scenery energy at a random clock level

Let $I_B : H_B \to L^2(\mathbb{P}_B)$ be the Wiener integral, so $B_s = I_B(\mathbf{1}_{[0,s]})$. Fix $s \geq 0$ and an orthonormal basis (e_j) of H_B. The partial sums

$$B_s^{(N)} = \sum_{j=1}^{N} \langle \mathbf{1}_{[0,s]}, e_j\rangle I_B(e_j)$$

are polynomial cylinders, with gradients equal to the orthogonal projections of $\mathbf{1}_{[0,s]}$. Wiener isometry and Parseval show convergence of both values and gradients. Thus B_s is in the original scenery domain, with derivative $\mathbf{1}_{[0,s]}$.

Now put $s = |W_T|$. The functions of the clock multiplying $I_B(e_j)$ are square integrable, because

$$|\langle \mathbf{1}_{[0,|W_T|]}, e_j \rangle|^2 \le |W_T|.$$

They are allowed L^2 parameters in the scenery lift. The random projected gradients converge to $\mathbf{1}_{[0,|W_T|]}$ in the full product L^2 field norm, by pointwise Parseval and domination by $|W_T|$. The same is true of the random partial-sum values by conditioning on W. Closedness proves the scenery part of Theorem 1.1. Conditioning also gives

$$\mathbb{E} Z_T^2 = \mathbb{E}|W_T| = \sqrt{2T/\pi};$$

the last equality is the elementary absolute first moment of a centered Gaussian of variance T. This establishes every identity in (2).

4 A Hilbert-valued section with Brownian metric

Set $\mathcal{H} = L^2(\mathbb{P}_B)$ and, for $c > 0$, define

$$F_c(x) = B_{c|x|} \in \mathcal{H}, \qquad x \in \mathbb{R}. \tag{8}$$

It is a continuous Hilbert-valued curve. For nonnegative x, y, Wiener isometry gives the exact metric

$$\|F_c(x) - F_c(y)\|_{\mathcal{H}}^2 = c|x - y|. \tag{9}$$

In particular $F_c \in L^2(\gamma; \mathcal{H})$ for standard one-dimensional Gaussian measure γ, since $\|F_c(x)\|_{\mathcal{H}}^2 = c|x|$.

Lemma 4.1. *If a continuous Banach-valued curve F on $[a, b]$, $a < b$, satisfies $\|F(x) - F(a)\|^2 \ge c(x - a)$ for $c > 0$, then it has no representation*

$$F(x) = v + \int_a^x g(s)\, ds$$

almost everywhere on $[a, b]$ with $g \in L^2([a, b])$ and a constant vector v.

Proof. The primitive is continuous, so almost-everywhere equality with the continuous F implies equality everywhere. At a this gives $v = F(a)$. Cauchy–Schwarz for the Bochner integral gives, for $x > a$,

$$c(x - a) \le \|F(x) - F(a)\|^2 \le (x - a) \int_a^x \|g(s)\|^2\, ds.$$

After division by $x - a$, the right side tends to zero as $x \downarrow a$, a contradiction. □

Gaussian weighting does not remove this local obstruction: on a compact interval its density is bounded above and bounded away from zero, so Gaussian L^2 derivative energy implies Lebesgue L^2 energy there.

Lemma 4.2. *Let $\mathcal{H}$ be a real separable Hilbert space and $F, g \in L^2(\gamma; \mathcal{H})$. Suppose every scalar pair $\langle e, F\rangle, \langle e, g\rangle$, $e \in \mathcal{H}$, satisfies the closed one-dimensional Gaussian derivative identity. Then on every finite interval $[a, b]$,*

$$F(x) = v + \int_a^x g(s)\, ds$$

almost everywhere, for some $v \in \mathcal{H}$.

Proof. The scalar graph identity implies

$$\int \langle e, F(x)\rangle \{xp(x) - p'(x)\}\, d\gamma(x) = \int \langle e, g(x)\rangle p(x)\, d\gamma(x)$$

for compactly supported smooth p. If φ is a compactly supported test function in (a, b), take $p = \varphi/\rho$, where ρ is the Gaussian density. Since $\rho' = -x\rho$, $xp - p' = -\varphi'/\rho$. Thus $\langle e, g\rangle$ is the ordinary weak derivative of $\langle e, F\rangle$ on (a, b). Both are locally L^2, so the one-dimensional weak fundamental theorem gives $\langle e, F(x)\rangle = c_e + \langle e, \int_a^x g(s)\, ds\rangle$ almost everywhere. This elementary theorem follows, for example, by subtracting the displayed primitive and observing that a locally integrable distribution with zero derivative is constant; convolution with compact smooth kernels proves that assertion directly.

Choose a countable dense sequence (e_j) in $\mathcal{H}$. Intersect their full-measure equality sets and choose one point x_0 in the intersection. The vector $v = F(x_0) - \int_a^{x_0} g(s)\, ds$ realizes every constant c_{e_j}. At any other point in the intersection, the inner products with all e_j of $F(x) - v - \int_a^x g$ vanish. Density makes the vector itself zero, proving the statement. □

Combining (9) on $[0, 1]$ with the two lemmas shows that F_c has no single $g \in L^2(\gamma; \mathcal{H})$ realizing all scalar closed Gaussian derivative pairs. This is a common-field obstruction. It does not assert that every individual scalar weak derivative fails to exist.

5 Recovering a common field from the full clock graph

We now connect the section obstruction to the actual infinite-dimensional clock operator. Let

$$h_T = T^{-1/2}\mathbf{1}_{[0,T]} \in H_W, \qquad G = W(h_T) = W_T/\sqrt{T}.$$

Then $\|h_T\| = 1$ and G is standard Gaussian. Complete h_T to an orthonormal basis of H_W. The clock Gaussian space decomposes into the coordinate G and an independent residual Gaussian family. Under this decomposition,

$$Z_T = F_{\sqrt{T}}(G) \tag{10}$$

as an actual product L^2 class, with F valued in the scenery Hilbert space.

Suppose, for contradiction, that $Z_T \in D(D_W)$, with full clock gradient V. Contract V with h_T and average only over the residual clock coordinates:

$$g(x)(\omega_B) = \mathbb{E}_{\text{residual}}\langle V(\omega_B, x, \cdot), h_T\rangle_{H_W}. \tag{11}$$

Hilbert-valued conditional integration and Jensen give

$$g \in L^2(\gamma; \mathcal{H}), \qquad \|g\|_{L^2(\gamma;\mathcal{H})} \le \|V\|_2.$$

This is one common scenery-valued field, rather than separately chosen scalar derivatives.

For $e \in \mathcal{H}$ and $p \in C_c^\infty(\mathbb{R})$, the clock test field $e(\omega_B)p(G)h_T$ has divergence

$$e(\omega_B)\{Gp(G) - p'(G)\}.$$

The formula follows on tensor polynomial cylinders from Gaussian integration by parts. Its test field and displayed divergence are in the relevant L^2 spaces; passage through the closed clock graph extends the identity to Z_T, V. Using (10) and conditional integration in (11), it becomes

$$\int \langle e, F_{\sqrt{T}}(x)\rangle\{xp(x) - p'(x)\}\, d\gamma(x) = \int \langle e, g(x)\rangle p(x)\, d\gamma(x).$$

The proof of Lemma 4.2 therefore supplies a common vector primitive for $F_{\sqrt{T}}$ on $[0,1]$. Lemma 4.1 excludes it by (9). This proves $Z_T \notin D(D_W)$.

The original joint intrinsic derivative in the mapped library is defined on the intersection of the scenery and clock domains, restricted to the closed subspace of actual IBM-observable L^2 classes. The point Z_T is observable by definition but fails the clock domain test. Thus the joint-domain conclusion requires no assertion about a new derivative domain or a new filtration. This completes Theorem 1.1.

6 A bounded smooth observable with the same obstruction

The preceding argument is not restricted to unbounded point values. We prove Theorem 1.2 using an exact first-chaos projection. Let Π be the orthogonal projection in $\mathcal{H} = L^2(\mathbb{P}_B)$ onto the closed first Gaussian chaos $I_B(H_B)$. For $s \geq 0$,

$$\Pi \sin B_s = e^{-s/2} B_s. \tag{12}$$

Indeed, for $h \in H_B$, joint Gaussian regression and one-dimensional integration by parts give

$$\mathbb{E}[I_B(h) \sin B_s] = \langle h, \mathbf{1}_{[0,s]}\rangle \mathbb{E} \cos B_s = e^{-s/2}\langle h, \mathbf{1}_{[0,s]}\rangle.$$

For $s > 0$, regression here follows simply by splitting h into its projection onto $\mathbf{1}_{[0,s]}$ and its orthogonal complement; the latter Gaussian is independent and centered. At $s = 0$ the identity is zero. The last expression is the inner product with $e^{-s/2}B_s$, proving (12).

For $c > 0$ set $S_c(x) = \sin B_{c|x|}$ as a scenery-valued Hilbert class. It is continuous and has norm at most one. Its projected curve is

$$\Pi S_c(x) = e^{-c|x|/2} B_{c|x|}, \qquad \Pi S_c(0) = 0.$$

On $0 \leq x \leq 1$, the exact source variance implies

$$\|\Pi S_c(x) - \Pi S_c(0)\|_{\mathcal{H}}^2 = e^{-cx} cx \geq ce^{-c} x. \tag{13}$$

If S_c had a common finite-energy vector primitive, applying the bounded operator Π to its Bochner integral would produce a primitive of ΠS_c with no larger derivative energy. Lemma 4.1 excludes this by (13). The recovery argument of the preceding section, with F_c replaced by S_c, would recover just such a primitive from a full clock gradient of $\sin Z_T$. Thus $\sin Z_T \notin D(D_W)$ at every positive time, and domain inclusion gives the joint intrinsic exclusion.

For the positive scenery statement, at each deterministic s the Taylor polynomials of $\sin B_s$ and their derivatives converge in Gaussian L^2, dominated by a constant times $e^{|B_s|}$. Their gradients converge to $\cos B_s\, \mathbf{1}_{[0,s]}$. Closedness gives that actual scenery graph pair. At the random level $s = |W_T|$ its field norm squared is bounded by s, and its value is bounded by one. Measurability follows from continuity of B and the interval indicator map. The exact fiber domain (7) therefore gives the scenery graph of S_T and its norm bound in Theorem 1.2. This is not an attempted clock chain rule applied to an input outside the clock domain.

7 Exact clock-noise energy and a critical sector obstruction

The clock section also gives quantitative information below ordinary first-order differentiability. We keep the statements in terms of an explicit coupling and the original orthogonal clock-chaos projections. This avoids assuming an identification with a fractional operator domain.

Fix $T > 0$, put $c = \sqrt{T}$ and $\kappa = \sqrt{2/\pi}$, and let

$$G = W_T/\sqrt{T}, \qquad G' = W_{T+1} - W_T.$$

The variables G, G' are independent standard normals, independent of the scenery. For $0 \le h \le 1$, define

$$G_h = (1-h)G + \sqrt{h(2-h)}G', \qquad Z_T^{(h)} = B_{c|G_h|}.$$

Then G_h is standard normal and $\mathbb{E}GG_h = 1-h$. The construction uses the future clock increment G'; it is a Gaussian noise probe and is not an adapted physical-time driver. The same full random scenery is retained in both Z_T and $Z_T^{(h)}$.

Theorem 7.1 (Exact noise formula)**.** *For $0 \le h \le 1$,*

$$\|Z_T - Z_T^{(h)}\|_E^2 = c\kappa r(h), \qquad r(h) = \sqrt{4-2h} + \sqrt{2h} - 2. \tag{14}$$

One has

$$\frac{\sqrt{2h}}{4} \le r(h) \le \sqrt{2h}, \qquad \lim_{h\downarrow 0} \frac{r(h)}{\sqrt{h}} = \sqrt{2}. \tag{15}$$

In particular, for any real a,

$$\int_0^1 \frac{\|Z_T - Z_T^{(h)}\|_E^2}{h^{1+a}}\, dh < \infty \quad \Longleftrightarrow \quad a < \tfrac{1}{2}. \tag{16}$$

Proof. Conditioning on the clock and using the exact scenery increment metric gives the squared norm $c\mathbb{E}\big||G| - |G_h|\big|$. For arbitrary real x, y, checking their signs and which has larger absolute value gives

$$\big||x| - |y|\big| = |x+y| + |x-y| - |x| - |y|.$$

The centered Gaussian variables $G + G_h$ and $G - G_h$ have variances $4-2h$ and $2h$, respectively. Since a centered normal with variance v has first absolute moment $\kappa\sqrt{v}$, this proves (14).

For $h > 0$, rationalization gives

$$r(h) = \sqrt{2h} - \frac{2h}{2+\sqrt{4-2h}}, \qquad \frac{r(h)}{\sqrt{h}} = \sqrt{2} - \frac{2\sqrt{h}}{2+\sqrt{4-2h}}.$$

The limit and upper bound follow. Moreover

$$\frac{r(h)}{\sqrt{2h}} = 1 - \frac{\sqrt{2h}}{2+\sqrt{4-2h}} \ge 1 - \frac{\sqrt{2}}{2} \ge \frac{1}{4} \quad (0 < h \le 1).$$

The case $h = 0$ is immediate. As $c\kappa > 0$, the integrand in (16) is bounded above and below by positive constant multiples of $h^{-a-1/2}$. Its integral at zero is finite exactly when $-a - 1/2 > -1$. There is no other endpoint singularity. □

Let Π_n^W be the original orthogonal projection onto the nth homogeneous chaos of the clock Gaussian source, acting as the identity on all scenery randomness. Put

$$E_n(T) = \|\Pi_n^W Z_T\|_E^2, \qquad n \geq 0.$$

These are energies of the actual endpoint, not of its scenery average. They are nonnegative and sum to $\|Z_T\|_E^2 = c\kappa$.

Lemma 7.2 (Clock-sector noise identity)**.** *For $0 \leq h \leq 1$,*

$$2\sum_{n\geq 0}(1-(1-h)^n)E_n(T) = c\kappa r(h). \tag{17}$$

Every odd clock projection of Z_T vanishes.

Proof. The section $\Phi(x) = B_{c|x|}$ belongs to $L^2(\gamma; L^2(\mathbb{P}_B))$, where γ is standard Gaussian measure. Its Hilbert-valued Hermite expansion is

$$\Phi(x) = \sum_{n\geq 0} b_n \frac{\mathrm{He}_n(x)}{\sqrt{n!}}, \qquad \sum_n \|b_n\|_{L^2(\mathbb{P}_B)}^2 = c\kappa.$$

Completeness follows by applying scalar Hermite completeness to the coordinates of a countable Hilbert basis of $L^2(\mathbb{P}_B)$; Parseval and nonnegative summation give the asserted expansion and energy equality. A scalar function of the single normalized clock direction G has its nth Hermite component in the original full clock chaos of order n. Finite Hermite expansions and continuity of the orthogonal projections therefore identify $E_n(T) = \|b_n\|^2$.

For standard normals X, Y of correlation ρ, their exponential generating functions give

$$\mathbb{E}[e^{sX-s^2/2}e^{tY-t^2/2}] = e^{\rho st}, \qquad \mathbb{E}[\mathrm{He}_n(X)\,\mathrm{He}_m(Y)] = \mathbf{1}_{n=m}n!\rho^n.$$

Differentiation at $s = t = 0$ is justified by Gaussian exponential integrability. For finite Hilbert-valued Hermite sums it follows that the squared difference at G, G_h is $2\sum_n(1-(1-h)^n)\|b_n\|^2$. Both pullbacks from $L^2(\gamma; L^2(\mathbb{P}_B))$ into the joint space are isometries, so the finite sums converge in both pullback norms. The series has nonnegative terms bounded by $2E_n(T)$. Passing to the limit and using (14) proves (17). Finally $\Phi(-x) = \Phi(x)$ whereas odd Hermite polynomials change sign. Thus $b_n = 0$ for odd n, and the corresponding original clock projections vanish. □

Theorem 7.3 (Critical weighted energy fails)**.** *For each $T > 0$,*

$$\sum_{n\geq 1}\sqrt{n}\,E_n(T) = \infty. \tag{18}$$

Consequently, for every $a \geq \frac{1}{2}$,

$$\sum_{n\geq 0}(1+n)^a E_n(T) = \infty.$$

Beyond every finite cutoff there is a nonzero even clock sector of Z_T.

Proof. For $n \geq 0$ and $0 < h \leq 1$, Bernoulli's inequality gives

$$0 \leq 1-(1-h)^n \leq \min(1, nh) \leq \sqrt{nh}.$$

For each fixed n the same bound by nh gives $(1-(1-h)^n)/\sqrt{h} \le n\sqrt{h} \to 0$. If the sum in (18) were finite, dominated convergence for counting measure, with dominating sequence $2\sqrt{n}E_n(T)$, would imply

$$\lim_{h\downarrow 0} 2\sum_{n\ge 0} \frac{1-(1-h)^n}{\sqrt{h}} E_n(T) = 0.$$

But (17) and (15) give the strictly positive limit $c\kappa\sqrt{2}$, a contradiction. For $a \ge 1/2$, $(1+n)^a \ge \sqrt{n}$, proving the second assertion. If all even projections beyond some cutoff vanished, odd-sector vanishing would make the critical sum finite. This proves the final assertion. □

8 The exact subcritical clock Bessel domain

The positive half of the energy threshold can also be proved for the original clock sectors. For $0 < a < 1/2$ and $n \ge 1$, define

$$I_a(n) = 2\int_0^1 \frac{1-(1-h)^n}{h^{1+a}}\, dh.$$

This is finite, since $1-(1-h)^n \le nh$ and $a < 1$. For $0 < h \le 1/(2n)$, the finite geometric sum and Bernoulli's inequality give

$$1-(1-h)^n = h\sum_{j=0}^{n-1}(1-h)^j \ge nh(1-h)^{n-1} \ge nh(1-(n-1)h) \ge nh/2.$$

Integration on this smaller interval proves the explicit lower bound

$$I_a(n) \ge n\int_0^{1/(2n)} h^{-a}\, dh = \frac{2^{a-1}}{1-a} n^a.$$

All summands in the clock-noise identity are nonnegative. Tonelli's theorem, followed by (16), therefore gives

$$\sum_{n\ge 1} I_a(n)E_n(T) = \int_0^1 \frac{\|Z_T - Z_T^{(h)}\|_E^2}{h^{1+a}}\, dh < \infty.$$

The lower bound proves $\sum_{n\ge 1} n^a E_n(T) < \infty$. Since $(1+n)^a \le 2^a n^a$ for $n \ge 1$, the same holds for the Bessel weights after adding the finite zeroth sector. For $a \le 0$, $(1+n)^a \le 1$ and Parseval gives summability immediately. Together with Theorem 7.3, this proves for every real a

$$\sum_{n\ge 0}(1+n)^a E_n(T) < \infty \quad\Longleftrightarrow\quad a < 1/2. \tag{19}$$

The argument uses positive series and a genuine integral, without a Tauberian assumption or an asserted coefficient asymptotic.

`IBMClockSubcritical`:
`modeIntegral_lower`, `weighted_mode_integrals_summable`,
`bessel_weight_summable`, `actual_bessel_summable_iff`.

To identify an actual closed operator, use the entire original joint Hilbert space E, not a scalar averaged endpoint. Its complete clock projections Π_n^W retain all scenery coordinates. For $s \in \mathbb{R}$

set

$$\mathcal{D}_s = \left\{ f \in E : \sum_{n\geq 0} (1+n)^s \|\Pi_n^W f\|_E^2 < \infty \right\}, \tag{20}$$

$$\mathcal{B}_s f = \sum_{n\geq 0} (1+n)^{s/2} \Pi_n^W f \quad (f \in \mathcal{D}_s). \tag{21}$$

Orthogonality and completeness prove convergence of this series exactly under the displayed domain condition and give $\|\mathcal{B}_s f\|^2 = \sum_n (1+n)^s \|\Pi_n^W f\|^2$. Its graph is the closed intersection of the coordinate equations $\Pi_n^W g = (1+n)^{s/2}\Pi_n^W f$. Finite sector truncations $f_N = \sum_{n\leq N} \Pi_n^W f$ belong to every $\mathcal{D}_s$ and converge to f in E. If $f \in \mathcal{D}_s$, the weighted square tail also tends to zero, so $(f_N, \mathcal{B}_s f_N) \to (f, \mathcal{B}_s f)$. Thus the operator is densely defined and these truncations form a graph core. This is not a claim that each sector is finite dimensional.

`GaussianClockSpectralPowers`:
`operator_domain`, `operator_isClosed`,
`operator_energy`, `graph_truncation_tendsto`.

The operator is positive and self-adjoint on this exact domain. For symmetry, its inner product with another domain input is the absolutely convergent sector pairing series, by Cauchy–Schwarz. If (g, b) is in its Hilbert-adjoint graph, test that pairing against arbitrary vectors in each individual clock sector. One obtains $\Pi_n^W b = (1+n)^{s/2}\Pi_n^W g$. Parseval for b then forces $g \in \mathcal{D}_s$ and $b = \mathcal{B}_s g$. Conversely symmetry supplies the other graph inclusion. Positivity follows from $\langle \mathcal{B}_s f, f\rangle = \sum_n (1+n)^{s/2}\|\Pi_n^W f\|^2 \geq 0$.

`GaussianClockSpectralAdjoint`:
`operator_adjoint`, `operator_ibp`, `operator_nonneg`.

Theorem 8.1 (Sharp actual endpoint regularity)**.** *At every $T > 0$, $Z_T \in \mathcal{D}_s$ if and only if $s < 1/2$. For $f \in \mathcal{D}_r$ one has*

$$\mathcal{B}_r f \in \mathcal{D}_s \iff f \in \mathcal{D}_{r+s}, \qquad \mathcal{B}_s \mathcal{B}_r f = \mathcal{B}_{r+s} f$$

whenever these equivalent conditions hold. In particular, for $\beta \geq 0$, $\mathcal{B}_{-2\beta}$ is a bounded operator on all of E and

$$\mathcal{B}_{-2\beta} Z_T \in \mathcal{D}_1 \iff \beta > 1/4.$$

Proof. The first assertion is (19) and the actual domain (20). For the shift assertion, $\Pi_n^W \mathcal{B}_r f = (1+n)^{r/2}\Pi_n^W f$, so its s-weighted squared energy is exactly the $(r+s)$-weighted energy of f. Multiplication of the scalar weights gives the composition after projection onto every sector; completeness identifies the outputs. When $\beta \geq 0$, the weights $(1+n)^{-\beta} \leq 1$ give an everywhere defined contraction. Applying the shift assertion to the actual endpoint gives $1 - 2\beta < 1/2$, equivalent to the strict stated condition. The endpoint $\beta = 1/4$ fails by the critical divergence. □

`IBMClockFractionalDomain`: `actual_bessel_domain`.
`IBMClockFractionalCalculus`:
`bessel_shift_domain`, `bessel_composition`,
`actual_regularized_domain`, `actual_smoothing_threshold`.

The hypothesis $f \in \mathcal{D}_r$ in the composition is essential. For $r > 0$ the product $\mathcal{B}_{-r}\mathcal{B}_r$ is the identity on $\mathcal{D}_r$, not an identity operator with domain all of E. No unqualified equality of those unbounded-operator domains is inferred.

The bridge back to the original first clock derivative

There is also an exact domain bridge, rather than just a comparison with a newly defined norm:

$$\mathcal{D}_1 = D(D_W), \qquad \|\mathcal{B}_1 f\|_E^2 = \|f\|_E^2 + \|D_W f\|^2. \tag{22}$$

Here both sides use the same original scenery-first joint space. The formal Bochner clock derivative uses the reversed factor order; the probability-factor flip is an isometry and must be applied before identifying its domain or vector output with this notation.

For completeness, the scalar Gaussian Hermite derivative lowers degree by one. On a normalized multi-index polynomial of total degree n, the sum of squared coordinate-gradient norms is n; orthogonality of the lowered Hermite basis also gives orthogonality of gradient outputs from different total degrees. Finite-coordinate polynomial approximation extends this to each whole clock sector, with derivative norm bounded by $\sqrt{n}$ times its value norm. Lifting to the scenery-valued space and using Tonelli preserves these identities and the factor flip. Consequently the original finite truncation graphs satisfy

$$\|D_W f_N\|^2 = \sum_{n \le N} n\|\Pi_n^W f\|^2, \qquad \|D_W f_M - D_W f_N\|^2 = \sum_{N < n \le M} n\|\Pi_n^W f\|^2 \quad (M \ge N).$$

For $f \in \mathcal{D}_1$ this proves that their gradients are Cauchy. Closedness of the original derivative puts f in $D(D_W)$ and gives the norm identity (22). Conversely, for a graph pair (f, v), scalar Gaussian derivative Parseval applied in almost every scenery fiber and nonnegative integration yield $\|v\|^2 = \sum_n n\|\Pi_n^W f\|^2$. Adding ordinary Parseval gives $f \in \mathcal{D}_1$ and the reverse norm identity. In particular the smoothed endpoint belongs to the *original* first clock domain precisely when $\beta > 1/4$.

`GaussianClockNativeDomain`:
`flipped_native_energy_hasSum`, `finite_gradient_energy`,
`bessel_native_domain`, `bessel_native_energy`,
`native_graph_truncation_tendsto`.

These statements retain the full random scenery and concern ambient Gaussian clock variables. Smoothing changes the observable; it does not put the unsmoothed Z_T in its excluded native domain or provide an observed-past driver. The noise parameter changes Gaussian correlation, not physical observation time. Gaussian spectral Sobolev theory and noise characterizations are classical. Geiss–Toivola [3], Section 2 (Gaussian Sobolev spaces and Definition 2.2), specify their scalar Gaussian gradient norm and real-interpolation Besov scale. We do not identify every such Besov scale or fine index with the particular Hilbert spectral domain above. The source-valued extension, its original operator bridge and the endpoint calculation were proved here explicitly; priority for this reflected-clock application remains unresolved.

9 Native smoothing can stay away from the observed right past

The preceding native-domain theorem has a separate causal limitation. Let $\mathcal{F}_u^Z = \sigma(Z_r : 0 \le r \le u)$ and, inside E, set

$$\mathcal{A}_T = \bigcap_{u > T} L^2(\mathcal{F}_u^Z).$$

Each past subspace is closed; completing its sigma algebra does not change its L^2 classes. Thus $\mathcal{A}_T$ is the completed observed right-past subspace used in the frozen construction. In particular $Z_T \in \mathcal{A}_T$. Denote its orthogonal projection by P_T. The following obstruction concerns this time-specific subspace, rather than the sigma algebra of all future observations.

Theorem 9.1. *There exist $T > 0$, $b \geq 1$ and $d > 0$ such that, for every real $\beta \geq b$, the smoothed actual endpoint $S_\beta = \mathcal{B}_{-2\beta} Z_T$ belongs to the original native clock derivative domain and*

$$\inf_{G \in \mathcal{A}_T} \|S_\beta - G\|_E \geq d/2 > 0.$$

Thus native Gaussian clock differentiability, even at arbitrarily large smoothing orders, does not imply observed right-past measurability.

Proof. The clock-vacuum component is the scenery-only variable

$$V_T = \Pi_0^W Z_T = \mathbb{E}_W[B_{|W_T|}],$$

where the expectation integrates the independent clock and retains the full scenery. This is a genuine Bochner $L^2(\mathbb{P}_B)$ integral by Cauchy–Schwarz. Projection onto clock chaos zero is precisely averaging that probability factor; its extension to the joint space is constant in the clock. Hence $V_T \in D(D_W)$ and $D_W V_T = 0$.

Let $\mathcal{H}_1 = \sigma(B_r : 0 \leq r \leq 1)$ and $Y = B_2 - B_1$. The centered Y is independent of $\mathcal{H}_1$. The actual Brownian scenery covariance, integrated over the clock, gives

$$\mathbb{E}[Y V_T] = \mathbb{E}_W[\min(|W_T|, 2) - \min(|W_T|, 1)] > 0 \quad (T > 0).$$

The integrand lies in $[0, 1]$ and equals one whenever $W_T > 2$, an event of positive probability for every nondegenerate Gaussian clock value. The pairing is legitimate by the L^2 bounds. Consequently V_T is not $\mathcal{H}_1$-measurable: otherwise independence and centering would force this pairing to vanish.

We now localize the actual observed history. Continuity of W and $W_0 = 0$ imply that

$$A_N = \{\sup_{0 \leq r \leq 1/N} |W_r| \leq 1\}, \qquad N \geq 1,$$

increase to a full-probability event. Their measurability follows by taking the supremum over rational times in the compact interval. Choose N with $\mathbb{P}_W(A_N) > 0$, set $S = 1/N$ and $T = S/2$. On A_N, every observed Z_r, $r \leq S$, uses only scenery on $[0, 1]$. More precisely, if a scenery-only $K \in L^2(\mathbb{P}_B)$ had a representative measurable with respect to the completed $\mathcal{F}_S^Z$, choose a raw-past representative and factor it measurably through $Z|_{[0,S]}$. Such a factorization is available because the continuous observed path takes values in the standard Borel space $C([0, S])$ and its evaluations generate that past sigma algebra. Fubini shows that for almost every fixed clock path this representative equals K for almost every scenery path. Pick one such clock in the positive-probability event A_N. Its entire observed history is a measurable function of $B|_{[0,1]}$, so the same factorization makes K $\mathcal{H}_1$-measurable, up to completion. This proves the required source-only restriction and also handles exceptional-set changes of representatives.

Applying the restriction to $K = V_T$ contradicts the positive future pairing above. Thus $V_T \notin L^2(\mathcal{F}_S^Z)$. As $S > T$, $\mathcal{A}_T \subset L^2(\mathcal{F}_S^Z)$, and therefore $V_T \notin \mathcal{A}_T$. Since this is a closed subspace,

$$d = \|V_T - P_T V_T\|_E > 0.$$

Finally, the spectral multiplier formula gives

$$\|S_\beta - V_T\|_E^2 = \sum_{n \geq 1} (1 + n)^{-2\beta} E_n(T) \leq 3^{-2\beta} \|Z_T\|_E^2.$$

The last bound uses the already proved vanishing of every odd sector; all surviving indices in this sum are at least two. Thus $S_\beta \to V_T$ strongly. For every $G \in \mathcal{A}_T$, the projection best-approximation property and the triangle inequality yield

$$\|S_\beta - G\|_E \geq d - \|S_\beta - V_T\|_E \geq d - 3^{-\beta}\|Z_T\|_E.$$

Choose $b \geq 1$ so that $3^{-b}\|Z_T\|_E \leq d/2$. This proves the uniform distance for every real $\beta \geq b$; the strict native threshold $\beta > 1/4$ already proved gives the native derivative domain simultaneously. □

`IBMClockVacuumPast`, namespace `IBMLocalPastRestriction`:
`past_sections_confined`, `source_only_past_requires_history`,
`exists_positive_confined_time`.
`IBMClockVacuumPast`:
`clock_vacuum_const`, `vacuumSource_pairing_positive`,
`vacuumSource_not_history`,
`exists_actual_adapted_clock_projection_failure`.
`IBMClockSmoothingPast`:
`smooth_tendsto_vacuum`, `smooth_native_iff`,
`positive_distance_from_all_adapted`,
`exists_actual_uniform_smoothing_obstruction`.

The proof reorganizes the source's measurable past-coefficient and section argument using continuous observed paths. Its displayed $3^{-\beta}$ convergence rate is an elementary human consequence of odd-sector vanishing and Parseval; the mapped convergence theorem uses dominated convergence without claiming this rate. The time T is existential, not asserted arbitrary, and b, d are not numerically estimated. No conclusion is drawn for every small positive smoothing order, for preservation of the derivative domain by P_T, or for membership in the full all-time intrinsic observable space. The result is an actual fixed-time causal obstruction, not a completion of physical-time stochastic calculus or a new abstract smoothing principle.

10 Graph-dual derivatives and escape of the original field norm

We give the closed-operator argument separately to make its topology and representability exact. Let $T : D(T) \subset E_0 \to F_0$ be a closed densely defined operator between real Hilbert spaces. Put

$$\mathcal{G} = D(T^*), \qquad \|v\|_{\mathcal{G}}^2 = \|v\|_{F_0}^2 + \|T^*v\|_{E_0}^2.$$

The adjoint is closed, so this is a Hilbert space. Define bounded maps

$$df(v) = \langle f, T^*v\rangle_{E_0}, \qquad jV(v) = \langle V, v\rangle_{F_0}. \tag{23}$$

Cauchy–Schwarz gives $\|df\| \leq \|f\|$ and $\|jV\| \leq \|V\|$. For $u \in D(T)$,

$$jTu = du. \tag{24}$$

Proposition 10.1. *The map j is injective, and $df = jV$ for some $V \in F_0$ if and only if $f \in D(T)$, in which case $V = Tf$. If $f \notin D(T)$, then $df \neq 0$.*

Proof. Write $\Gamma = \operatorname{graph} T$ in $E_0 \oplus F_0$. By the definition of the adjoint and density of $D(T)$,

$$\Gamma^{\perp} = \{(-T^*v, v) : v \in D(T^*)\}.$$

The identity $df = jV$ says exactly that (f, V) is orthogonal to this orthogonal complement. Since Γ is closed, $\Gamma^{\perp\perp} = \Gamma$; thus (f, V) belongs to the original graph. The converse is (24). If $jV = 0$, the same orthogonality argument puts $(0, V)$ in Γ, so $V = T0 = 0$. This also proves that $D(T^*)$ is dense in F_0. Finally $df = 0 = j0$ would put f in $D(T)$, proving the last assertion. □

For any $u \in D(T)$ and $f \in E_0$, the graph-dual error bound is immediate:

$$\|jTu - df\|_{\mathcal{G}^*} = \|d(u - f)\|_{\mathcal{G}^*} \leq \|u - f\|_{E_0}. \tag{25}$$

Density of $D(T)$ supplies such approximating values for every f. This convergence is in the operator norm of the graph-test dual, not in the original field space.

Proposition 10.2. *If $u_n \in D(T)$, $u_n \to f$ in E_0, and $f \notin D(T)$, then $\|Tu_n\|_{F_0} \to \infty$.*

Proof. If the norms did not tend to infinity, some subsequence would have $\|Tu_{n_j}\| \leq C$ for a finite C. For $v \in D(T^*)$ take the limit in $\langle u_{n_j}, T^*v\rangle = \langle Tu_{n_j}, v\rangle$:

$$|\langle f, T^*v\rangle| \leq C\|v\|_{F_0}.$$

Since $D(T^*)$ is dense in F_0, this functional extends continuously to all of F_0. Riesz representation provides $V \in F_0$ with $\langle f, T^*v\rangle = \langle V, v\rangle$ for every adjoint-domain test. Proposition 10.1 would put f in $D(T)$, a contradiction. □

Apply these results to $T = D_W$, the closed densely defined ambient clock lift of Section 2, and $f = Z_T$ or S_T (with the subscript $T > 0$ now denoting time). Theorems 1.1 and 1.2 give nonmembership, and (23), (25) and Proposition 10.2 prove Theorem 1.3. The bound on the weak derivative is explicitly $\|d_W Z_T\| \leq (2T/\pi)^{1/4}$, while $\|d_W S_T\| \leq 1$. Every original-domain value approximation has divergent clock energy, although its embedded derivatives converge in a weaker, precisely defined norm.

Nonlinear composition in the full adjoint-graph dual

The same topology permits a precise nonlinear limit without a pointwise product of a random coefficient and a distribution. Return to the actual ambient clock derivative D_W on the original joint probability space. Use its L^2 field space and the full graph-test space $\mathcal{G} = D(D_W^*)$ with the norm above. Let $\phi \in C^1(\mathbb{R})$ satisfy

$$|\phi(x)| \leq A, \qquad |\phi'(x)| \leq L \quad (x \in \mathbb{R}), \qquad \phi' \text{ continuous},$$

with finite $A, L \geq 0$. For every $f \in L^2$ define $d_W(\phi(f))$ directly by (23). The composition is an actual L^2 class because the joint law is a probability measure. If $u_n \in D(D_W)$ and $u_n \to f$ in L^2, the original closed Gaussian chain rule gives

$$\phi(u_n) \in D(D_W), \qquad D_W\phi(u_n) = \phi'(u_n)D_W u_n.$$

The product on the right is formed in the genuine original field L^2 space at each finite n, since ϕ' is bounded. The chain rule follows by approximating in the Gaussian graph core, selecting an

almost-everywhere convergent value subsequence, and passing the gradients to the limit. More explicitly, if $u_j \to u$ and $D_W u_j \to V$ in their L^2 spaces, then

$$\|\phi'(u_j)D_W u_j - \phi'(u)V\|_2 \le L\|D_W u_j - V\|_2 + \|(\phi'(u_j) - \phi'(u))V\|_2 \longrightarrow 0.$$

For the second term, continuity of ϕ' gives convergence almost everywhere along the selected subsequence and the square is dominated by $4L^2|V|^2$. The value convergence follows from the mean-value bound $|\phi(x) - \phi(y)| \le L|x - y|$. Closedness then retains the original derivative graph. This is the classical Gaussian Sobolev chain rule, applied in actual product sections; no independent measurable choice of section derivatives is required.

Apply (25) to $\phi(u_n)$ and $\phi(f)$. We obtain the quantitative full-dual convergence

$$\|j(\phi'(u_n)D_W u_n) - d_W(\phi(f))\|_{\mathcal{G}^*} \le L\|u_n - f\|_2 \longrightarrow 0. \tag{26}$$

The limit is independent of the approximating sequence. On every $v \in D(D_W^*)$, its literal stochastic pairing is

$$d_W(\phi(f))(v) = \mathbb{E}[\phi(f)D_W^* v].$$

This integral is absolutely integrable by Cauchy–Schwarz. For two values $f, g \in L^2$, the same argument gives

$$\|d_W(\phi(f)) - d_W(\phi(g))\|_{\mathcal{G}^*} \le L\|f - g\|_2.$$

No assertion that $\phi'(f)d_W f$ is a defined graph-dual product is needed for either formula. In particular boundedness of a coefficient alone does not say that multiplication preserves $D(D_W^*)$ with its graph norm.

For a concrete actual-process consequence choose $f = Z_T$, $\phi(x) = \sin x$, and any original-domain approximation $u_n \to Z_T$ in L^2. Such approximations exist by density; the constructed spatial mollifications are one example. Then (26) holds with $L = 1$ and field $\cos(u_n)D_W u_n$. Nevertheless

$$\|\cos(u_n)D_W u_n\|_2 \longrightarrow \infty. \tag{27}$$

Indeed $\sin u_n \to \sin Z_T$ in L^2, and Theorem 1.2 places $\sin Z_T$ outside $D(D_W)$ for every $T > 0$. Proposition 10.2, applied to the native values $\sin u_n$, proves (27). Thus the same genuine fields converge after embedding in $\mathcal{G}^*$ while their original norms diverge. The limiting functional is nonzero and has no original L^2 field representative by Proposition 10.1. This explicit sine escape is a human combination of the mapped chain rule, domain obstruction and escape theorem, rather than a separate named formal assertion. None of these statements constructs an observed-filtration time integral or a pointwise multiplication rule for arbitrary graph-dual distributions.

`GaussianClosedChainRule.bounded_product_chain_graph`.

`IBMClockWeakChain.chain_graph`.

`IBMClockWeakChain.chain_dual_error`.

`IBMClockWeakChain.chain_dual_limit`.

`IBMClockWeakChain.actual_pairing`.

11 Physical spatial kernels and the white-noise domain boundary

The negative norm in the preceding section is an adjoint-graph norm. There is also a separate, actual physical-space $H^{-1}(\mathbb{R})$ construction. It has a different variable, operator and norm, which must be specified before comparing limits. In this section $x \in \mathbb{R}$ is the spatial coordinate of the two-sided scenery. It is not the clock Gaussian coordinate or the physical observation time.

An isonormal map fixed by the original scenery

In $L^2(\mathbb{R}, dx; \mathbb{R})$ define the oriented indicator

$$v_r(x) = \mathbf{1}_{\{x \le r\}} - \mathbf{1}_{\{x \le 0\}}.$$

For $r \ge 0$ this is $\mathbf{1}_{(0,r]}$, and for $r < 0$ it is $-\mathbf{1}_{(r,0]}$. Direct integration gives

$$\|v_r\|_2^2 = |r|, \qquad \|v_r - v_s\|_2^2 = |r - s|, \qquad \langle v_r, f \rangle = \int_0^r f(x)\, dx.$$

The span of these vectors is dense. Indeed, a vector orthogonal to them has primitive $\int_0^r f = 0$ for every r. Local integrability and the almost-everywhere fundamental theorem of calculus then give $f = 0$. Thus the orthogonal complement of the span is zero.

The actual two-sided scenery satisfies the same norm and increment identities: $\mathbb{E}B_r^2 = |r|$ and $\mathbb{E}(B_r - B_s)^2 = |r - s|$. Polarization gives equality of the Gram matrices of (v_r) and (B_r). Therefore

$$\mathcal{I}\left(\sum_j a_j v_{r_j}\right) = \sum_j a_j B_{r_j}$$

is well defined and isometric on their finite span. Density extends it uniquely to an isometry from all physical $L^2(\mathbb{R})$ into the original scenery Gaussian L^2 space. Its image lies in first chaos; it is not asserted to exhaust every unused ambient Gaussian coordinate. The map is fixed by $\mathcal{I}v_r = B_r$, rather than chosen by an unrelated basis identification. Finite Gaussian laws and passage to L^2 limits show

$$\mathbb{E}\mathcal{I}f = 0, \qquad \mathbb{E}[(\mathcal{I}f)(\mathcal{I}g)] = \langle f, g \rangle, \qquad \mathcal{I}f \sim N(0, \|f\|_2^2).$$

This is the original physical scenery white-noise integral. It is a map with deterministic L^2 input; a distribution-valued realization of white noise does not automatically enlarge that input space. Hairer [5], Example 4.63 and formula (4.16), provide the classical isonormal covariance context.

Strong physical kernels and anchored Dirac limits

Let ρ_n be the actual nonnegative normalized smooth compactly supported mollifiers, with support in $[-\varepsilon_n, \varepsilon_n]$, where $\varepsilon_n = 2/(n+1)$. Define the physical L^2 kernel

$$K_{n,r} = \int_{\mathbb{R}} \rho_n(r - s) v_s\, ds.$$

The Bochner integral exists: the parameter is confined to a compact interval and $s \mapsto v_s$ is L^2 continuous. Bounded linearity of $\mathcal{I}$ gives the actual source equality $\mathcal{I}K_{n,r} = \int \rho_n(r - s) B_s\, ds$. Normalization and the indicator metric give the uniform estimate

$$\|K_{n,r} - v_r\|_2 \le \int \rho_n(r - s)|s - r|^{1/2} ds \le \varepsilon_n^{1/2}. \tag{28}$$

This elementary rate is a human deduction; the formal source supplies actual strong physical L^2 convergence and source realization.

Use the actual Bessel space $H^{-1}(\mathbb{R}; \mathbb{C})$, with real scalars for maps from real kernels. Spatial distributional differentiation is a bounded map $\partial_x : L^2(\mathbb{R}) \to H^{-1}(\mathbb{R})$: the Fourier multiplier for a derivative, divided by the Bessel weight, is bounded. Write C_∂ for its fixed norm; no convention-

dependent sharp numerical value is claimed. A single Dirac mass belongs to this space because its Fourier transform has constant modulus and $\int_{\mathbb{R}}(1+\xi^2)^{-1}d\xi < \infty$.

Testing against a Schwartz function and using the oriented integral gives

$$\partial_x v_r = \delta_0 - \delta_r.$$

For the mollified kernel, interchange the bounded distributional test with its compactly supported Bochner integral. Since $\int \rho_n = 1$, the same identity yields

$$\partial_x K_{n,r} = \delta_0 - \rho_n(r - \cdot), \qquad \rho_n(r - \cdot) \longrightarrow \delta_r \quad \text{in } H^{-1}. \tag{29}$$

The last assertion follows from boundedness of ∂_x and (28), with error at most $C_\partial \varepsilon_n^{1/2}$. The zero-order embedding of the scalar mollifier into H^{-1} is understood in this formula. The physical derivative, zero-order embedding and a Dirac mass are three different objects.

These limits also hold strongly in $L^2(\Omega_W; L^2(\mathbb{R}_x))$ and $L^2(\Omega_W; H^{-1}(\mathbb{R}_x))$ at the actual random clock $r = |W_t|$. The uniform bound (28) gives the same error bound after the clock expectation. The target indicator field is square integrable since $\mathbb{E}\|v_{|W_t|}\|_2^2 = \mathbb{E}|W_t| < \infty$; the mollified fields are then square integrable by the triangle inequality. Strong measurability follows from continuity in r and the actual measurable clock. Applying the original scenery isometry in each independent clock section realizes $B_{|W_t|}$ and its mollified values in the same source space. The derivative limit is precisely $\delta_0 - \delta_{|W_t|}$ in the physical variable. These statements do not supply a derivative in the clock Gaussian coordinate.

The same Dirac limit cannot be fed into the Wiener map

At $r = 0$, put $p_n(x) = \rho_n(-x)$. The actual zero-order embedded profiles satisfy $p_n \to \delta_0$ in H^{-1} by (29). However

$$1 = \left| \int_{-\varepsilon_n}^{\varepsilon_n} p_n(x)dx \right| \le (2\varepsilon_n)^{1/2}\|p_n\|_2, \qquad \|\mathcal{I}p_n\|_{L^2(\Omega_B)} = \|p_n\|_2 \ge \frac{\sqrt{n+1}}{2} \longrightarrow \infty.$$

This is just Cauchy–Schwarz on the shrinking support and the original isometry. It rules out convergence to any original Gaussian L^2 random variable. More strongly, there is no map $F : H^{-1}(\mathbb{R}) \to L^2(\Omega_B)$, even nonlinear, continuous at δ_0 and agreeing with $\mathcal{I}$ on all ordinarily embedded real L^2 kernels. Otherwise continuity along this sequence would force $\mathcal{I}p_n \to F(\delta_0)$, contradicting norm escape. The source formalizes this actual nonextension and escape; the explicit $\sqrt{n+1}/2$ lower bound is a labelled human consequence of the particular support radius.

The formal physical-kernel, Dirac and white-noise modules supply the original isometry, exact source equalities, strong random-clock Bochner limits and the nonextension theorem. This construction does not manufacture a random variable $\mathcal{I}\delta_r$ in the old L^2 space. Nor does the physical negative norm become the preceding clock Bessel norm or the adjoint-graph-dual norm. The positive physical distributional limit and the negative original-noise statement are compatible precisely because their operators have different domains and topologies.

12 Scope, antecedents and formal correspondence

The Brownian-time model is classical [1]. Gaussian derivatives act on its independent sources; Z itself is not Gaussian. Standard derivative/adjoint conventions appear in [2]. The Brownian-time

change-of-measure work [4] addresses a different question; none of the exclusions here rules out every possible Brownian-time calculus. Novelty remains unresolved.

All formal correspondences use frozen verified commit

`bab688bb6140cd6316c80680095a914ea0d346db.`

The accompanying theorem map records exact sources for the folded-section metric, primitive and contraction arguments, point and sine exclusions, clock-noise and critical/subcritical sector energies, exact closed spectral domains and the original first-clock domain bridge, and generic graph-dual representation/error/escape theorems. The ambient clock application remains distinct from the separate native adjoint-graph-dual time-average results.

The formal realization may contain unused independent scenery coordinates, including a negative-time branch. The half-line model retains the same process and partial domains on its observables: graph pairs tensor with constants in unused coordinates; conversely averaging those coordinates contracts the graph test identities. They do not provide a substitute clock gradient.

Value regularization alone cannot produce a convergent original clock-gradient sequence at the point or sine; closedness and the norm-escape theorem exclude it. Scalar tests without one common finite-energy field do not meet the recovered-field hypothesis. Neither a source-variable divergence nor an adjoint-graph test functional is identified as an observed-past physical-time integrand. No physical-time Itô/Clark–Ocone completion or changing-past integrated point-regularizer limit is proved.

References

[1] H. Allouba and W. Zheng, Brownian-time processes: the PDE connection and the half-derivative generator, *Ann. Probab.* **29** (2001), 1780–1795. `https://arxiv.org/abs/1005.3801`.

[2] T. Sottinen and L. Viitasaari, Stochastic analysis of Gaussian processes via Fredholm representation, *Int. J. Stoch. Anal.* **2016**, article 8694365. doi:10.1155/2016/8694365.

[3] S. Geiss and A. Toivola, On fractional smoothness and L_p-approximation on the Gaussian space, *Ann. Probab.* **43** (2015), 605–638. doi:10.1214/13-AOP884. `https://arxiv.org/abs/1206.5415`.

[4] B. Abdurakhmanov and H. Allouba, Brownian-time change of measure, preprint, arXiv:2605.07044 (2026). `https://arxiv.org/abs/2605.07044`.

[5] M. Hairer, *An Introduction to Stochastic PDEs*, primary lecture notes, arXiv:0907.4178v2 (2023), Example 4.63 and formula (4.16). `https://arxiv.org/pdf/0907.4178`.