

Original Research Article

Trends of human influenza infection in Chhattisgarh: a retrospective
observational study

Raj Sharma, Rupendra K. Bharti*, Sanat Sharma, Mahendra K. Jaiswal

Department of Pharmacology,
Late BRKM Government
Medical College, Dimrapal,
Jagdalpur, Chhattisgarh, India

Received: 21 March 2019

Revised: 04 April 2019

Accepted: 03 May 2019

*Correspondence to:
Dr. Rupendra K. Bharti,
Email: rbharti000@gmail.com

Copyright: the author,
publisher and licensee Medip
Academy. This is an openaccess
article distributed under
the terms of the Creative
Commons Attribution NonCommercial
License, which
permits unrestricted noncommercial
use, distribution,
and reproduction in any
medium, provided the original
work is properly cited.

ABSTRACT

Background: Human influenza virus was recognized as a pandemic in 2009 by the World Health Organization. Since then many newer incidences was

recognized in India, but there was no sufficient data from all state of India. This study will provide data from the Chhattisgarh state of India.

Methods: It was a retrospective observational study from December 2015 to November 2017. All patient samples with suspected influenza infection were collected and analysed by Real-time reverse polymerase chain reaction (RT-PCR).

Results: 341 patients' sample was collected and analysed; among these samples, 07.9% of patients have all three serotype of influenza positive. Raipur district has the highest incidence of influenza A followed by Durg and Raigarh district of Chhattisgarh. There was no significant difference between male and female who was affected by the influenza virus.

Conclusions: The incidence of Human influenza virus is lesser in Chhattisgarh as compare to the average states of India and the state capital has a higher rate of sample collection as well as positive influenza infection.

Keywords: Chhattisgarh, Influenza, Incidence, Oseltamivir, Pandemic

INTRODUCTION

Human influenza virus is a pig transmitted infectious disease and it is appeared to be highest mortality causing disease since past decade. In 2009, it was recognized as pandemic infectious disease by World health Organization, which was transmitted via human to human by aerosol. Influenza was first time recognized as Spanish flu in 1918-1919, which was also pandemic in nature. Illness oftenly present in farmers who had swine in their farms and that was observed by veterinarian J. S. Koen. The isolation of pig derived H1N1 virus was first time obtained during 1930 by Shope and Lewis then in 1974 human

derived H1N1 virus was isolated, which confirm the hypothesis that pig derived H1N1 virus could infect human too.

Human influenza or H1N1 influenza or Swine flu is a respiratory tract disorder comprised of flu like symptoms which include sore throat, rhinorrhoea, fever, chills, vomiting, restlessness, anorexia, headache, arthralgia and diarrhea. In patients with immune deficit, the clinical presentation of swine flu could be worsen and may lead to altered sensorium and respiratory collapse. Till date five serotype of H1N1 influenza was isolated which are H1N1, H1N2, H2N3, H3N1, and H3N2. Among these only H1N1 causes human infection and rest are seen only in pigs.

Swine flu is an orthomyxo RNA virus, having two surface antigens and N1 (Neuraminidase type 1).¹

In India, 27,236 laboratory confirmed cases with 981 deaths were reported during 2009.^{4,5} The post pandemic period 62,000 laboratory confirmed swine flu with 5000 deaths was reported by the Ministry of Health and Family Welfare and also stated that the incidence was high during monsoon and low during winter.⁴⁻⁸ The incidences of H1N1 influenza were keep rising worldwide. In India there was lack in sufficient data of human influenza virus infection from the many states. This study will provide the newly laboratory diagnosed cases of human influenza virus infection in the state of Chhattisgarh.

METHODS

It was hospital-based retrospective descriptive, recordbased analysis from December 2015 to November 2017.

This study was carried out in the Department of Pharmacology,
Late
BRKM
Government
Medical

College, Dimrapal, Jagdalpur, Chhattisgarh. Patients with the sign and symptoms of influenza viral infection who are falling under category B and C and Confirm case (tested positive for influenza infection) by real-time reverse transcription polymerase chain reaction were included in this study.

Patients from category B and C were administered proper therapy for influenza viral infection and samples were sent for confirmation, if the sample was negative, treatment was discontinued.

The
data
related
to
patients'

demographics from the H1N1 visiting screening centers, swine flu outpatient departments and swine flu isolation wards and hospitals from Chhattisgarh were collected. Testing of influenza virus was carried out at microbiology department of Late BRKM Government Medical College, Dimrapal, Jagdalpur.

Collection of samples and confirmation

Throat or nasal swab samples of suspected cases with influenza-like sign and symptoms were collected in Viral Transport Medium and sent to laboratories maintaining cold-chain. RNA was extracted by Purelink Viral RNA/DNA blood Mini Kit. RNA of each isolate was tested by separate primer/probe sets for InfA, H1N1 and H1 subtype as per the CDC real-time RTPCR protocol (ABI Step One Plus RT-PCR instrument Applied Biosystems, USA).¹⁰ All the data related to demographic characteristics, was analysed using Epi Info v7.2 and Yates corrected chi-square test. P-value <0.05 was considered statistically significant.

RESULTS

Total 341 samples were collected from different hospitals and medical colleges in the state with suspected H1N1

infection. Out of total 341 sample, 54% patients were male and the mean age was 30.39 19.2, years-wise comparison depicted in Table 1. Trends of suspected cases of influenza infection between both the gender, higher percentage among female 31 to 40 were observed between 21 to 30 year of age and in males at 0-10 years were the highest. Between Dec 2015 to Nov 2017, only 7.9% patients were found to be positive of all three subtypes of human influenza stain like H1N1, Influenza-A and H1-subtype, while 46.3% patients was found positive for influenza-A, 17.3% for H1N1, 24.9% for H1 subtype infection by RT PCR and all three was positive in 7.9%. Populated region of Chhattisgarh has higher rate of suspected cases such as Raipur, and Durg, even the most Tribble region of Chhattisgarh has comparable higher rate than other district. Maximum number of sample was send from the Raipur district followed by Durg and Raigarh.

Table 1: Age wise distribution of Human Influenza

patients.

Age in decades

Male

Female

P Value

: Incidence of newer diagnosed human

influenza subtypes from December 2015November 2017.

The 41.8% sample was positive for Influenza A, 39% for H1 subtype infection in district Raipur, whereas in district Durg and Raigarh has 8.9%, 11.9%, 10.6% and 2.5%, 13.6%, 11.8% sample was positive for Influenza A, H1N1 and H1 subtype, respectively.

H1 subtupe

H1 N1

Influenza A

Incidence of Human Influenza

Sharma R and others Int J Basic Clin Pharmacol. 2019 Jun;8 :1288-1291

H1 subtupe

: Distribution of percentage of sample sends

by the district of Chhattisgarh.

: Distribution of Sero-positive influenza

subtype in most occurring districts of Chhattisgarh.

: Comparison of positive influenza serotype in

between gender.

Raipur has significant number of positive cases of influenza A, H1N1 and H1 subtype than other district of Chhattisgarh state. There was slightly higher incidence of Influenza A, H1N1, and H1 subtype observed in male patients than female. The mean percentage of Influenza- The positive test for influenza A, H1N1, and H1

subtype was high in males than female patients. A was positive in male patient was 51.3% as compare to female patient was 48.7%, while 52.5% male patients were positive for H1N1 as compare to 47.5% female patients and H1 subtype was positive in 50.6% male patient vs. 49.4% in female patients.

DISCUSSION

The incidence of human influenza viral infection in Chhattisgarh is comparable from other state of India. The collections of sample were higher during rainy season as reported in national health profile of India. From December 2015 to November 2017, a total number of 341 samples were sent to our center, out of which, highest number of sample was received from Raipur district followed by Durg district, which was comparable with the incidence of human influenza infection in the district of Rajasthan.

Among these samples 07.9% was seropositive for all three subtypes of influenza virus such as Influenza-A, H1N1, and H1 Subtype.

The incidence of Influenza-A was 46.3%, H1N1, and H1 subtype was 24.9% from Dec-2015-2017. Among higher reporting districts of Chhattisgarh, the overall newer case of from Raipur was 41.8% and Baster

for Influenza-A subtype. Similarly, H1N1 is also most common seen in Raipur followed by Raigarh and, H1 Subtype incidences were also high in Raipur followed by Durg. These results signify the higher rate of incidence commonly seen in the state capital as also observed by Malhotra et al. The mean age of male patients those samples was obtained was 30.39 19.2 years as compare to female patients was 30.77 18.8. In female, common age group presenting Influenza like symptoms was 21-30 years, while the common age group in male patients was 31-40 years, age-wise distribution is shown in the table-1, and similar distribution was also observed by choudhary et al.

The management of Human Influenza virus is requires isolation of infected person with antiviral therapy and curative management of other complications. Antiviral therapy mainly includes Oseltamivir, Zanamivir,

Peramivir, Laninamivir which are neuraminidase

inhibitors and Adamantanes such as Amantadine and Rimantadine.^{13,14} Nowadays only NAls is effective against swine flu due development of resistance against adamantanes.^{13,14} In India, Oseltamivir and Zanamivir is used for the treatment as well as for the prophylaxis of human influenza. According to WHO, Uncomplicated cases does not require antiviral therapy, only patients with suspected symptoms in high risk group were advocated for antiviral therapy. All patients including pregnant females were included by WHO for antiviral therapy.

Durg
9%
Mahasamund
5%

Raipur
37%
Rajnandgaon
3%

Dhantari

Bilashpur

Gariyaban

Baster

7%
Raigarh

BalodaBazar

Rest of the

Bemetara

H1 subtype

Influenza A
80

H1 Subtype
Influenza A
H1 N1

Number of Sample tested

Positive

Percentage of incidence

Sharma R and others Int J Basic Clin Pharmacol. 2019 Jun;8 :1288-1291

In adults, oseltamivir is prescribed 75 mg twice daily regimen for 5 days, while Zanamivir is given in a dose of 10 mg twice daily, both drugs can be used as prophylaxis of swine flu in an once daily regimen. Chemoprophylaxis should be given to the healthcare professionals, Oseltamivir is given once daily for 10 days and may be extended upto 6 weeks depending upon duration of close contact to the patients. WHO also recommended the vaccine for the healthcare professionals, the response of the vaccine will come after 2-3 weeks after vaccination. There were some limitation in our study; the power of study was low due to less sample size; the mortality was not calculated and we unable to find the commonest sign and symptoms of human influenza virus in Chhattisgarh region of India.

CONCLUSION

The case of Human influenza virus in Chhattisgarh from the period of December 2015 to November 2017 was only 341 which was less than many states of Chhattisgarh. Higher incidences were observed in the state capital then neighbouring district like Durg and Raigarh. There was no significant difference in between gender affected by human influenza, while Influenza A was more common than H1N1 and H1 subtype among the all received samples. The treatment remains the same for swine flu,

which includes neuraminidase inhibitors and Adamantanes.

ACKNOWLEDGEMENTS

Authors would like to thank Department of Microbiology,
Late BRKM GMC Jagdalpur, Chhattisgarh.

Funding: No funding sources

Conflict of interest: None declared

Ethical approval: The study was approved by the
Institutional Ethics Committee

REFERENCES

1. Rawal G, Yadav S, Kumar R, Sujana R. Swine Flu

: A concise review. Indian J
Immunol Resp Med. April-June;2 :29-32.

2. Myers KP, Olsen CW, Gray GC. Cases of Swine

Influenza in Humans: A Review of the Literature. Clin
Infect Dis. 2007 Apr 15;44 :1084-8.

3. Yadav S, Rawal G. Swine flu-have we learnt any

lesson from the past? Pan Afr Med J. 2015 Oct
12;22:118.

4. Central Bureau of Health Intelligence. National Health

Profile
of
India.
2009.
Available
at:

<http://cbhidghs.nic.in/writereaddata/linkimages/8%20Health%20Status%20Indicators4950277739.pdf>.

5. Press Information Bureau, Government of India,

Ministry of Health and Family. Preventive measures
on
swine
flu.
Available
at:

<http://pib.nic.in/newsite/PrintRelease.aspx?relid=115710>.

6. Central Bureau of Health Intelligence. National Health

Profile
of
India.
2010.
Available
at:

<http://cbhidghs.nic.in/writereaddata/mainlinkFile/File1012.pdf>

7. Choudhry AI, Singh S, Khare S, Rai A, Rawat DS,

Aggarwal RK, and others Emergence of pandemic 2009 influenza A H1N1, India. Indian J Med Res. 2012;135 :534-7.

8. Broor S, Krishnan A, Roy DS, Dhakad S, Kaushik S,

Mir MA, and others Dynamic patterns of circulating seasonal and pandemic A pdm09 influenza viruses from 2007-2010 in and around Delhi, India. PLoS One. 2012;7 :e29129.

9. Guidelines on Categorization of Influenza a H1N1

Cases during Screening for Home Isolation, Testing
Treatment,
and
Hospitalization.
Available
at:

<http://www.mohfw.nic.in/WriteReadData/l892s/804456402Categorisation.pdf>

10. CDC Protocol of Real Time RTPCR for Influenza A

.
Available
at:

http://www.who.int/csr/resources/publications/swineflu/CDCRealtimeRTPCR_SwineH1Assay2009_20090430.pdf

11. Malhotra B, Singh R, Sharma P and others Epidemiological

& clinical profile of influenza A 2009 virus infections during 2015 epidemic in Rajasthan. Indian J Med Res 2016;144 :918-23.

12. Choudhary SR, Momin MH, Chauhan NT. Profile of

influenza a positive patients of surat district. Int J Contemporary Med Res. 2017;4 :504-506.

13. Singh V, Sood M. Swine Flu - A comprehensive view.

14. Mukherjee S, Sen S, Nakate PC, Moitra S.

Management of swine flu outbreak and its treatment guidelines. Community Acquir Infect. 2015;2 :71-8.

[View publication stats](#)

Cite this article as: Sharma R, Bharti RK, Sharma S, Jaiswal MK. Trends of human influenza infection in Chhattisgarh: a retrospective observational study. *Int J Basic Clin Pharmacol* 2019;8:1288-91.