

---

## MATH 230A

**Name:** Quin Darcy  
**Instructor:** Dr. Domokos

**Due Date:** 9/10/20  
**Assignment:** Homework 1

---

1.1 Let  $A, B, C \subseteq X$ . Prove the following:

(a)  $A \cup (B \cap C) = (A \cup B) \cap (A \cup C)$ .

**Proof.**

$$\begin{aligned} x \in A \cup (B \cap C) &\Leftrightarrow (x \in A) \vee (x \in B \cap C) \\ &\Leftrightarrow (x \in A) \vee (x \in B \wedge x \in C) \\ &\Leftrightarrow (x \in A \vee x \in B) \wedge (x \in A \vee x \in C) \\ &\Leftrightarrow (x \in A \vee x \in B) \cap (x \in A \vee x \in C) \\ &\Leftrightarrow x \in (A \cup B) \cap (A \cup C). \end{aligned}$$

□

(c)  $A \setminus (B \cup C) = (A \setminus B) \cap (A \setminus C)$ .

**Proof.**

$$\begin{aligned} x \in A \setminus (B \cup C) &\Leftrightarrow (x \notin A) \wedge (x \in B \cup C) \\ &\Leftrightarrow (x \in A) \wedge (x \notin B \wedge x \notin C) \\ &\Leftrightarrow (x \in A \wedge x \notin B) \wedge (x \in A \wedge x \notin C) \\ &\Leftrightarrow (x \in A \setminus B) \wedge (x \in A \setminus C) \\ &\Leftrightarrow x \in (A \setminus B) \cap (A \setminus C). \end{aligned}$$

□

(e)  $(A \cup B)^c = A^c \cap B^c$ .

**Proof.**

$$\begin{aligned} x \in (A \cup B)^c &\Leftrightarrow x \notin A \cup B \\ &\Leftrightarrow (x \notin A) \wedge (x \notin B) \\ &\Leftrightarrow (x \in A^c) \wedge (x \in B^c) \\ &\Leftrightarrow x \in A^c \cap B^c. \end{aligned}$$

□

1.2 Prove that

(a)

$$\bigcup_{n=1}^{\infty} \left[ \frac{1}{n}, 1 \right] = (0, 1]$$

**Proof.** Let  $x$  be an element of the LHS. Then by Archimedes Principle, there exists  $n \in \mathbb{N}$  such that  $1/n < x$ . Thus,  $x \in [1/n, 1] \subseteq (0, 1]$ . Hence,

$$\bigcup_{n=1}^{\infty} \left[ \frac{1}{n}, 1 \right] \subseteq (0, 1].$$

Now assume  $x \in (0, 1]$ . Then  $x > 0$  and so by Archimedes Principle, there exists  $n \in \mathbb{N}$  such that  $1/n < x$  and thus

$$x \in \left[ \frac{1}{n}, 1 \right] \subseteq \bigcup_{n=1}^{\infty} \left[ \frac{1}{n}, 1 \right] \leftrightarrow (0, 1] \subseteq \bigcup_{n=1}^{\infty} \left[ \frac{1}{n}, 1 \right].$$

Therefore,

$$\bigcup_{n=1}^{\infty} \left[ \frac{1}{n}, 1 \right] = (0, 1].$$

□

(b)

$$\bigcap_{n=1}^{\infty} \left( 0, \frac{n+1}{n} \right) = (0, 1]$$

**Proof.** Note that for all  $n$ ,  $1 \leq (n+1)/n$  and so we immediately get that

$$(0, 1] \subseteq \bigcap_{n=1}^{\infty} \left( 0, \frac{n+1}{n} \right).$$

Thus, if  $x$  is an element of the intersection, then  $x \in (0, 1]$ . Hence,

$$\bigcap_{n=1}^{\infty} \left( 0, \frac{n+1}{n} \right) \subseteq (0, 1].$$

Therefore,

$$\bigcap_{n=1}^{\infty} \left( 0, \frac{n+1}{n} \right) = (0, 1].$$

□

1.3 Prove that  $\sqrt{2} \notin \mathbb{Q}$ .

**Proof.** Assume, for contradiction, that  $\sqrt{2} \in \mathbb{Q}$ . Then there exists  $a, b \in \mathbb{Z}$  such that  $b \neq 0$ ,  $\gcd(a, b) = 1$  and  $\sqrt{2} = a/b$ . Taking the square of both sides we obtain

$$2 = \frac{a^2}{b^2} \rightarrow 2b^2 = a^2. \quad (1)$$

This means that  $a^2$  is an even number. Note that if  $a$  is odd, i.e.,  $a = 2k + 1$  for some  $k \in \mathbb{Z}$ , then  $a^2 = 4k^2 + 4k + 1$  which is an odd number. Hence, if  $a^2$  is even, then  $a$  is even. Thus we can write  $a = 2m$  for some  $m \in \mathbb{Z}$ . Substituting into (1), we get

$$2b^2 = (2m)^2 = 4m^2 \rightarrow b^2 = 2m^2.$$

Hence,  $b$  is an even number. Thus,  $2 \mid a$  and  $2 \mid b$  which implies  $2 \mid \gcd(a, b)$ . Thus,  $\gcd(a, b) \neq 1$  which is a contradiction. Therefore  $\sqrt{2} \notin \mathbb{Q}$ .  $\square$

1.4 Let  $f: X \rightarrow Y$  be a function. Show that  $f$  is one-to-one if and only if for every  $y \in Y$ , the pre-image  $f^{-1}(\{y\})$  contains at most one element.

**Proof.** ?  $\square$

1.5 Let  $f: X \rightarrow Y$  be a function. Show that  $f$  is onto if and only if for every  $y \in Y$ , the pre-image  $f^{-1}(\{y\}) \neq \emptyset$ .

**Proof.** Assume  $f$  is onto. Then by Definition 1.5, for every  $y \in Y$  there exists  $x \in X$  such that  $f(x) = y$ . Hence, if  $y \in Y$ , then  $f^{-1}(\{y\}) = \{x \in X \mid f(x) \in \{y\}\}$  which is necessarily nonempty by the previous line. Now assume that for every  $y \in Y$ ,  $f^{-1}(\{y\}) \neq \emptyset$ . Let  $y \in Y$  and let  $x \in f^{-1}(\{y\})$ , then  $f(x) = y$ . Thus for every  $y \in Y$ , there exists  $x \in X$  such that  $f(x) = y$ . Therefore,  $f$  is onto.  $\square$

1.6 Let  $f: X \rightarrow Y$  be a function. Show that  $f$  is one-to-one if and only if  $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$  for every pair of sets  $A_1, A_2 \subseteq X$ .

**Proof.** Assume that  $f$  is one-to-one. Let  $A_1, A_2 \subseteq X$ . By Proposition 1.4.(b), we know  $f(A_1 \cap A_2) \subseteq f(A_1) \cap f(A_2)$ . Let  $y \in f(A_1) \cap f(A_2)$ . Then  $y \in f(A_1) \wedge y \in f(A_2)$ . This implies that there is some  $x_1 \in A_1$  and  $x_2 \in A_2$  such that  $f(x_1) = y = f(x_2)$ . Since  $f$  is one-to-one, then  $x_1 = x_2$ . Hence,  $x_1 \in A_1 \wedge x_1 \in A_2$ . Thus  $x_1 \in A_1 \cap A_2$ . Thus,  $f(x_1) = y \in f(A_1 \cap A_2)$ . Therefore,  $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$ .

Assume that  $f(A_1 \cap A_2) = f(A_1) \cap f(A_2)$  for all  $A_1, A_2 \subseteq X$ . Let  $x_1, x_2 \in X$  and assume  $y \in f(\{x_1\}) \cap f(\{x_2\})$ . Then  $f(x_1) = y = f(x_2)$ . Additionally, we have that  $y \in f(\{x_1\} \cap \{x_2\})$ . Thus either  $\{x_1\} \cap \{x_2\} = \emptyset$  or  $x_1 = x_2$ . The latter cannot be true since this would imply  $f(\emptyset) = y$ . Hence,  $x_1 = x_2$  and therefore  $f$  is one-to-one.  $\square$

- 1.7 Let  $f: X \rightarrow Y$  be a function. Show that  $f$  is one-to-one if and only if  $f^{-1}(f(A)) = A$  for every set,  $A \subseteq X$ .

**Proof.** Assume that  $f$  is one-to-one and that  $A \subseteq X$ . If  $a \in A$ , then  $f(a) \in f(A)$ . The latter is true iff  $a \in f^{-1}(f(A))$ . Hence,  $A \subseteq f^{-1}(f(A))$ . Now let  $x \in f^{-1}(f(A))$ . This implies that  $f(x) \in f(A)$ . Thus there exists  $a \in A$  such that  $f(x) = f(a)$  and since  $f$  is one-to-one, then  $x = a$ . Thus  $x \in A$  and so  $f^{-1}(f(A)) \subseteq A$ . Therefore  $f^{-1}(f(A)) = A$ .

Assume  $f^{-1}(f(A)) = A$  for all  $A \subseteq X$ . Let  $x_1, x_2 \in X$  such that  $f(x_1) = f(x_2)$ . By our assumption we have that  $f^{-1}(f(\{x_1\})) = \{x_1\}$  and  $f^{-1}(f(\{x_2\})) = \{x_2\}$ . Since  $f(x_1) = f(x_2)$ , it follows that  $f(x_1), f(x_2) \in f(\{x_1\})$ . Similarly,  $f(x_1), f(x_2) \in f(\{x_2\})$ . If there were some  $y \neq f(x_1) = f(x_2)$  such that  $y \in f(\{x_1\})$ , then  $f(x_1) = y \neq f(x_1)$  which would contradict that  $f$  is well-defined. Hence  $f(\{x_1\}) = f(x_1)$  and  $f(\{x_2\}) = f(x_2)$ . Thus

$$f^{-1}(f(\{x_1\})) = f^{-1}(f(x_1)) = \{x_1\} = \{x_2\} = f^{-1}(f(x_2)) = f^{-1}(f(\{x_2\}))$$

Which implies that  $x_1 = x_2$ . Therefore  $f$  is one-to-one.  $\square$

- 1.8 Let  $f: X \rightarrow Y$  be a function. Show that  $f$  is onto if and only if  $f(f^{-1}(B)) = B$  for all  $B \subseteq Y$ .

**Proof.** Assume  $f$  is onto. Let  $y \in B$ . Then since  $f$  is onto,  $f(f^{-1}(y)) = y$  and  $f^{-1}(y) \in f^{-1}(B)$ . Thus,  $y = f(f^{-1}(y)) \in f(f^{-1}(B))$ . Hence  $B \subseteq f(f^{-1}(B))$ . Now let  $y \in f(f^{-1}(B))$ . Then  $f^{-1}(y) \in f^{-1}(B)$ . Thus there is some  $y' \in B$  such that  $f^{-1}(y) = f^{-1}(y')$ . Since  $f$  is onto then  $f(f^{-1}(y)) = y = y' = f(f^{-1}(y'))$ . Hence  $y = y' \in B$  and so  $f(f^{-1}(B)) \subseteq B$ . Therefore  $f(f^{-1}(B)) = B$ .

Assume  $f(f^{-1}(B)) = B$  for all  $B \subseteq Y$  and let  $y \in Y$ . Then  $\{y\} \subseteq Y$  and  $f(f^{-1}(\{y\})) = \{y\}$ . Thus either there exists  $x \in f^{-1}(\{y\})$  such that  $f(x) \in \{y\}$ , or  $f^{-1}(\{y\}) = \emptyset$ . If the latter were true, then for all  $x \in f^{-1}(\{y\})$ ,  $f(x) \in \{y\}$  is true vacuously. Therefore, there exists  $x \in f^{-1}(\{y\}) \subseteq X$  such that  $f(x) = y$ . Thus  $f$  is onto.  $\square$

- 1.9 Let  $f: X \rightarrow Y$  be a function. Show that  $f$  is one-to-one and onto if and only if  $f(A^c) = (f(A))^c$  for all  $A \subseteq X$ .

**Proof.** Assume  $f$  is a bijection. Let  $y \in (f(A))^c$ . Then  $y \notin f(A)$  which implies that for all  $x$ , if  $f(x) = y$ , then  $x \in A^c$  which implies  $f(x) \in f(A^c)$ . Since  $y$  was arbitrary, then the previous statement holds for all  $y \in (f(A))^c$ . Moreover, since  $f$  is bijective, then for all  $y \in (f(A))^c$ , there exists  $x \in X$  such that  $f(x) = y$ . Hence, if  $y \in (f(A))^c$ , then  $y = f(x) \in f(A^c)$ . Thus,  $(f(A))^c \subseteq f(A^c)$ . Now we argue the contrapositive. Assume  $y \notin (f(A))^c$ . Then  $y \in f(A)$  which implies there exists  $x \in A$  such that  $f(x) = y$ . This implies that there exists  $x \notin A^c$  (which is unique since  $f$  is bijective) such that  $f(x) = y$  and  $y \notin f(A^c)$ . Hence,  $f(A^c) \subseteq (f(A))^c$ . Therefore  $f(A^c) = (f(A))^c$ .

Assume  $f(A^c) = (f(A))^c$  for all  $A \subseteq X$ . Let  $x_1, x_2 \in X$  such that  $x_1 \neq x_2$ . Then  $x_1 \in X \setminus \{x_2\}$  and  $x_2 \in X \setminus \{x_1\}$ . Thus  $f(\{x_1\}) \subseteq f(X \setminus \{x_2\}) \subseteq Y \setminus \{f(x_2)\}$ . Hence  $f(x_1) \neq f(x_2)$  and so  $f$  is one-to-one. Finally since  $f(X \setminus \emptyset) = f(X) = Y \setminus f(\emptyset) = Y \setminus \emptyset = Y$ , then the codomain of  $f$  is all of  $Y$ . Hence  $f$  is onto.  $\square$

- 1.10 Let  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  be one-to-one and onto functions. Show that  $g \circ f: X \rightarrow Z$  is invertible and

$$(g \circ f)^{-1} = f^{-1} \circ g^{-1}.$$

**Proof.** Let  $x_1, x_2 \in X$  such that  $(g \circ f)(x_1) = (g \circ f)(x_2)$ . Then it follows that  $g(f(x_1)) = g(f(x_2))$  and since  $g$  is one-to-one, this implies that  $f(x_1) = f(x_2)$ . Similarly, since  $f$  is one-to-one, then this implies  $x_1 = x_2$ . Hence,  $g \circ f$  is one-to-one. Now let  $z \in Z$ . Then since  $g$  is onto, there exists  $y \in Y$  such that  $g(y) = z$ . Similarly, since  $y \in Y$  and  $f$  is onto, then there exists some  $x \in X$  such that  $f(x) = y$ . Thus,  $g(f(x)) = z$  and so  $g \circ f$  is onto. Therefore  $g \circ f$  is invertible. Finally, let  $z \in Z$ . Then  $(g \circ f)^{-1}(z) = x$  for some  $x \in X$ . This means that  $f(x) = y$  for  $y \in Y$  and  $g(y) = z$ . Hence  $f^{-1}(y) = x$  and  $g^{-1}(z) = y$ . Thus

$$(f^{-1} \circ g^{-1})(z) = f^{-1}(g^{-1}(z)) = f^{-1}(y) = x.$$

Therefore  $(g \circ f)^{-1}(z) = (f^{-1} \circ g^{-1})(z)$  for all  $z \in Z$  which implies the left and right hand side are equal.  $\square$

- 1.11 Prove Proposition 1.14.

- (a) **Proof.** Let  $f: X \rightarrow X$  be defined by  $f(x) = x$ . Then  $f$  is a bijection and therefore  $X \sim X$ .  $\square$
- (b) **Proof.** Assume that  $X \sim Y$ . Then there exists a bijection  $f: X \rightarrow Y$  and since  $f$  is a bijection, then  $f^{-1}: Y \rightarrow X$  exists. Furthermore, we have that  $f^{-1}(x) = f^{-1}(y)$  gives  $f(f^{-1}(x)) = f(f^{-1}(y))$  and so  $x = y$ . Thus  $f^{-1}$  is one-to-one. Lastly, if  $x \in X$ , then  $f^{-1}(f(x)) = x$ . Hence,  $f^{-1}$  is onto. Therefore  $f^{-1}$  is one-to-one and onto and thus  $Y \sim X$ .  $\square$
- (c) **Proof.** Assume  $X \sim Y$  and  $Y \sim Z$ . Then  $f: X \rightarrow Y$  and  $g: Y \rightarrow Z$  are each bijections and by problem 1.10,  $g \circ f: X \rightarrow Z$  is a bijection. Therefore  $X \sim Z$ .  $\square$

- 1.12 Prove that the union of two countable sets is countable.

**Proof.** Let  $A$  and  $B$  be countable sets and let  $S = A \cup B$ . Since  $A$  and  $B$  are both countable, we can denote each as  $\{a_k\}$  and  $\{b_k\}$ , respectively, for  $k = 1, 2, \dots$ . Define  $f$  by

$$f(n) = \begin{cases} a_n & \text{if } 2 \mid n \\ b_n & \text{if } 2 \nmid n. \end{cases}$$

Now assume  $2 \mid n$  and that  $n = m$ . Then  $f(n) = a_n$  and  $f(m) = a_m$  from which it follows  $a_n = a_m$ . Similarly, if  $2 \nmid n$ , then  $b_n = b_m$ . Hence,  $f(n) = f(m)$  for all  $n, m \in \mathbb{N}$

and so  $f: \mathbb{N} \rightarrow S$  is well-defined. Now take some  $x_i \in S$ . Then if  $x_i \in A$ , it follows that  $2 \mid i$  and  $f(i) = x_i$ . Similarly, if  $x_i \in B$ , then  $2 \nmid i$  and  $f(i) = x_i$ . Hence,  $f$  is onto.

Finally, define  $g: S \rightarrow \mathbb{N}$  by  $g(x) = n$  such that  $n$  is the smallest natural number such that  $f(n) = x$ . This  $n$  exists by the well-ordering principle. Assume  $g(x_i) = g(x_j)$ . Then  $f(g(x_i)) = x_i = x_j = f(g(x_j))$ . Therefore  $g$  is one-to-one. Moreover since  $g(S) \subseteq \mathbb{N}$  and  $S \sim g(S)$  then by Lemma 1.15,  $S \sim g(S) \sim \mathbb{N}$ .  $\square$

1.13 Prove that the union of countably many countable sets is countable.

**Proof.** Let  $A_1, A_2, \dots$  be a countable collection of countable sets. Then we want to show that  $\bigcup_{n=1}^{\infty} A_n \sim \mathbb{N}$ . Define  $B_1 = A_1$ ,  $B_2 = A_2 \setminus A_1$ ,  $B_3 = A_3 \setminus (A_1 \cup A_2)$ ,  $\dots$ . Now note that for any  $i, j \in \mathbb{N}$  such that  $i < j$ , if  $a \in B_i$ , then

$$x \in A_i \setminus (A_1 \cup \dots \cup A_{i-1})$$

which implies that  $x \in A_i$ . However, since

$$B_j = A_j \setminus (A_1 \cup \dots \cup A_i \cup \dots \cup A_{j-1})$$

then it follows that  $x \notin B_j$ . Hence  $B_i \cap B_j = \emptyset$ .

Now we note that if  $b \in \bigcup_{n=1}^{\infty} B_n$ , then there exists some  $i \in \mathbb{N}$  such that  $b \in B_i$  and by definition, this implies that  $b \in A_i \subseteq \bigcup_{n=1}^{\infty} A_n$ . Hence,

$$\bigcup_{n=1}^{\infty} B_n \subseteq \bigcup_{n=1}^{\infty} A_n.$$

Next, we let  $a \in \bigcup_{n=1}^{\infty} A_n$  and define  $S = \{n \in \mathbb{N} \mid a \in A_n\}$ . Clearly,  $S \subseteq \mathbb{N}$  and so by the Well-Ordering Principle,  $S$  has a smallest element, call it  $n_0 = \min(S)$ . It follows that  $a \in A_{n_0}$  and  $a \notin (A_1 \cup \dots \cup A_{n_0-1})$ . Therefore

$$a \in A_{n_0} \setminus (A_1 \cup \dots \cup A_{n_0-1}) = B_{n_0} \subseteq \bigcup_{n=1}^{\infty} B_n.$$

Hence,  $\bigcup_{n=1}^{\infty} A_n \subseteq \bigcup_{n=1}^{\infty} B_n$  and therefore

$$\bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n.$$

Now we will let denote the prime numbers as  $p_1, p_2, \dots$  and define the following function:

$$f: \bigcup_{n=1}^{\infty} B_n \rightarrow \mathbb{N}$$

where, since each  $B_n$  is countable then there is a bijection  $g_n: B_n \rightarrow \mathbb{N}$ , and so for all  $b \in \bigcup_{n=1}^{\infty} B_n$ ,  $f(b) = p_n^{g_n(b)}$ . Note that  $g_n(b) \in \mathbb{N}$  which means  $f(b) \in \mathbb{N}$ . Now let  $b, b' \in \bigcup_{n=1}^{\infty} B_n$  such that  $f(b) = f(b')$ . Since  $b$  and  $b'$  are elements of the union, then

there exists  $i, j \in \mathbb{N}$  such that  $b \in B_i$  and  $b' \in B_j$ . Since we showed that  $B_i \cap B_j = \emptyset$  if  $i \neq j$ , then either  $i = j$  or  $B_i = B_j$ . If  $i \neq j$ , then  $b \neq b'$ . However, by assumption

$$f(b) = p_i^{g_i(b)} = p_j^{g_j(b')} = f(b')$$

which implies  $p_i \mid p_j^{g_j(b')}$  which further implies that for some  $k \leq g_j(b')$ ,  $p_i = p_j^m$  which is a contradiction since  $p_i \neq p_j$ . Hence,  $B_i = B_j$ . Thus  $f(b) = p_i^{g_i(b)} = p_i^{g_i(b')} = f(b')$ . This equality only holds if  $g_i(b) = g_i(b')$  and since  $g_i$  is one-to-one, then it follows that  $b = b'$ . Therefore,  $f$  is one-to-one.

Moreover, we have that

$$\bigcup_{n=1}^{\infty} B_n \sim f\left(\bigcup_{n=1}^{\infty} B_n\right) \subseteq \mathbb{N}.$$

Thus if  $f(\bigcup_{n=1}^{\infty} B_n)$  is finite, then  $\bigcup_{n=1}^{\infty} B_n$  is finite and hence countable. Otherwise, if  $f(\bigcup_{n=1}^{\infty} B_n)$  is infinite, then  $f(\bigcup_{n=1}^{\infty} B_n) \sim \mathbb{N}$  and by Proposition 1.14,  $\bigcup_{n=1}^{\infty} B_n \sim \mathbb{N}$  and therefore countable. Finally, since  $\bigcup_{n=1}^{\infty} B_n = \bigcup_{n=1}^{\infty} A_n$ , then the latter is countable.  $\square$

1.14 Let  $A$  be the collection of all sequences of the digits 0 and 1, for which the number of digits 1 is finite. Show that  $A$  is countable.

**Proof.** First we need to show that  $\mathbb{N}^n$  is countable for all  $n \in \mathbb{N}$ . Define  $f_n: \mathbb{N}^n \rightarrow \mathbb{N}$  by  $(a_1, \dots, a_n) \mapsto 2^{a_1} 3^{a_2} \cdots p_n^{a_n}$ , where  $p_n$  is the  $n$ th prime number. Now assume that  $f_n((a_1, \dots, a_n)) = f_n((b_1, \dots, b_n))$ . Then  $2^{a_1} \cdots p_n^{a_n} = 2^{b_1} \cdots p_n^{b_n}$ . Rewrite the left and right hand sides by only including powers greater than or equal to 1. From this we get

$$p_i^{a_i} \cdots p_j^{a_j} = p_r^{b_r} \cdots p_s^{b_s}.$$

Wlog, assume that  $j - i \leq s - r$ . Clearly,  $p_i^{a_i}$  divides the left and right hand sides and so either  $p_i^{a_i} \mid p_r^{b_r}$  or  $p_i^{a_i} \nmid p_r^{b_r}$ . If the former is true, then  $p_i = p_r$  and  $a_i \leq b_r$ . Thus dividing both sides by  $p_i^{a_i}$  we obtain

$$1 \cdots p_j^{p_j} = p_r^{b_r - a_i} \cdots p_s^{b_s}.$$

However, if  $a_i < b_r$  then the right hand side and left hand side are both multiples of  $p_r$ , but this is a contradiction since the left hand side no longer contains any power of  $p_r$ . Hence, if  $p_i^{a_i} \mid p_r^{b_r}$ , then  $a_i = b_r$ . Additionally, if  $p_i^{a_i} \nmid p_r^{b_r}$ , then either  $p_i < p_r$  or  $p_i > p_r$ . If  $p_i < p_r$ , then the left hand side is a multiple of  $p_i$  whereas the right is not which is a contradiction. Similarly, the same argument holds if  $p_i > p_r$ . Thus,  $p_i = p_r$  and  $a_i = b_r$ . Finally, if  $j - i < s - r$ , then dividing the left and right hand side by the left hand side we obtain

$$1 = p_t^{b_t} \cdots p_v^{b_v}$$

which is a contradiction since all exponents were greater than or equal to 1. Hence,  $j - i = s - r$  and thus  $a_i = b_i$  for all  $1 \leq i \leq n$ . Therefore,  $f_n$  is one-to-one. Finally, letting  $x \in \mathbb{N}$ , by the fundamental theorem of arithmetic, we can

uniquely express  $x$  as  $2^{a_1} \cdots p_n^{a_n}$  and hence  $f_n((a_1, \dots, a_n)) = x$ . Thus  $f_n$  is onto. Thus  $\mathbb{N}^n \sim \mathbb{N}$  and so  $\mathbb{N}^n$  is countable.

Let  $A_n \subseteq A$  be the set of all sequences in  $A$  which contain  $n$  many 1's. Then define the function  $g_n: A_n \rightarrow \mathbb{N}^n$  in the following way: If  $a \in A$  and  $a = a_1, a_2, \dots$ , then

$$a_1, a_2, \dots \mapsto (\alpha_1, \alpha_2, \dots, \alpha_n)$$

if and only if  $a_{\alpha_i} = 1$  for all  $i = 1, 2, \dots, n$  and 0 otherwise. Assume that  $g_n(a) = g_n(b)$  for some  $a, b \in A_n$ . Then  $(\alpha_1, \dots, \alpha_n) = (\beta_1, \dots, \beta_n)$  which implies that  $\alpha_i = \beta_i$  for all  $i$ . Hence, for each  $a_i$  and  $b_i$ , we have that either  $a_i = b_i = 0$  or  $a_{\alpha_i} = b_{\alpha_i} = 1$ . Therefore,  $a = b$  and  $g_n$  is one-to-one. Next, if we take  $(\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ , then if  $a$  is the sequence for which  $a_{\alpha_1}, \dots, a_{\alpha_n}$  are all 1 and the rest are 0, then  $g_n(a) = (\alpha_1, \dots, \alpha_n)$  and  $g_n$  is onto. Thus  $A_n \sim \mathbb{N}^n$  and so  $A_n$  is countable for all  $n$ .

The last part of this argument is to show that  $A = \bigcup_{n=1}^{\infty} A_n$ . If  $a \in A$ , then  $a$  contains  $m$  many 1's for some  $m \in \mathbb{N}$  and thus  $a \in A_m \subseteq \bigcup_{n=1}^{\infty} A_n$ . Conversely, if  $a \in \bigcup_{n=1}^{\infty} A_n$ , then for some  $m \in \mathbb{N}$ ,  $a \in A_m$  and thus  $a$  is a sequence of digits 0 and 1 for which there are  $m$  many digits 1. Hence,  $a \in A$  and the equality has been shown. Thus, since for each  $n \in \mathbb{N}$ ,  $A_n$  is countable, then by problem 1.13,

$$A = \bigcup_{n=1}^{\infty} A_n$$

is countable. □

- 1.15 Let  $X$  be the set of all numbers from  $[0, 1]$  whose decimal expansion contains only the digits 3 or 5. Is  $X$  countable or uncountable?

**Solution.** Assume that  $X$  is countable. Then we can list each element of  $X$  in the following way:  $x_1, x_2, \dots$ , where  $x_i = 0.x_{i1}x_{i2}\dots$  for all  $i \in \mathbb{N}$ . In other words,  $x_{mn}$  is the  $n$ th digit of the  $m$ th decimal expansion. Now define a new decimal expansion  $y = 0.y_1y_2\dots$  such that

$$y_i = \begin{cases} 3 & \text{if } x_{ii} = 5 \\ 5 & \text{if } x_{ii} = 3 \end{cases}.$$

We can see that  $y \in [0, 1]$  and that each of its digits are either 3 or 5. Hence,  $y \in X$ . However, we claim that  $y$  is not in the list provided above. For if we assume that there is some  $x_t$  such that  $y = x_t$ , then this implies  $y_i = x_{ti}$  for all  $i$ . But this is not the case since by construction  $y_t \neq x_{tt}$ . Therefore  $y$  is not in the list above and the elements of  $X$  cannot be enumerated. Hence  $X$  is uncountable.



1.17 Show that if  $\text{card}(X) = n$ , then  $\text{card}(2^X) = 2^n$ .

**Proof.** We proceed by induction on  $n \in \mathbb{N}$ . Define

$$P(n) := \text{card}(X) = n \rightarrow \text{card}(2^X) = 2^n.$$

For the base case, we assume  $\text{card}(X) = 0$ . Then  $X = \emptyset$  and thus  $2^X = \{\emptyset\}$ . Hence  $\text{card}(2^X) = 1 = 2^0$ . Thus  $P(1)$  holds.

Now assume  $P(k)$  holds for some  $k \geq 2$ . Then  $\text{card}(X) = k$  implies  $\text{card}(2^X) = 2^k$ . Let  $Y$  be a set such that  $\text{card}(Y) = k + 1$  and take  $y \in Y$  and set  $\hat{Y} = Y \setminus \{y\}$ . Then clearly,  $\text{card}(\hat{Y}) = k$  and hence  $\text{card}(2^{\hat{Y}}) = 2^k$ . It follows that for any subset  $S \subseteq Y$ , that  $y \in S$  or  $y \notin S$ . Moreover, in collecting all the subsets  $S \subseteq Y$  such that  $y \notin S$ , we find that there are  $2^k$  of them since this collection is precisely  $2^{\hat{Y}}$ . It follows then that  $Y \setminus \hat{Y}$  contains all the elements of  $\hat{Y}$  but with  $y$  as a member, of which there are  $2^k$  such elements. Hence,  $\text{card}(Y) = 2 \cdot 2^k = 2^{k+1}$ . Thus  $P(k+1)$  holds. Therefore  $P(n)$  holds for all  $n \in \mathbb{N}$ .  $\square$

1.18 Show that  $2^{\mathbb{N}} \sim \mathbb{R}$ .

**Proof.** First note that since  $\mathbb{N} \in 2^{\mathbb{N}}$ , then  $2^{\mathbb{N}}$  is not finite. Next, assume for contradiction that  $2^{\mathbb{N}} \sim \mathbb{N}$ . Then there exists a bijection  $h: \mathbb{N} \rightarrow 2^{\mathbb{N}}$  such that  $h(n) = S_n$ . Now define a function  $f: 2^{\mathbb{N}} \rightarrow [0, 1]$  in the following way: for any  $S_i \in 2^{\mathbb{N}}$

$$f(S_i) = \sum_{n=1}^{\infty} \frac{s_{in}}{10^n}; \quad s_{in} = \begin{cases} 3 & \text{if } n \in S_i \\ 5 & \text{if } n \notin S_i \end{cases}.$$

We want to show this function is one-to-one. Assume for  $S_i, S_j \in 2^{\mathbb{N}}$  that  $f(S_i) = f(S_j)$ . Then  $0.s_{i1}s_{i2}\dots = 0.s_{j1}s_{j2}\dots$  which implies  $s_{ik} = s_{jk}$  for all  $k \in \mathbb{N}$ . Thus, if  $k \in S_i$ , then  $s_{ik} = 3 = s_{jk}$  and hence  $k \in S_j$ . Thus  $S_i \subseteq S_j$ . Similarly, it follows by the same reasoning that  $S_j \subseteq S_i$  and thus  $S_i = S_j$ . Therefore  $f$  is one-to-one.

Define  $r = 0.g_1g_2\dots$ , where  $g_i = 3$  if  $s_{ii} = 5$  or  $g_i = 5$  if  $s_{ii} = 3$ . Assume there exists some  $S_i \in 2^{\mathbb{N}}$  such that  $f(S_i) = r$ . Then  $0.g_1g_2\dots g_i\dots = 0.s_{i1}s_{i2}\dots s_{ii}\dots$ , which implies  $g_i = s_{ii}$ , but this is not possible by construction of  $r$  and hence for all  $S_i \in 2^{\mathbb{N}}$ ,  $f(S_i) \neq r$ . Finally, define  $S' = \{n \in \mathbb{N} \mid g_n = 3\}$ , where  $g_n$  is the  $n$ th digit in the decimal expansion  $r$ . Then it follows by construction that  $f(S') = r$ . However,  $S' \in 2^{\mathbb{N}}$  yet  $f(S') = r \neq f(S_i)$  for all  $i \in \mathbb{N}$  which implies  $S' \neq S_i$  for all  $i \in \mathbb{N}$ . Hence  $S' \notin h(\mathbb{N})$  which is a contradiction.

Therefore  $2^{\mathbb{N}} \not\sim \mathbb{N}$ . Thus, by the Continuum Hypothesis,  $\text{card}(2^{\mathbb{N}}) \geq \text{card}(\mathbb{R})$ . However, since  $f$  is one-to-one, then  $2^{\mathbb{N}} \sim f(2^{\mathbb{N}}) \subseteq [0, 1] \sim \mathbb{R}$  and hence  $\text{card}(2^{\mathbb{N}}) \not\geq \text{card}(\mathbb{R})$  which implies  $\text{card}(2^{\mathbb{N}}) = \text{card}(\mathbb{R})$ . Therefore  $2^{\mathbb{N}} \sim \mathbb{R}$ .  $\square$

## 1.19 Prove Proposition 1.27.

**Proof.** Assume  $\alpha^* = \inf A$ . Then by Definition 1.26,  $\alpha^*$  is a lower bound of  $A$  (part (i) of Proposition 1.27) and if  $\alpha$  is a lower bound of  $A$ , then  $\alpha \leq \alpha^*$ . Let  $B$  be the set of all lower bounds of  $A$ . By definition, for any  $b \in B$ ,  $b \leq \alpha^*$ . Conversely, if  $b \leq \alpha^*$ , then  $b$  is a lower bound of  $A$  and hence  $b \in B$ . Now let  $\varepsilon > 0$ . It follows that  $\alpha^* < \alpha^* + \varepsilon$ . Thus  $\alpha^* + \varepsilon \notin B$ , otherwise  $\alpha^* + \varepsilon \leq \alpha^*$  which is not the case. Hence,  $\alpha^* + \varepsilon$  is not a lower bound of  $A$ . Thus there exists some  $a \in A$  such that  $a < \alpha^* + \varepsilon$ . Moreover, since  $a \in A$  then  $\alpha^* \leq a$  and thus  $\alpha^* \leq a < \alpha^* + \varepsilon$  (part (ii) of Proposition 1.27).

Conversely, assume that  $\alpha^*$  is a lower bound of  $A$  and that for all  $\varepsilon > 0$ , there exists  $a_\varepsilon \in A$  such that  $\alpha^* \leq a_\varepsilon < \alpha^* + \varepsilon$ . From the second condition it follows that  $\alpha^* - \varepsilon \leq \alpha^*$  for all  $\varepsilon > 0$ . Then if  $b$  is any lower bound of  $A$ , then there exists  $\varepsilon > 0$  such that  $b + \varepsilon = \alpha^*$  and hence  $b = \alpha^* - \varepsilon \leq \alpha^*$ . Therefore  $\alpha^* = \inf A$ .  $\square$

## 1.20 Prove Proposition 1.29.

**Proof.** Let  $\beta^* = \sup A$ . Then  $\beta^*$  is an upper bound of  $A$  (condition (i) of the proposition) and for any upper bound  $a$  of  $A$ ,  $\beta^* \leq a$ . Let  $B$  be the set of all upper bounds of  $A$ . Then by definition if  $b \in B$ ,  $\beta^* \leq b$ . Conversely, if  $\beta^* \leq b$ , then  $b$  is an upper bound of  $A$  and hence  $b \in B$ . Let  $\varepsilon > 0$ . Then  $\beta^* - \varepsilon < \beta^*$  and hence  $\beta^* - \varepsilon \notin B$ . Thus  $\beta^* - \varepsilon$  is not an upper bound of  $A$ . This implies that there exists  $a \in A$  such that  $\beta^* - \varepsilon < a$ . But since  $a \in A$ , then  $a \leq \beta^*$ . Hence  $\beta^* - \varepsilon < a \leq \beta^*$  (part (ii) of the proposition).

Conversely, assume that  $\beta^*$  is an upper bound of  $A$  and that for all  $\varepsilon > 0$ , there exists  $b_\varepsilon \in A$  such that  $\beta^* - \varepsilon < b_\varepsilon \leq \beta^*$ . Then  $\beta^* \leq \beta^* + \varepsilon$  for all  $\varepsilon > 0$ . Thus if  $b$  is any upper bound of  $A$  then  $\beta^* \leq b$  and so there exists  $\varepsilon > 0$  such that  $\beta^* - \varepsilon = b$ . Hence  $\beta^* \leq b = \beta^* - \varepsilon$ . Therefore  $\beta^* = \sup A$ .  $\square$

1.21 Let  $A \subseteq \mathbb{R}$  be a nonempty set which is bounded from above. Show that if  $\sup A \notin A$ , then for all  $\varepsilon > 0$  the open interval  $(\sup A - \varepsilon, \sup A)$  contains infinitely many terms of  $A$ .

**Proof.** Let  $\varepsilon > 0$  and define  $S = (\sup A - \varepsilon, \sup A)$ . Assume for contradiction that for some  $n \in \mathbb{N}$   $S$  is finite and  $S = \{s_1, s_2, \dots, s_n\}$ . Being finite implies that  $S$  has a maximum, call it  $s = \max(S)$ . Let  $R = \{\sup A - s_i : s_i \in S\}$ . Note that  $R \subseteq \mathbb{R}$  and  $R$  is nonempty since otherwise  $\sup A - (\sup A - \varepsilon) = 0$  which would imply  $\varepsilon = 0$ . Further, note that  $R$  is bounded below by 0 since if for some  $1 \leq i \leq n$ ,  $\sup A - s_i < 0$  then  $\sup A < s_i$  which implies  $\sup A$  is not an upper bound of  $A$ . Therefore by Axiom 5,  $\inf R$  exists.

We claim that  $\inf R = \sup A - s$ . First, let  $r \in R$ , then for some  $s_i \in S$ ,  $r = \sup A - s_i$ . If  $\sup A - s_i < \sup A - s$  then  $s < s_i$  which is a contradiction since  $s = \max(S)$ . Thus  $\sup A - s$  is a lower bound of  $R$ . Let  $b$  be a lower bound of  $R$ . Then for all  $s_i \in S$ ,  $b \leq \sup A - s_i$  and hence  $b \leq \sup A - s$ . Therefore  $\sup A - s = \inf R$ . Finally, by Proposition 1.20 there exists  $s' \in S$  such that  $\sup A - s < s' \leq \sup A$ . Thus  $s > \sup A - s'$ . But this contradicts that  $s = \inf R$ . Therefore  $S$  is not finite and  $S$  contains infinitely many terms of  $A$ .  $\square$

1.21 Let  $A, B \subseteq \mathbb{R}$  be nonempty bounded sets, and let  $c \in \mathbb{R}$ . Define the following sets:

$$A + B = \{a + b \mid a \in A, b \in B\}$$

$$A - B = \{a - b \mid a \in A, b \in B\}$$

$$A \cdot B = \{ab \mid a \in A, b \in B\}$$

$$cA = \{ca \mid a \in A\}.$$

Prove that:

(1)  $\inf(A + B) = \inf(A) + \inf(B)$

**Proof.** Let  $\alpha = \inf(A + B)$ ,  $\alpha_1 = \inf A$ , and  $\alpha_2 = \inf B$ . Assume  $\alpha < \alpha_1 + \alpha_2$ . Let  $a \in A, b \in B$ . Then  $\alpha \leq a + b$ ,  $\alpha_1 \leq a$ , and  $\alpha_2 \leq b$ . Thus  $\alpha_1 + \alpha_2 \leq a + b$ . Hence  $\alpha_1 + \alpha_2$  is a lower bound of  $A + B$ . Thus  $\alpha_1 + \alpha_2 \leq \alpha$  which is a contradiction. Assume  $\alpha_1 + \alpha_2 < \alpha$ . Let  $a \in A, b \in B$ . Then  $\alpha \leq a + b$  and  $\alpha_2 \leq b$ . Hence  $\alpha - \alpha_2 \leq a$ . Thus  $\alpha - \alpha_2$  is a lower bound of  $A$  which implies  $\alpha - \alpha_2 \leq \alpha_1$ . Thus  $\alpha \leq \alpha_1 + \alpha_2$  which is a contradiction. Therefore  $\alpha = \alpha_1 + \alpha_2$ , or  $\inf(A + B) = \inf A + \inf B$ .  $\square$

(3)  $\sup(-A) = -\inf(A)$

**Proof.** Let  $\alpha = \sup(-A)$  and  $\beta = -\inf(A)$ . For any  $a \in A$  we have that  $-a \leq \alpha$  and  $-\beta \leq a$  which implies  $-a \leq \beta$ . Thus  $\beta$  is an upper bound of  $-A$  and so  $\alpha \leq \beta$ . Similarly,  $-\alpha \leq a$  and so  $-\alpha$  is a lower bound of  $A$ . Thus  $-\alpha \leq -\beta$ . Hence  $\beta \leq \alpha$ . Therefore  $\alpha = \beta$ .  $\square$

(5)  $\sup(A - B) = \sup A - \inf B$ .

**Proof.** Let  $C = -B$ . Then by (1) and (2) it follows that

$$\sup(A - B) = \sup(A + C) = \sup A + \sup C = \sup A + \sup(-B) = \sup A - \inf B.$$

$\square$

(7)  $\sup(cA) = c \sup A$  if  $c > 0$ .

**Proof.** Let  $ca \in cA$ . Then  $ca \leq \sup(cA)$ . Similarly, for any  $a \in A$  we have that  $a \leq \sup A$  and thus for any  $c > 0$ ,  $ca \leq c \sup A$ . Thus  $c \sup A$  is an upper bound of  $cA$  and hence  $\sup(cA) \leq c \sup A$ . Also note that since  $ca \leq \sup(cA)$  for all  $c > 0$  and  $a \in A$ , then  $a \leq \frac{1}{c} \sup(cA)$  and hence  $\frac{1}{c} \sup(cA)$  is an upper bound of  $A$ . As such it follows that  $\sup A \leq \frac{1}{c} \sup(cA)$  and hence  $c \sup A \leq \sup(cA)$ . Therefore  $c \sup A = \sup(cA)$ .  $\square$

(9)  $\sup(cA) = c \inf A$  if  $c < 0$ .

**Proof.** Let  $ca \in cA$ . Then  $ca \leq \sup(cA)$  and so  $\frac{1}{c} \sup(cA) \leq a$ . Hence  $\frac{1}{c} \sup(cA)$  is a lower bound of  $A$ . Thus  $\frac{1}{c} \sup(cA) \leq \inf A$ . Additionally, if  $a \in A$  then  $\inf A \leq a$  and so  $ca \leq c \inf A$ . Thus  $c \inf A$  is an upper bound of  $cA$  which implies  $\sup(cA) \leq c \inf A$ . Hence  $\sup A \leq \frac{1}{c} \sup(cA)$ . Since  $\inf A \leq \sup A$  then  $\inf A \leq \frac{1}{c} \sup(cA)$ . Thus  $\frac{1}{c} \sup(cA) = \inf A$  and hence  $\sup(cA) = c \inf A$ .  $\square$

(11) Is it true that  $\sup(A \cdot B) = \sup A \cdot \sup B$ ?

**Solution.** Let  $A = [-2, -1]$  and  $B = [0, 1]$  then  $A \cdot B = [-2, 0]$ . Thus  $\sup(A \cdot B) = 0$  and  $\sup A \cdot \sup B = (-1)(1) = -1$ . Thus this claim is not true.

1.23 State and prove the density property of the irrational numbers.

**Proof.** If  $x, y \in \mathbb{R}$  and  $x < y$ , then there exists  $i \in \mathbb{I}$  such that  $x < i < y$ .

By Theorem 1.33, for any  $x, y \in \mathbb{R}$  with  $x < y$  there exists  $p \in \mathbb{Q}$  such that  $x < p < y$ . Moreover, by Corollary 1.32 there exists  $n \in \mathbb{N}$  such that  $\frac{1}{n} < \frac{y-p}{2}$ . Thus  $p + \frac{\sqrt{2}}{n} < p + \frac{2}{n} < y$  and since  $x < p$  then  $x < p + \frac{\sqrt{2}}{n}$ . Hence  $x < p + \frac{\sqrt{2}}{n} < y$ . Now assume that  $p + \frac{\sqrt{2}}{n} \in \mathbb{Q}$ . Then there exists  $a, b \in \mathbb{Z}$  such that  $b \neq 0$ ,  $\gcd(a, b) = 1$ , and  $p + \frac{\sqrt{2}}{n} = \frac{a}{b}$ . Hence  $\sqrt{2} = \frac{(a-bp)n}{b} \in \mathbb{Q}$ . However this contradicts problem 1.3. Therefore  $p + \frac{\sqrt{2}}{n} \in \mathbb{I}$ .  $\square$