


Modelling palaeoecological community data: a state-space approach

Quinn Asena, Anthony Ives, Jack Williams. and Jonathan Johnson

UW-Madison

2023-08-06

Is the past recoverable from the data?



Palaeoecologists are concerned with questions such as:

- what are the drivers of community change?
- role species-environment interactions?
- role of density dependence and species interactions?
- how can palaeoecological information inform present and future ecosystem states?

Palaeoecological proxy data

- Typically comprise multiple time-series (response):
 - e.g., pollen, diatoms...
- Include environmental covariates (predictors):
 - e.g., isotopic data, charcoal data, lake level...

Palaeoecological data pose many statistical challenges:

- Uneven sampling through time
- Time averaging
- Measurement uncertainty
- Relative abundances

Descriptive approaches

Many descriptive approaches exist for analysing multivarivariate time-series:

- Cluster analyses: CONISS
- Ordination: PCA, NMDS...
- Machine learning: MVRT, LDA (fancy cluster analyses)

Descriptive methods allow us to extract patterns from the data but not determine potential causes of those patterns.

Find image of cluster analysis

Beyond pattern recognition

The cutting edge in palaeoecology is to establish potential causes of observed patterns in species relative abundances. For example, are observed patterns driven by:

- species interactions?
- climate variability?
- fire regime?
- ...

This is what we want to know if we are to use palaeoecology to inform management of contemporary ecosystems or inform potential future ecosystem states. No easy task!

State-space modelling

State-space modelling goes beyond descriptive approaches and attempts to estimate:

- autoregressive / density dependent processes
- interspecific interactions (C matrix)
- species-environment interactions (B vector)
- combinations of the above

State-space modelling

State-space modelling attempts to predict the “true” unobservable state of a system from observable variables. It does so via two equations, one that models the *process of the system*:

Process equation:

Something not right here Tony...:

$$Y = B0 + C(Y - B0 - BX) + BX$$

and one that models the *observations* from the system:

Observation equation:

$$Y_t = Multinomial(z_t)$$

State-space modelling

State-space models are not new to ecology and have been used for:

- estimating animal populations
- animal movement
- plant cover data
- and much more

However, state-space models are not well explored in palaeoecology.

State-space modelling

include eq on slide:

This new variant of a state-space model:


- uses a multinomial distribution: accepts raw count data
- accepts multiple predictor data streams in the same model: e.g., isotopic data, charcoal accumulation rates, fungal spores...
- simultaneously fits multiple coefficients
- models autocorrelation structure

Can be used to assess a range of possible causes of observed patterns in palaeo-data.

Empirical example

Demonstrating a three-taxon model from Sunfish Pond:

- unpublished data: not presenting the dataset, focusing on the modelling approach (Johnson *et al.*, unpub)
- ACES project interested in abrupt transitions between dominant species



Jonathon Johnson (JJ)




Empirical example

This example is a three-taxon model:

- two focal species (*Betula* and *Quercus*)
- third 'species' is an aggregate of all other species
- we fit species interactions
- estimate species change through time

Remember, this is a multinomial problem so the we do not estimate the aggregate species directly.

Time-forward model enables even intervals




Johnson *et al.*, unpublished

Species interaction estimates

C matrix

	other	Quercus	Betula
other	-0.074	0.000	0.000
Quercus	0.000	-0.006	0.121
Betula	0.000	-0.702	-0.459

- columns = abundance; rows = change in abundance
- density dependence on the diagonal
- *Quercus-Quercus* near 0 means abundance of *Quercus* has little effect on the change in *Quercus*
- *Quercus-Betula* -0.7 means that abundance of *Quercus* affects the change in *Betula* abundance



Estimating the effect of covariates


Estimate of change over time

B vector

```
other Quercus Betula  
0    0.048 -0.532
```

Overall:

- *Quercus* increases with time
- *Betula* decreases with time
- estimates are relative!



Evaluating the model with simulated data

- We cannot determine the accuracy of fitted coefficients empirically from palaeoecological data.
- Simulation experiments are used to assess the success of recovering parameters:
 1. data are simulated under known conditions with replication
 2. the model is fit to the simulated replicate datasets
 3. fitted models are assessed for how well input parameters are recovered

Evaluating the model with simulated data

C matrix fit to the pollen data

	other	Quercus	Betula
other	-0.074	0.000	0.000
Quercus	0.000	-0.006	0.121
Betula	0.000	-0.702	-0.459

B vector fit to the pollen data

other	Quercus	Betula
0	0.048	-0.532

C matrix fit to the simulated data

oh

B vector fit to the simulated data

no!

- 100 reps
- average difference between input and fitted parameters

Hypothesis testing

We cannot determine with certainty, outside of simulation, causation from palaeo-data.

What we can do is:

- set up multiple working hypotheses (Chamberlin, 1897)
- couple descriptive methods with inferential ones
- assess which results lend support to the likelihood of each hypothesis being true

Given the data at hand the interaction matrix (C) indicates some competition between *Quercus* and *Betula*. Such an inference lends support to one hypothesis.

Acknowledgements

Aces team: - Jack Williams, Tony Ives, Angie Perotti, Nora Schlenker, Sam Wiles, Amanda Toomey, Adrian George, Bryan Schuman, David Nelson, Jonathon Johnson

- National Science Foundation
- UW-Madison

