

Assignment 1

Quinn Perfetto, 104026025
60-454 Design and Analysis of Algorithms

January 24, 2017

Question 1. (i). We want to prove that when Algorithm Summation terminates its execution,

$$y = \sum_{j=0}^n a_j x^j$$

Proof. We shall apply induction to prove that after the m th iteration of the for loop the following invariant holds true:

$$y = \sum_{j=0}^{m-1} a_{n-j} x^{n-i-j}$$

(Induction basis) First, we note that y is initialized to 0 in Line 1. When $m = 1$ and $i = n$ in Line 3,

$$\begin{aligned} y &= a_n + x * y \\ &= a_n + x * 0 \\ &= a_n \\ &= a_n x^0 \\ &= \sum_{j=0}^0 a_{n-j} x^{n-i-j} \end{aligned}$$

(Induction Hypothesis) Suppose for iteration $m - 1 < n$, $y = \sum_{j=0}^{m-2} a_{n-j} x^{n-i-j-1}$
(Induction Step) When Line 3 is executed for the m th time,

$$\begin{aligned}
y &= a_{n-i} + x \sum_{j=0}^{m-2} a_{n-j} x^{n-i-j-1} \\
&= a_{n-i} + \sum_{j=0}^{m-2} a_{n-j} x^{n-i-j} \\
&= a_{n-i} x^0 + \sum_{j=0}^{m-2} a_{n-j} x^{n-i-j}
\end{aligned}$$

(Induction Hypothesis)

$$= \sum_{j=0}^{m-1} a_{n-j} x^{n-i-j}$$

Therefore we can conclude that the invariant $y = \sum_{j=0}^{m-1} a_{n-j} x^{n-i-j}$ holds $\forall m > 0$.

On the $n+1$ th iteration (where $i = 0$) we then have,

$$\begin{aligned}
y &= \sum_{j=0}^n a_{n-j} x^{n-j} \\
&= a_n x^n + a_{n-1} x^{n-1} + \dots + a_1 x^1 + a_0 x^0 \\
&= \sum_{j=0}^n a_j x^j
\end{aligned}$$

(By reversing the order of the terms)

□

Question 1. (ii).

Key Operation: Multiplication of real numbers.

Input size: n (The size of the input array)

The Summation algorithm must perform the key operation on each of the n elements of the input array.

1. Worst-case Time Complexity: $T(n) = n$
2. Average-case Time Complexity: $T_{ave}(n) = n$