

Praktikum: P4 Gruppe: 22

☒ **Mo** ☐ **Mi**
Zutreffendes bitte ausfüllen

WS20/21

Namen: Paul Filip useba[at]student.kit.edu

Namen: Janic Beck

Versuch: Mößbauer-Effekt

Betreuer: Paras Koundal Durchgeführt am: 29.11.20

Wird vom Betreuer ausgefüllt.

1. Abgabe am: _____

Rückgabe am: _____ Kommentar:

2. Abgabe am: _____

Ergebnis: + / 0 / - Handzeichen: _____

Datum: _____ Kommentar:


Contents

1	Theory & Preparation	1
1.1	Mössbauer effect	1
1.2	Mössbauer spectroscopy	2
2	Experiment & Evaluation	3

1. Theory & Preparation

1.1 Mössbauer effect

The process of **resonant absorption** in nuclear physics describes the phenomenon of subsequent de- and excitation of two equal atoms to the same energy levels via one γ -quant. Consider for example an excited state of ^{57}Fe , that emits a photon with energy (roughly) 14.4 keV during its transition to the ground state.



In principle, one could use this emitted photon to excite another ^{57}Fe atom to the higher energy state. The photon is absorbed resonantly by the second atom during this process.

In reality, resonant absorption such as the Na-D-line only occurs under certain circumstances. Due to conservation laws the energy E_γ of the emitted photon does not exactly equal the transition energy E_0 , but is instead shifted downward by the nuclear recoil energy. A similar analysis finds that the energy for absorption of the same atom is shifted upwards.

$$\underbrace{E_\gamma = E_0 - \frac{p_\gamma^2}{2m}}_{\text{Emission}} \qquad \underbrace{E_\gamma = E_0 + \frac{p_\gamma^2}{2m}}_{\text{Absorption}} \qquad (1.1)$$

With the photon impulse p_γ and atom mass m . If the line width introduced by natural broadening or other effects does not exceed the energy gap, resonant absorption cannot occur (see Figure 1.1). It is also notable that the energy gap between emission and absorption spectrum can be increased by additional effects. This will be further discussed in chapter 2.




Figure 1.1: (a) The natural linewidth Γ_0 is not sufficient for a sizeable overlap of both spectra. Resonance absorption is not possible. (b) The line width of both spectra can be increased by other effects such as Doppler broadening. In such cases the spectra with linewidth $\Gamma_0 + \Gamma_D$ can overlap and resonant absorption is possible.

As it turns out, the above rules stating when resonant absorption can and cannot occur are not strictly true. Experiments in the 1960s conducted by R. Mössbauer ([?]) showed that resonant absorption in a crystal lattice happens much more readily than one would expect based on the previous discussion. The difference is the tight binding of the atoms in the crystal lattice. Instead of the individual atom recoiling, different phonons can be created (or destroyed) by the emission and absorption. In a sense, the entire crystal absorbs the recoil energy, effectively substituting the atom mass m in equation Equation 1.1 by the mass M of the entire crystal. Because $m \ll M$, the energy gap between emission and absorption spectrum drastically decreases. This phenomenon of recoilless nuclear resonant absorption is named **Mössbauer effect**.

1.2 Mössbauer spectroscopy

Es war einmal ...

2. Experiment & Evaluation