

Modern Algebra 1 - MATH 6302

Joel Sleeba
University of Houston
joelsleeba1@gmail.com

September 30, 2024

Contents

Contents	1
1	2
1.1 Internal Direct products	2
2	4

Chapter 1

1.1 Internal Direct products

Suppose G is a group and $H, K \leq G$. Define

$$HK = \{hk : h \in H, k \in K\}$$

Theorem 1.1.1. Suppose $|H|, |K| < \infty$, then $|HK| = \frac{|H||K|}{|H \cap K|}$.

Proof. We write $HK = \cup_{h \in H} hK$. Then $h_1K = h_2K$ if and only if $h_2^{-1}h_1 \in K$. This is equivalent to saying $h_2^{-1}h_1 \in H \cap K$. verify \square

Theorem 1.1.2. If $H, K \leq G$ then $HK \leq G$ if and only if $HK = KH$.

Corollary 1.1.2.1. If H is a normal subgroup of G , then $HK \leq G$

Theorem 1.1.3. If $H, K \leq G$, $|H|, |K| < \infty$, H, K normal in G , and $H \cap K = \{e\}$, then $HK \cong H \times K$. In this case HK is called the internal direct product of H and K .

Proof. By the corollary, $HK \leq G$. Therefore consider the map $\phi : HK \rightarrow H \times K : hk \rightarrow (h, k)$.

- ϕ is well defined. We know that $|HK| = \frac{|H||K|}{|H \cap K|} = |H||K|$. Hence every element in HK has a unique representation as hk . Thus we see that ϕ is well defined.
- ϕ is a homomorphism. Let $h_1k_1, h_2k_2 \in HK$. Want to show that $\phi(h_1k_1h_2k_2) = \phi(h_1k_1)\phi(h_2k_2) = (h_1, k_1)(h_2, k_2) = (h_1h_2, k_1k_2)$. Equivalently we want to show that $h_1k_1h_2k_2 = h_1h_2k_1k_2$, which is again equivalent to showing $hk = kh$ for all $h \in H, k \in K$. This is again equivalent to showing $[h, k] = h^{-1}k^{-1}hk = e$ for all $h \in H, k \in K$.

To see why $[h, k] = \{e\}$. see that $[h, k] = h^{-1}(k^{-1}hk) = h^{-1}h' \in H$ for some $h' \in H$. Similarly $[h, k] = (h^{-1}k^{-1}h)k = k'k \in K$ for some $k' \in K$.


Both of these are by the normality of H and K . But since we know that $H \cap K = \{e\}$, we see that $[h, k] = e$ for all h, k . Hence we are done.

□

Chapter 2

Example 2.0.1. Let $G = D_{12}$, $H = \langle r \rangle$, $K = \langle s \rangle$. Then $H \cap K = \{e\}$, so $|HK| = 12$ and $HK \cong D_{12}$, but $HK \not\cong H \times K$ since $H \times K$ is Abelian and D_{12} is not.

Theorem 2.0.1 (Diamond Isomorphism Theorem). *Suppose $A, B \leq G$ and $A \subseteq N_G(B)$, then $AB \leq G$, $B \triangleleft AB$, $A \cap B \trianglelefteq A$ and $AB/B \cong A/A \cap B$*



Proof. verify

□

Theorem 2.0.2. *Let $H, K \trianglelefteq G$ and $H \leq K$, then $H \trianglelefteq K$, $K/H \trianglelefteq G/H$,*

$$(G/H)/(K/H) \cong G/K$$