

Professional Anatomical Treatment Plan

Analysis Details:

Target Muscle Group: Corrugator Supercilii

Analysis Type: weakness

Primary Treatment: Dermal Fillers


Treatment Intensity: moderate

Session Frequency: weekly

Primary symptoms: numbness

Severity Scale: 3

Follow-up Plan: 6_months



PROFESSIONAL ANATOMICAL TREATMENT PLAN

Target Area: CORRUGATOR SUPERCILII

Analysis Type: weakness

Primary Treatment: dermal fillers

CLINICAL ASSESSMENT:

The patient presents with symptoms affecting the corrugator supercilii region. Based on the weakness analysis, the following treatment protocol is recommended.

Primary Symptoms: numbness

Severity Rating: 3

TREATMENT PROTOCOL:

1. Initial Assessment

- Comprehensive evaluation of corrugator supercilii function and condition
- Baseline documentation with clinical photography
- Patient history and contraindication screening

2. Primary Treatment: dermal fillers

- Treatment Intensity: moderate
- Session Frequency: weekly
- Progressive monitoring of muscle response and patient tolerance
- Adjustment of treatment parameters based on individual response

3. Monitoring and Follow-up

- Regular assessment of treatment efficacy
- Documentation of progressive improvements
- 6_months

PATIENT EDUCATION:

- Expected timeline for visible results: 7-14 days
- Potential side effects and their management
- Post-treatment care instructions
- Activity restrictions and recommendations

SAFETY CONSIDERATIONS:

- Patient should be advised of all potential risks and contraindications
- Informed consent must be obtained prior to treatment
- Emergency protocols should be reviewed with patient
- Contact information for post-treatment concerns provided

This treatment plan should be reviewed and adjusted based on individual patient response and clinical judgment.