Averox Pvt Ltd
Averox Diplomatic Enclave Islamabad
315548282020
hassan@averox.com

Professional Anatomical Treatment Plan



Analysis Details:
Target Muscle Group: Corrugator Supercilii
Analysis Type: hyperactivity
Primary Treatment: Dermal Fillers
Treatment Intensity: moderate
Session Frequency: twice_weekly
Primary symptoms: facial_spasms
Severity Scale: 3
Follow-up Plan: 2_weeks

PROFESSIONAL ANATOMICAL TREATMENT PLAN
Target Area: CORRUGATOR SUPERCILII
Analysis Type: hyperactivity
Primary Treatment: dermal fillers
CLINICAL ASSESSMENT:
The patient presents with symptoms affecting the corrugator supercilii region. Based on the hyperactivity analysis, the following treatment protocol is recommended.
Primary Symptoms: facial_spasms
Severity Rating: 3
TREATMENT PROTOCOL:
1. Initial Assessment
- Comprehensive evaluation of corrugator supercilii function and condition
- Baseline documentation with clinical photography
- Patient history and contraindication screening
2. Primary Treatment: dermal fillers
- Treatment Intensity: moderate
- Session Frequency: twice_weekly
- Progressive monitoring of muscle response and patient tolerance
- Adjustment of treatment parameters based on individual response
3. Monitoring and Follow-up
- Regular assessment of treatment efficacy
- Documentation of progressive improvements
- 2_weeks
PATIENT EDUCATION:
- Expected timeline for visible results: 7-14 days
- Potential side effects and their management
- Post-treatment care instructions
- Activity restrictions and recommendations
SAFETY CONSIDERATIONS:
- Patient should be advised of all potential risks and contraindications
- Informed consent must be obtained prior to treatment
- Emergency protocols should be reviewed with patient
- Contact information for post-treatment concerns provided
This treatment plan should be reviewed and adjusted based on individual patient response and clinical judgment.